

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/309445345>

# HOW TO USE CROP GROWTH MODEL WOFOST FOR FORECASTING GROWTH AND YIELD OF A CROP

Article in Journal of AgriSearch · March 2016

DOI: 10.21921/jas.v3i1.4107

CITATIONS  
0

6 authors, including:

 **Lal Singh**  
Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir  
**49 PUBLICATIONS 283 CITATIONS**  
[SEE PROFILE](#)

 **Raihana Kanth**  
Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir  
**144 PUBLICATIONS 768 CITATIONS**  
[SEE PROFILE](#)

READS  
1,470

 **Parmeet Singh**  
Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir  
**19 PUBLICATIONS 86 CITATIONS**  
[SEE PROFILE](#)

 **Purshotam Singh**  
Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir  
**61 PUBLICATIONS 257 CITATIONS**  
[SEE PROFILE](#)



## How to use crop growth model WOFOST for forecasting growth and yield of a crop

LAL SINGH\*, PARMEET SINGH, RAIHANA HABIB KANTH, PURUSHOTAM SINGH,  
SABIA AKHTER AND AMAL SAXENA

Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, J&K, India

### ABSTRACT

WOFOST version 7.1.3 is a computer model that simulates the growth and production of annual field crops. All the run options are operational through a graphical user interface named WOFOST Control Center version 1.8 (WCC). WCC facilitates selecting the production level, and input data sets on crop, soil, weather, crop calendar, hydrological field conditions, soil fertility parameters and the output options. The files with crop, soil and weather data are explained, as well as the run files and the output files. A general overview is given of the development and the applications of the model. Its underlying concepts are discussed briefly.

ARTICLE INFO	
Received on :	28.02.2016
Accepted on :	03.03.2016
Published online :	10.03.2016

**Keywords:** computer model, crop data simulation, crop production, production ecology, soil data, system analysis, weather data

### INTRODUCTION

WOFOST (WOrldFOodSTudies) is a simulation model for the quantitative analysis of the growth and production of annual field crops. It is a mechanistic model that explains crop growth on the basis of the underlying processes, such as photosynthesis, respiration and how these processes are influenced by environmental conditions. With WOFOST, you can calculate attainable crop production, biomass, water use, etc. for a location given knowledge about soil type, crop type, weather data and crop management factors (e.g. sowing date). WOFOST has been used by many researchers over the World and has been applied for many crops over a large range of climatic and management conditions. Moreover, WOFOST is implemented in the Crop Growth Monitoring System which is used operationally to monitor arable crops in Europe and to make crop yield forecasts for the current growing season. WOFOST originated in the framework of interdisciplinary studies on world food security and on the potential world food production by the Center for World Food Studies (CWFS) in cooperation with the Wageningen Agricultural University, Department of Theoretical Production Ecology (WAU-TPE) and the DLO-Center for Agrobiological Research and Soil Fertility (AB-DLO), Wageningen, the Netherlands. After cessation of CWFS in 1988, the DLO Winand Staring Centre (SC-DLO) has continued development of the model in co-operation with AB-DLO and WAU-TPE. Currently, the WOFOST model is maintained and further developed by Alterra in co-operation with the Plant Production Systems Group of Wageningen University (<http://www.pps.wur.nl/UK>) and the Agri4Cast unit of the Joint Research Centre in Italy (<http://mars.jrc.it/mars/About-us/AGRI4CAST>).

**Purpose of application:** Crop growth monitoring with agrometeorological data set. We can forecast the yield of different crops for any location well in advance with the help of

previous crop, soil and weather data, and expected weather data of the particular location. This may help in decision making of agricultural production system in time.

#### Input data requirement:

##### 1. Crop information like as :

- \*\* Crop identity : Rapeseed (*Brassica napus* L. ssp. *oleifera* (Metzg.)Sinsk.)
- \*\* CRPNAM : Brown sarson, Shalimar, Kashmir'
- \*\* Emergence
- TBASEM : Lower threshold temperature for emergence [cel]
- TEFFMX : Maximum effective temperature for emergence [cel]
- TSUMEM : Temperature sum from sowing to emergence [cel d]
- \*\* Phenology
- IDSL : Indicates whether pre-anthesis development depends on temperature (=0), daylength (=1), or both (=2)
- DLO : Optimum daylength for development [hr]
- DLC : Critical daylength (lower threshold) [hr]
- TSUM1 : Temperature sum from emergence to anthesis [cel d]
- TSUM2 : Temperature sum from anthesis to maturity [cel d]
- DTSMTB : Daily increase in temp. sum as function of av. temp. [cel; cel d]
- DVSEND : Development stage at harvest (= 2.0 at maturity [-])
- \*\* Initial
- TDWI : Initial total crop dry weight [ $\text{kg ha}^{-1}$ ]
- LAIEM : Leaf area index at emergence [ $\text{ha ha}^{-1}$ ]
- RGRLAI : Maximum relative increase in LAI

\* Corresponding author email:[drlalsingh@rediffmail.com](mailto:drlalsingh@rediffmail.com)

	[ha ha <sup>-1</sup> d <sup>-1</sup> ]		
** Green area			depletion [-]
SLATB	: Specific leaf area as a function of DVS [-; ha kg <sup>-1</sup> ]	IAIRDU	: Air ducts in roots present (=1) or not (=0)
SPA	: Specific pod area [ha kg <sup>-1</sup> ]	** Rooting	
SSATB	: Specific stem area [ha kg <sup>-1</sup> ] as function of DVS	RDI	: Initial rooting depth [cm]
SPAN	: Life span of leaves growing at 35 Celsius [d]	RRI	: Maximum daily increase in rooting depth [cm d <sup>-1</sup> ]
TBASE	: Lower threshold temp. for ageing of leaves [cel]	RDMCR	: Maximum rooting depth [cm]
** Assimilation		** Nutrients	
KDIFTB	: Extinction coefficient for diffuse visible light [-] as function of DVS	** Maximum and minimum concentrations of N, P, and K in storage organs and in vegetative organs [kg kg <sup>-1</sup> ]	
EFFTB	: Light-use efficiency single leaf [kg ha <sup>-1</sup> hr <sup>-1</sup> j <sup>-1</sup> m <sup>2</sup> s] as function of	YZERO	: Max. amount veg. organs at zero yield [kg ha <sup>-1</sup> ]
dailymean temperature		NFIX	: Fraction of N-uptake from biol. fixation [kg kg <sup>-1</sup> ]
AMAXTB	: Maximum leaf CO <sub>2</sub> assimilation-rate function of DVS [-; kg ha <sup>-1</sup> hr <sup>-1</sup> ]		
** Conversion of assimilates into biomass			
CVL	: Efficiency of conversion into leaves [kg kg <sup>-1</sup> ]	2. Soil information	
CVO	: Efficiency of conversion into storage org. [kg kg <sup>-1</sup> ]	SOLNAM	: 'EC3-medium fine' (It depends on your site information file name)
CVR	: Efficiency of conversion into roots [kg kg <sup>-1</sup> ]	** Physical soil characteristics	
CVS	: Efficiency of conversion into stems [kg kg <sup>-1</sup> ]	** Soil water retention	
** Maintenance respiration		SMTAB	: Vol. soil moisture content as function of pF [log (cm); cm <sup>3</sup> cm <sup>-3</sup> ]
Q10	: Relative increase in respiration rate per 10 Cel temp. incr. [-]	SMW	: Soil moisture content at wilting point [cm <sup>3</sup> /cm <sup>3</sup> ]
RML	: Rel. maintenance respiration rate leaves [kg CH <sub>2</sub> O kg <sup>-1</sup> d <sup>-1</sup> ]	SMFCF	: Soil moisture content at field capacity [cm <sup>3</sup> /cm <sup>3</sup> ]
RMO	: Rel. maint. resp. rate storage organ [kg CH <sub>2</sub> O kg <sup>-1</sup> d <sup>-1</sup> ]	SM0	: Soil moisture content at saturation [cm <sup>3</sup> /cm <sup>3</sup> ]
RMR	: Rel. maint. resp. rate roots [kg CH <sub>2</sub> O kg <sup>-1</sup> d <sup>-1</sup> ]	CRAIRC	: Critical soil air content for aeration [cm <sup>3</sup> /cm <sup>3</sup> ]
RMS	: Rel. maint. resp. rate stems [kg CH <sub>2</sub> O kg <sup>-1</sup> d <sup>-1</sup> ]	** Hydraulic conductivity	
RFSETB	: Red. factor for senescence as function of DVS[-;-]	CONTAB	: 10-log hydraulic conductivity as function of pF [log (cm); log (cm/day)]
** Partitioning		K0	: Hydraulic conductivity of saturated soil [cm day <sup>-1</sup> ]
FRTB	: Fraction of total dry matter to roots as a function of DVS[-; kg kg <sup>-1</sup> ]	SOPE	: Maximum percolation rate root zone [cm day <sup>-1</sup> ]
FLTB	: Fraction of above-gr. DM to leaves as a function of DVS[-; kg kg <sup>-1</sup> ]	KSUB	: Maximum percolation rate subsoil [cm day <sup>-1</sup> ]
FSTB	: Fraction of above-gr. DM to stems as a function of DVS[-; kg kg <sup>-1</sup> ]	** Soil workability parameters	
FOTB	: Fraction of above-gr. DM to stor. org. as a function of DVS[-; kg kg <sup>-1</sup> ]	SPADS	: 1 <sup>st</sup> topsoil seepage parameter deep seedbed
** Death rates		SPODS	: 2 <sup>nd</sup> topsoil seepage parameter deep seedbed
PERDL	: Max. rel. death rate of leaves due to water stress	SPASS	: 1 <sup>st</sup> topsoil seepage parameter shallow seedbed
RDRRTB	: Rel. death rate of stems as a function of DVS [-; kg kg <sup>-1</sup> d <sup>-1</sup> ]	SPOSS	: 2 <sup>nd</sup> topsoil seepage parameter shallow seedbed
RDRSTB	: Rel. death rate of roots as a function of DVS [-; kg kg <sup>-1</sup> d <sup>-1</sup> ]	DEFLIM	: Required moisture deficit deep seedbed
** Water use			
CFET	: Correction factor transpiration rate [-]		
DEPNR	: Crop group number for soil water		

### 3. Weather information




Daily weather data of solar radiation, minimum temperature, maximum temperature and rainfall is required to run the program. The WOFOST model support InfoCrop format weather files with a minor alteration. Therefore we convert our daily weather data from exel format to InfoCrop format by conversion in weather master of InfoCrop and copy that converted file pasting in WOFOST weather folder. After

doing required change save it. In addition to daily data of these parameters the below mentioned information's are also required.

Station name : Shalimar (It depends on your site information file name)

\* Longitude :74 Latitude:34 Altitude: 1587

Column	Daily Value	Units
* 1	Station number	
* 2	Year	
* 3	Day	
* 4	Irradiance	KJ m <sup>-2</sup>
* 5	Min Temperature	°C
* 6	Max Temperature	°C
* 7	Early Morning VP	kPa
* 8	Mean Wind Speed	m s <sup>-1</sup>
* 9	Precipitation	mm d <sup>-1</sup>



\*\*WCCDESCRIPTION=Shalimar, Kashmir

\*\*WCCFORMAT=2

These additional information should be pasted



\*\*WCCYEARNR=2014

in InfoCrop file to make it WOFOST weather file

How to convert weather data from excel format to InfoCrop format



Column	Daily Value	Units
1	Station number	
2	Year	
3	Day	
4	Irradiance	KJ m <sup>-2</sup>
5	Min Temperature	°C
6	Max Temperature	°C
7	Early Morning VP	kPa
8	Mean Wind Speed	m s <sup>-1</sup>
9	Precipitation	mm d <sup>-1</sup>



**WOFOST Control Center - Session: WCC**

**Select variables reruns**

**Detailed output - WCC**

**WOFOST Control Center - Session: WCC**

Results for 'POTENTIAL CROP PRODUCTION' in table format are shown for different time steps ('DAY') according to the value you specified in the 'Output interval in days' in the input tab 'General'. The results contain:

- 'IDSEM' : number of days since emergence (d);
  - 'DVS' : development stage of crop (-);
  - 'TSUM' : thermal time since emergence (Cd);
  - 'WLW' : dry weight of living leaves ( $\text{kg ha}^{-1}$ );
  - 'WST' : dry weight of living stems ( $\text{kg ha}^{-1}$ );
  - 'WSO' : dry weight of living storage organs ( $\text{kg ha}^{-1}$ );
  - 'TAGP' : total above ground production (dead and living plant organs) ( $\text{kg ha}^{-1}$ );
  - 'LAI' : leaf area index: (leaf area)/(soil area) ( $\text{ha ha}^{-1}$ );
  - 'TRA' : transpiration rate ( $\text{mm d}^{-1}$ );
  - 'GASS' : gross assimilation rate ( $\text{kg} (\text{CH}_2\text{O}) \text{ ha}^{-1} \text{ d}^{-1}$ );
  - 'MRES' : maintenance respiration rate ( $\text{kg} (\text{CH}_2\text{O}) \text{ ha}^{-1} \text{ d}^{-1}$ );
  - 'DMI' : rate of dry matter increase ( $\text{kg ha}^{-1} \text{ d}^{-1}$ ).
- A 'SUMMARY' of results is given for the potential production:
- 'HALT' : day number at harvest (day of the year);
  - 'ANTH' : duration of pre-anthesis phase (d);
  - 'TWRT' : total dry weight of roots (dead and living) ( $\text{kg ha}^{-1}$ );
  - 'TWLV' : total dry weight of leaves (dead and living) ( $\text{kg ha}^{-1}$ );
  - 'TWST' : total dry weight of stems (dead and living) ( $\text{kg ha}^{-1}$ );
  - 'TWSO' : total dry weight of storage organs (dead and

- 'TAGP' : living) ( $\text{kg ha}^{-1}$ );
- 'HINDEX' : total above ground production (dead and living plant organs) ( $\text{kg ha}^{-1}$ );
- 'HINDEX' : harvest index: weight of storage organs / weight of total above ground crop (-);
- 'TRANSP' : total transpiration (cm water);
- 'TRC' : transpiration coefficient rate (kg (water) / kg (dry matter));
- 'GASST' : total gross assimilation ( $\text{kg} (\text{CH}_2\text{O}) \text{ha}^{-1}$ );
- 'MREST' : total maintenance respiration ( $\text{kg} (\text{CH}_2\text{O}) \text{ha}^{-1}$ ).

The 'Graph' button enables you to draw graphs of specified output data of the potential crop growth (Fig.). These are the same output data as shown in the table of the result tab 'Potential'. There is also an option to present one graph showing the total above ground production (dry weight of living and dead plant organs) and its components: dry weight of living stems, dry weight of living stems and dry weight of living storage organs. The 'Graph' window offers you to load another output file ('Output file 2...'). With the buttons 'Y-axis 1' and 'Y-axis 2' you can control the position of the y-axis (left and right).



## SUMMARY POTENTIAL PRODUCTION

Summary results for potential production are given in table format. One line is given for each growing season and run. The following variables are found in the table:

SUMMARY POTENTIAL PRODUCTION											
YR	RUNNAM	SOW	EM	ANT	FLWR	DUR	HALT	PWT	TWLV	TWST	TWSO
1988	WCC1	99	0	1	169	169	228	223	1,271	2914	727
1988	WCC1	99	0	1	142	143	205	206	618	1,326	440
1989	WCC1	99	0	1	141	142	202	203	845	1,869	598

Means over 3 years:  
Standard deviation:  
Variation coefficients:

- 'YR' : simulation year;
- 'RUNNAM' : name of the simulation run;
- 'SOW' : sowing date (day in year) (in case of fixed emergence value is -99);
- '-->' : days between sowing and emergence (d) (in case of fixed emergence value is 0);
- 'EM' : emergence date (day in year);
- 'ANT' : duration of pre-anthesis phase (d);
- 'FLWR' : day of flowering (day in year);
- 'DUR' : duration of simulation period (d);
- 'HALT' : day number at harvest (day in year);

- 'TWRT' : total dry weight of roots (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWLV' : total dry weight of leaves (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWST' : total dry weight of stems (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWSO' : total dry weight of storage organs (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TAGP' : total above ground production (dead and living plant organs) ( $\text{kg ha}^{-1}$ );
- 'LAIM' : maximum leaf area index ( $\text{ha ha}^{-1}$ );
- 'HINDEX' : harvest index: weight of storage organs / weight of total above ground crop (-);
- 'TRC' : transpiration coefficient rate (kg (water) / kg (dry matter));
- 'GASST' : total gross assimilation ( $\text{kg} (\text{CH}_2\text{O}) \text{ha}^{-1}$ );
- 'MREST' : total maintenance respiration ( $\text{kg} (\text{CH}_2\text{O}) \text{ha}^{-1}$ );
- 'TRANSP' : total transpiration (cm water);
- 'EVTSOL' : total evaporation from soil surface (cm water).

A table is filled with means ('Means over x years'), standard deviation ('Standard deviation'), and variation coefficients ('Variation coefficients') when a run for more than two weather years is made. These statistics are given for:

- 'DUR' : duration of simulation period (d);
- 'TWLV' : total dry weight of leaves (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWST' : total dry weight of stems (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWSO' : total dry weight of storage organs (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TAGP' : total above ground production (dead and living plant organs) ( $\text{kg ha}^{-1}$ );
- 'LAIM' : maximum leaf area index ( $\text{ha ha}^{-1}$ );
- 'HINDEX' : harvest index: weight of storage organs / weight of total above ground crop (-);
- 'TRC' : transpiration coefficient rate (kg (water) / kg (dry matter));
- 'TRANSP' : total transpiration (cm water);
- 'EVTSOL' : total evaporation from soil surface (cm water).

## SUMMARY OUTPUT

The different fields in the table are (first five field are repeated in second table!):

- 'PS' : production level (PP=potential; WP=water-limited);
- 'RUNNAM' : name of the simulation run;
- 'IZT' : influence of ground water (0 = no influence; 1 = influence);
- 'SOW' : sowing date (day in year) (in case of fixed emergence value is -99);
- 'EM' : emergence date (day in year);
- 'DUR' : duration of simulation period (d);
- 'TWLV' : total dry weight of leaves (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWST' : total dry weight of stems (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TWSO' : total dry weight of storage organs (dead and

- 'var%'	: living) ( $\text{kg ha}^{-1}$ ); variation coefficient of total dry weight of storage organs (dead and living) ( $\text{kg ha}^{-1}$ );
- 'TAGP'	: total above ground production (dead and living plant organs) ( $\text{kg ha}^{-1}$ );
- 'var%'	: variation coefficient of total above ground production (dead and living plant organs) ( $\text{kg ha}^{-1}$ );
- 'LAIM'	: maximum leaf area index ( $\text{ha ha}^{-1}$ );
- 'HINDEX'	: harvest index: weight of storage organs / weight of total above ground crop (-);
- 'RYLD'	: relative yield: water-limited yield / potential yield (%);
- 'RAGP'	: relative total above ground production: water-limited production / potential production (%);
- 'TRC'	: transpiration coefficient rate ( $\text{kg (water)} / \text{kg (dry matter)}$ );
- 'RDMSOL'	: maximum rooting depth allowed by soil (cm).

## CONCLUSION

WOFOST is a tool for the quantitative analysis of the growth and production of annual field crops. As with all tools, you should know what you could do with it and what not. Like all mathematical models of agricultural production, WOFOST is a simplification of reality. In practice, crop yield is a result of the interaction of ecological, technological and socio-economic factors. In WOFOST, only ecological factors are considered under the assumption that optimum

management practices are applied.

## Limitations of WOFOST

It should be stressed again that WOFOST is a model, hence a simplification of reality. The user always has to be cautious when drawing conclusions from the simulation results. Keep in mind that the quality of the model results cannot surpass the quality of the input data. A model merely elucidates the consequences of the user's opinions and data (Driessen, 1986). Therefore, the careful selection of the input data is of utmost importance. As a general rule, you should not simulate crop growth without experimentation. Experimentation is needed to obtain specific parameters and to calibrate and verify the model results.

In the validation process, crop growth as observed in reality can be compared with the simulation results. This provides an impression of the adequacy of WOFOST's predictions. When differences occur, you may need to adapt values of model-parameters (calibration). Then, the model should again be checked, against an independent set of observations. A problem in this respect is that there are an enormous number of parameters while only a few can be validated at a time. One should also realize that although calibration may improve the model results for a specific purpose, it can diminish its general applicability.

A weakness of WOFOST and all other crop growth simulation models is that some parameters are fixed whereas in practice they are known to vary, i.e. those concerning the relation between development stage and partitioning (Passioura, 1996). This problem arises when the processes at a lower integration level are insufficiently known.

Passioura JB. 1996. *Simulation Models: Science, Snake oil, Education, or Engineering?* Agronomy Journal.88:690-4.

<http://www.pps.wur.nl/UK>

<http://mars.jrc.it/mars/About-us/AGRI4CAST>

## REFERENCE

- Driessen PM. 1986. *The collection and treatment of basic data. Introduction*. In: Van Keulen and Wolf, 1986. 203-207.

## Citation:

Singh L, Singh P, Kanth R H, Singh P, Akhter S and Saxena A. 2016. How to use crop growth model WOFOST for forecasting growth and yield of a crop. *Journal of AgriSearch*3 (1):63-68