

pfsl22.tex

Lecture 22. 28.11.2012

This counts the Poisson points in B – and is a Poisson process with rate (parameter) $\nu(B)$. All this reverses: starting with an $e = (e(t) : t \geq 0)$ whose counting processes over Borel sets B are Poisson $P(\nu(B))$, then – as no point can contribute to more than one count over disjoint sets – disjoint counting processes never jump together, so are independent by above, and $\phi := \sum_{t \geq 0} \delta_{(e(t), t)}$ is a Poisson measure with intensity $\mu = \nu \times dt$.

Lévy Processes; Lévy-Khintchine Formula; Lévy-Itô decomposition.

We can now sketch the close link between the general Lévy process on the one hand and the general infinitely-divisible law given by the Lévy-Khintchine formula (LK) on the other.

First, if $X = (X_t)$ is Lévy, the law of each X_1 is infinitely divisible, so

$$E \exp\{iuX_1\} = \exp\{-\Psi(u)\} \quad (u \in \mathbf{R})$$

with Ψ a Lévy exponent as in (LK). Similarly,

$$E \exp\{iuX_t\} = \exp\{-t\Psi(u)\} \quad (u \in \mathbf{R}),$$

for rational t at first and general t by approximation and càdlàg paths. Then Ψ is called the *Lévy exponent*, or *characteristic exponent*, of the Lévy process X . Conversely, given a Lévy exponent $\Psi(u)$ as in (LK), construct a Brownian motion, and an independent Ppp $\Delta = (\Delta_t : t \geq 0)$ with characteristic measure μ , the Lévy measure in (LK). Then $X_1(t) := at + \sigma B_t$ has CF

$$E \exp\{iuX_1(t)\} = \exp\{-t\Psi_1(t)\} = \exp\left\{-t(iau + \frac{1}{2}\sigma^2 u^2)\right\},$$

giving the non-integral terms in (LK). For the ‘large’ jumps of Δ , write

$$\Delta_t^{(2)} := \Delta_t \text{ if } |\Delta_t| \geq 1, \quad 0 \text{ else.}$$

Then $\Delta^{(2)}$ is a Poisson point process with characteristic measure $\mu^{(2)}(dx) := I(|x| \geq 1)\mu(dx)$. Since $\int \min(1, |x|^2)\mu(dx) < \infty$, $\mu^{(2)}$ has finite mass, so $\Delta^{(2)}$, a $Ppp(\mu^{(2)})$, is discrete and its counting process

$$X_t^{(2)} := \sum_{s \leq t} \Delta_s^{(2)} \quad (t \geq 0)$$

is compound Poisson, with Lévy exponent

$$\Psi^{(2)}(u) = \int (1 - e^{iux}) I(|x| \geq 1) \mu(dx) = \int (1 - e^{iux}) \mu^{(2)}(dx).$$

There remain the ‘small jumps’,

$$\Delta_t^{(3)} := \Delta_t \text{ if } |\Delta_t| < 1, \quad 0 \text{ else,}$$

a $Ppp(\mu^{(3)})$, where $\mu^{(3)}(dx) = I(|x| < 1) \mu(dx)$, and independent of $\Delta^{(2)}$ because $\Delta^{(2)}, \Delta^{(3)}$ are Poisson point processes that never jump together. For each $\epsilon > 0$, the ‘compensated sum of jumps’

$$X_t^{(\epsilon,3)} := \sum_{s \leq t} I(\epsilon < |\Delta_s| < 1) \Delta_s - t \int x I(\epsilon < |x| < 1) \mu(dx) \quad (t \geq 0)$$

is a Lévy process with Lévy exponent

$$\Psi^{(\epsilon,3)}(u) = \int (1 - e^{iux} + iux) I(\epsilon < |x| < 1) \mu(dx).$$

Use of a suitable maximal inequality allows passage to the limit $\epsilon \downarrow 0$ (going from finite to possibly countably infinite sums of jumps): $X_t^{(\epsilon,3)} \rightarrow X_t^{(3)}$, a Lévy process with Lévy exponent

$$\Psi^{(3)}(u) = \int (1 - e^{iux} + iux) I(|x| < 1) \mu(dx),$$

independent of $X^{(2)}$ and with càdlàg paths. Combining:

Theorem (Lévy-Itô decomposition). For a Lévy exponent

$$\Psi(u) = iau + \frac{1}{2} \sigma^2 u^2 + \int (1 - e^{iux} + iux I(|x| < 1)) \mu(dx),$$

the construction above yields a Lévy process

$$X = X^{(1)} + X^{(2)} + X^{(3)}$$

with Lévy exponent $\Psi = \Psi^{(1)} + \Psi^{(2)} + \Psi^{(3)}$. Here the $X^{(i)}$ are independent Lévy processes, with Lévy exponents $\Psi^{(i)}$; $X^{(1)}$ is Gaussian, $X^{(2)}$ is a compound Poisson process with jumps of modulus ≥ 1 ; $X^{(3)}$ is a compensated sum of jumps of modulus < 1 . The jump process $\Delta X = (\Delta X_t : t \geq 0)$ is a

$Ppp(\mu)$, and similarly $\Delta X^{(i)}$ is a $Ppp(\mu^{(i)})$ for $i = 2, 3$.

Stable processes. A stable process has (to within location and scale) a Lévy exponent involving two parameters, $\alpha \in (0, 2]$, called the *index*, and $\beta \in [-1, 1]$, called the *skewness parameter*:

$$\Psi(u) = |u|^\alpha (1 - i\beta(\operatorname{sgn} u) \tan(\frac{1}{2}\pi\alpha)) \quad (\alpha \neq 1), \quad |u|(1 + i\beta(\operatorname{sgn} u) \frac{2}{\pi} \log |u|) \quad (\alpha = 1)$$

(for $\alpha = 2$, β drops out as $\tan \pi = 0$, so $\Psi(u) = u^2$, giving the normal (Gaussian) distribution). The case $\alpha = 1$ is the *Cauchy* case; the asymmetric Cauchy case $\alpha = 1, \beta \neq 0$ is awkward, and we do not consider it further.

The Lévy measure μ in the stable case is absolutely continuous, with density ν , $\mu(dx) = \nu(x)dx$, where

$$\nu(x) = c_+/x^{1+\alpha} \quad (x > 0), \quad c_-/|x|^{1+\alpha} \quad (x < 0) \quad (c_\pm \geq 0, c_+ + c_- > 0).$$

Here

$$\beta = (c_+ - c_-)/(c_+ + c_-).$$

The calculations are simpler in the symmetric case, $c_+ = c_- = c$ say. Then

$$\Psi'(u) = 2cu^{\alpha-1}I \quad (u > 0), \quad I := \int_0^\infty v^{-\alpha} \sin v dv.$$

So $\Psi(u) = 2cIu^\alpha/\alpha$ for $u > 0$, and similarly for $u < 0$: $\Psi(u) = |u|^\alpha .2cI/a$. But (see e.g. M2PM3 L30 on my website: there $t = 1 - \alpha \in (0, 1)$, but we can extend by analytic continuation to $-1 < t < 1$, $\alpha \in (0, 2)$) $I = \Gamma(1 - \alpha) \cos(\frac{1}{2}\pi\alpha)$ (here $\alpha \neq 1$: $\Gamma(z)$ has a pole at $z = 0$, and the Cauchy case $\alpha = 1$ is dealt with separately above). So one choice is $c := \sigma/(2I)$, and then $\Psi(u) = |u|^\alpha$.

Example. The symmetric stable law with $\alpha = 3/2$ is called the *Holtmark distribution*, proposed by the Danish physicist J. Holtmark in 1919 as a model for the distribution of galaxies in space (here $3/2$ comes from the 3 dimensions of space and the 2 in Newton's Inverse Square Law of Gravity). Since $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ and $\Gamma(1 + x) = x\Gamma(x)$, $\Gamma(1 - \alpha) \cos(\frac{1}{2}\pi\alpha)$ here is $\int_0^\infty v^{3/2} \sin v dv = \Gamma(-\frac{1}{2}) \cos(3\pi/4) = (-2\sqrt{\pi}).(-1/\sqrt{2}) = \sqrt{2\pi}$.

Subordinators.

We resort to complex numbers in the CF $\phi(u) = E(e^{iuX})$ because this always exists – for all real u – unlike the ostensibly simpler moment-generating

function (MGF) $M(u) := E(e^{uX})$, which may well diverge for some real u . However, if the random variable X is *non-negative*, then for $s \geq 0$ the *Laplace-Stieltjes transform* (LST)

$$\psi(s) := E[e^{-sX}] \leq E(1) = 1$$

always exists. For $X \geq 0$ we have both the CF and the LST to hand, but the LST is usually simpler to handle. We can pass from CF to LST formally by taking $u = is$, and this can be justified by analytic continuation.

Some Lévy processes X have increasing (i.e. non-decreasing) sample paths; these are called *subordinators*. From the construction above, subordinators can have no negative jumps, so μ has support in $(0, \infty)$ and no mass on $(-\infty, 0)$. Because increasing functions have FV, one must have paths of (locally) finite variation, the condition for which can be shown to be

$$\int \min(1, |x|) \mu(dx) < \infty.$$

Thus the Lévy exponent must be of the form

$$\Psi(u) = -idu + \int_0^\infty (1 - e^{iux}) \mu(dx),$$

with $d \geq 0$. It is more convenient to use the Laplace exponent $\Phi(s) = \Psi(is)$:

$$E(\exp\{-sX_t\}) = \exp\{-t\Phi(s)\} \quad (s \geq 0), \quad \Phi(s) = ds + \int_0^\infty (1 - e^{-sx}) \mu(dx).$$

Example. The Stable Subordinator. Here $d = 0$, $\Phi(s) = s^\alpha$, ($0 < \alpha < 1$),

$$\mu(dx) = dx / (\Gamma(1 - \alpha)x^{\alpha-1})$$

(in one normalization, the one convenient here). The special case $\alpha = 1/2$ is particularly important: this arises as the first-passage time of Brownian motion over positive levels, and gives rise to the *Lévy density* (see Problems).

Classification.

IV (Infinite Variation). The sample paths have infinite variation on finite time-intervals, a.s. This occurs iff

$$\sigma > 0 \quad \text{or} \quad \int \min(1, |x|) \mu(dx) = \infty.$$