

INV-202 (Human Trial)

inverse agonist

SKYE-(Nimacimab)

Preferred term, n (%)	INV-202 (N = 20)	Placebo (N = 17)	Total (N = 37)
At least one TEAE	16 (80)	8 (47)	24 (65)
Nausea	7 (35)	1 (6)	8 (22)
Decreased appetite ^a	7 (35)	1 (6)	8 (22)
Diarrhoea ^b	5 (25)	0	5 (14)
Frequent urination ^c	4 (20)	1 (6)	5 (14)
Hot flush	3 (15)	1 (6)	4 (11)
Hiccups	3 (15)	0	3 (8)
Nervous system disorders SOC	8 (40)	3 (18)	11 (30)
Headache	2 (10)	2 (12)	4 (11)
Dizziness	2 (10)	0	2 (5)
Presyncope	2 (10)	0	2 (5)
Taste disorder	2 (10)	0	2 (5)
Dysgeusia	0	1 (6)	1 (3)
Loss of consciousness	1 (5)	0	1 (3)
Syncope	1 (5)	0	1 (3)
Psychiatric disorders SOC	4 (20)	1 (6)	5 (14)
Depressed mood	2 (10)	0	2 (5)
Insomnia	2 (10)	0	2 (5)
Irritability	1 (5)	0	1 (3)
Nervousness	0	1 (6)	1 (3)
Sleep disorder	1 (5)	0	0

Abbreviations: SOC, system organ class; TEAE, treatment-emergent adverse event.

^a Includes early satiety.

^b Includes faeces soft/loose.

^c Includes pollakiuria and micturition urgency.

Nimacimab was observed in clinical studies, a safety issue assessment from the disease ("NAFLD") patients, no early termination events, no early termination concern occurring in an exploratory biomarker study. Patients receiving nimacimab v placebo had similar pharmacokinetic assessments over 4 weeks, potentially allowing for once-a-week subcutaneous dosing.

The promising PK data suggest that nimacimab can be dosed at a lower dose than competitive advantage receptor agonists or eCB agonists, supporting a tolerability profile.

Nimacimab is considered a CB1 receptor antagonist across multiple studies, with no direct binding and inhibition of CB1 from allosteric modulation.

Nimacimab effectively inhibits CB1 receptor signaling in clinical studies, is devoid of the CB1 receptor agonist effects. Worldwide rights to nimacimab are held by SKY, including claims to cannabinoid receptor antagonists.

In summary, leptin and CB1 receptor inhibiting lipogenesis mechanisms and obesity, adiposity and contrast.


Inverse agonist

CRBP-CRB 913

erved to have very limited accumulation in the CNS in pre-clinical
in earlier generations of CB1 inhibitors. Safety and tolerability
completed Phase 1b study of nimaclamab in non-alcoholic fatty liver
patients with diabetes or prediabetes demonstrated no serious adverse
nations of treatment due to adverse events, and no adverse events of
dose-dependent manner. Encouraging trends were observed in
s including cholesterol, liver enzymes and liver function in patients
versus placebo after the four-week dosing period. Moreover,
ssment of nimaclamab highlighted a half-life of approximately three
wing for monthly dosing, which would offer a competitive advantage
cutaneous dosing of current peptidic GLP-1 receptor agonists.

a of approximately 21 days from the Phase 1 study suggests that
sed once-a-month subcutaneously, which would potentially offer a
e over once-a-week subcutaneous dosing of current peptidic GLP-1
ven orally dosed GLP-1 receptor agonists due to their less desirable

CRB-913: Marked ↓ Brain and ↑
Lean and Obese Mice



Brain	
single acute dose	CRB-913 (ng/g)
10 mg/Kg	31.4

stently described as a "negative allosteric modulator" of the CB1
ltiple sources ^{1 2 3 4 5}. An inverse agonist would imply
nhibition of the receptor's constitutive activity, which is different
ulation.

nhibits CB1 signaling and, based on preclinical and early clinical
CNS liabilities typically seen by small molecule drugs that
because it does not cross the blood-brain barrier. Skye owns the
acimab, with patents issued in the U.S. and other territories
nabinoid 1 receptor antibodies with inverse agonist function.


exerts direct catabolic effects on adipose tissue by promoting lipolysis
sis/adipogenesis, and inducing thermogenesis. Targeting these
vercoming leptin resistance could be an effective strategy to reduce
rol excess leptin levels associated with obesity. ^{1 2 3 4}

inverse agonist

Rimnabant- Inverse Agonist

Phase 1

Peripheral Exposure Vs. Rimonabant in Bot



events, which occurred more frequently with 20 mg rimonab
ressive disorders [1.9 vs. 0.8%], nausea [1.4 vs. 0.1%], mo
vs. 0.6%], and anxiety [1.0 vs. 0.3%]). A thorough review
its was performed.

receptor, rimonabant was s
antagonist of the μ -opioid r

ase 3

concentration (ng/g)		
CRB-913 (lean mice)	Monlunabant (lean mice)	Rimonabant (lean mice)
26*	319**	561*
1:12		
1:21		

comparable versus placebo (most commonly, mood alterations with depressive symptoms) and no significant psychiatric and neurological adverse events.

Subsequently also found to act as an NMDA receptor.

Taranabant

Inverse agonist

Taranabant

- **CNS penetration:** Taranabant is designed to be a CNS-acting agent, but it is not intended to reduce the risk of CNS-related adverse events. Since it is not a CNS-acting agent, its penetration is thought to be lower than that of Rimonabant.
- **Bioavailability:** Taranabant has moderate oral bioavailability of around 30-40% in humans, which may contribute to its efficacy.
- **Brain-to-plasma ratio:** Taranabant has a low brain-to-plasma ratio compared to Rimonabant, indicating that it may not accumulate in the brain.

	Placebo N = 417	Taranabant 2 mg N = 414
One or more clinical AE	361 (86.6%)	353 (85.3%)
Drug-related ^c clinical AE	152 (36.5%)	185 (44.7%)
Serious clinical AE	25 (6.0%)	33 (8.0%)
Drug-related ^c serious clinical AE	0	1
Discontinued due to clinical AE	40 (9.6%)	53 (12.8%)
Discontinued due to drug-related ^c clinical AE	29 (7.0%)	42 (10.2%)
Discontinued due to serious clinical AE	4 (1.0%)	3 (0.7%)
Discontinued due to drug-related ^c serious clinical AE	0	1
Deaths ^d	1 (0.2%)	1 (0.2%)
One or more laboratory AE	16 (3.9%)	12 (2.9%)
Drug-related ^c laboratory AE	4 (1.0%)	3 (0.7%)
Serious laboratory AE	0	1
Discontinued due to laboratory AE	2 (0.5%)	1 (0.2%)
Discontinued due to drug-related ^c laboratory AE	2 (0.5%)	1 (0.2%)

Table 5 Number (%) of patients with pre-specified clinical adverse experiences

Clinical adverse event	Placebo N = 417	Taranabant 2 mg N = 414
Gastrointestinal disorders SOC	119 (28.5%)	173 (41.8%)**
Diarrhea	30 (7.2%)	51 (12.3%)*
Frequent bowel movements	2 (0.5%)	6 (1.4%)
Nausea	27 (6.5%)	69 (16.7%)**
Vomiting	14 (3.4%)	23 (5.6%)
Expanded nervous system disorders SOC ^b	105 (25.2%)	94 (22.7%)
Dizziness grouping ^c	24 (5.8%)	23 (5.6%)
Cognition grouping ^d	6 (1.4%)	5 (1.2%)
Sedation grouping ^e	4 (1.0%)	7 (1.7%)
Epilepsy, convulsion	0	0
Stroke grouping ^f	0	0
Syncope grouping ^g	4 (1.0%)	3 (0.7%)
Vertigo grouping ^h	0	2 (0.5%)
Sensory disorder grouping ⁱ	11 (2.6%)	4 (1.0%)
Motor disorder grouping ^j	0	3 (0.7%)
Vision grouping ^k	4 (1.0%)	0
Headache grouping ^l	59 (14.1%)	43 (10.4%)
Psychiatric disorders SOC plus irritability	85 (20.4%)	117 (28.3%)**
Altered affect grouping ^m	11 (2.6%)	19 (4.6%)
Depression grouping ⁿ	27 (6.5%)	37 (8.9%)
Depression, major depression	12 (2.9%)	13 (3.1%)
Depressed mood, depressive symptom	15 (3.6%)	24 (5.8%)
Irritability	7 (1.7%)	22 (5.3%)**
Nervousness	3 (0.7%)	3 (0.7%)
Anxiety grouping ^o	14 (3.4%)	20 (4.8%)
Anxiety	14 (3.4%)	16 (3.9%)
Anxiety disorder	0	0

Anxiety disorder	0	0
Panic disorder, panic attack	0	4 (1.0%)
Insomnia grouping ^p	25 (6.0%)	27 (6.5%)
Aggression, anger	3 (0.7%)	9 (2.2%)
Disorientation, confusional state	0	1 (0.2%)
Suicidal grouping ^q	0	0
Suicidal ideation ^r	0	0
Skin and Subcutaneous disorders SOC minus	62 (14.9%)	71 (17.1%)
hypoesthesia facial		
Hyperhidrosis	2 (0.5%)	8 (1.9%)
Pruritus grouping ^s	12 (2.9%)	19 (4.6%)
Vascular disorders SOC	22 (5.3%)	28 (6.8%)
Flushing, hot flush	4 (1.0%)	11 (2.7%)

Table 4 Clinical adverse experiences classified by MedDRA system organ class

	Placebo N = 417	Taranaba N = 4
<i>SOCs and specific clinical AEs with incidences $\geq 5\%$^c</i>		
Eye disorders	23 (5.5%)	14 (3.4%)
Gastrointestinal disorders	119 (28.5%)	173 (41.5%)
Diarrhea	30 (7.2%)	51 (12.0%)
Nausea	27 (6.5%)	69 (16.2%)
Vomiting	14 (3.4%)	23 (5.5%)
General disorders and administrative site conditions	58 (13.9%)	65 (15.2%)
Irritability	7 (1.7%)	22 (5.2%)
Infections and Infestations	225 (54.0%)	206 (48.5%)
Influenza	19 (4.6%)	34 (8.0%)
Nasopharyngitis	67 (16.1%)	49 (11.5%)
Sinusitis	23 (5.5%)	22 (5.2%)
Upper respiratory tract infection	50 (12.0%)	59 (14.0%)
Injury, poisoning and procedural complications	66 (15.8%)	63 (15.0%)
Metabolism and nutrition disorders	15 (3.6%)	23 (5.5%)
Musculoskeletal and connective tissue disorders	122 (29.3%)	117 (28.0%)
Arthralgia	29 (6.7%)	25 (6.0%)
Back pain	35 (8.4%)	30 (7.2%)
Pain in extremity	23 (5.5%)	20 (4.8%)
Nervous system disorders	102 (24.5%)	94 (22.2%)
Dizziness	24 (5.8%)	23 (5.5%)
Headache	52 (12.5%)	38 (9.0%)
Psychiatric disorders	80 (19.2%)	108 (25.8%)
Anxiety	14 (3.4%)	16 (3.8%)
Depressed mood	12 (2.9%)	21 (5.0%)
Insomnia	25 (6.0%)	27 (6.4%)
Renal and urinary disorders	11 (2.6%)	11 (2.6%)
Reproductive system and breast disorders	22 (5.3%)	29 (6.9%)
Respiratory, thoracic and mediastinal disorders	56 (13.4%)	62 (14.8%)
Skin and subcutaneous tissue disorders	62 (14.9%)	72 (17.2%)
Pruritus	12 (2.9%)	19 (4.5%)
Vascular disorders	22 (5.3%)	28 (6.7%)
Hypertension	15 (3.6%)	9 (2.1%)
<i>Drug-related^d SOC and clinical AEs with incidences $\geq 2\%$^e</i>		
Gastrointestinal disorders	54 (12.9%)	93 (22.2%)
Diarrhea	11 (2.6%)	20 (4.8%)
Frequent bowel movements	1 (0.2%)	3 (0.7%)
Nausea	16 (3.8%)	49 (11.5%)
Vomiting	3 (0.7%)	5 (1.2%)
General Disorders and administrative site conditions	22 (5.3%)	28 (6.7%)
Fatigue	6 (1.4%)	7 (1.7%)
Irritability	7 (1.7%)	16 (3.8%)
Metabolism and nutrition disorders	2 (0.5%)	10 (2.4%)
Nervous system disorders	38 (9.1%)	34 (8.0%)
Dizziness	11 (2.6%)	10 (2.4%)
Headache	17 (4.1%)	6 (1.4%)
Nervous system disorders	102 (24.5%)	94 (22.7%)
Dizziness	24 (5.8%)	23 (5.6%)
Headache	52 (12.5%)	38 (9.2%)
Psychiatric disorders	80 (19.2%)	108 (26.1%)
Anxiety	14 (3.4%)	16 (3.9%)
Depressed mood	12 (2.9%)	21 (5.1%)
Insomnia	25 (6.0%)	27 (6.5%)

to have limited CNS penetration, which is
 d side effects. While it can still enter the
 r than that of Rimonabant.

oral bioavailability, with a reported absolute
 s. This is lower than Rimonabant's
 reduced CNS penetration.

lower brain-to-plasma ratio compared to
 imulate as much in the brain.

Rimonabant N = 414	Taranabant 4 mg N = 415	Taranabant 6 mg (fixed and titrated) ^b N = 1256	Placebo N = 244	Taranabant 2 mg N = 264	Taranabant 4 mg N = 260
85.3%)	377 (90.8%)	1110 (88.4%)	192 (78.7%)	189 (71.6%)	193 (74.2%)
44.7%)	236 (56.9%)	689 (54.9%)	18 (7.4%)	25 (9.5%)	38 (14.6%)
8.0%)	29 (7.0%)	91 (7.2%)	11 (4.5%)	16 (6.1%)	17 (6.5%)
0	1 (0.2%)	11 (0.9%)	0	1 (0.4%)	0
12.8%)	82 (19.8%)	263 (20.9%)	3 (1.2%)	11 (4.2%)	11 (4.2%)
10.1%)	66 (15.9%)	216 (17.2%)	2 (0.8%)	9 (3.4%)	7 (2.7%)
0.7%)	6 (1.4%)	18 (1.4%)	0	1 (0.4%)	2 (0.8%)
0	1 (0.2%)	10 (0.8%)	0	1 (0.4%)	0
0.2%)	0	0	0	0	0
2.9%)	16 (3.9%)	51 (4.1%)	8 (3.3%)	7 (2.7%)	11 (4.2%)
0.8%)	5 (1.2%)	20 (1.6%)	1 (0.4%)	2 (0.8%)	2 (0.8%)
0	0	0	0	0	0
0	1 (0.2%)	0	0	1 (0.4%)	0
0	1 (0.2%)	0	0	0	0

and groupings of interest at year 1 (week 52) and year 2 (week 104)

Group	Year 1		Year 2 ^a		
	Taranabant 4 mg N = 415	Taranabant 6 mg (fixed and titrated) N = 1256	Placebo N = 254	Taranabant 2 mg N = 270	Taranabant 4 mg N = 264
**	193 (46.5%)*	578 (46.0%)*	62 (24.4%)	45 (16.7%)*	58 (22.0%)
	57 (13.7%)*	174 (13.9%)*	18 (7.1%)	9 (3.3%)	14 (5.3%)
	12 (2.9%)*	20 (1.6%)	0	0	1 (0.4%)
**	89 (21.4%)*	301 (24.0%)*	5 (2.0%)	6 (2.2%)	12 (4.5%)
	35 (8.4%)*	91 (7.3%)*	3 (1.2%)	4 (1.5%)	6 (2.3%)
	138 (33.3%)*	358 (28.5%)	30 (11.8%)	35 (13.0%)	45 (17.0%)
	43 (10.4%)*	111 (8.8%)*	8 (3.1%)	10 (3.7%)	8 (3.0%)
	9 (2.2%)	23 (1.8%)	1 (0.4%)	1 (0.4%)	4 (1.5%)
	12 (2.9%)*	27 (2.2%)	0	2 (0.7%)	1 (0.4%)
	0	1 (0.1%)	0	0	0
	1 (0.2%)	7 (0.6%)	0	0	1 (0.4%)
	14 (3.4%)*	26 (2.1%)	0	1 (0.4%)	2 (0.8%)
	0	3 (0.2%)	1 (0.4%)	0	0
	20 (4.8%)	53 (4.2%)	3 (1.2%)	5 (1.9%)	9 (3.4%)
	3 (0.7%)	12 (1.0%)*	2 (0.8%)	0	0
	0	4 (0.3%)	1 (0.4%)	1 (0.4%)	0
	57 (13.7%)	137 (10.9%)	17 (6.7%)	16 (5.9%)	20 (7.6%)
*	167 (40.2%)*	478 (38.1%)*	17 (6.7%)	23 (8.5%)	27 (10.2%)
	28 (6.7%)*	91 (7.3%)*	0	0	2 (0.8%)
	44 (10.6%)*	141 (11.2%)*	7 (2.8%)	8 (3.0%)	9 (3.4%)
	20 (4.8%)	57 (4.5%)	2 (0.8%)	2 (0.7%)	5 (1.9%)
	27 (6.5%)	84 (6.7%)*	4 (1.4%)	6 (2.2%)	3 (1.1%)
	36 (8.7%)*	85 (6.8%)*	1 (0.4%)	1 (0.4%)	3 (1.1%)
	8 (1.9%)	24 (1.9%)	0	2 (0.7%)	0
	46 (11.1%)*	117 (9.3%)*	3 (1.2%)	2 (0.7%)	8 (3.0%)
	41 (9.9%)*	105 (8.4%)*	3 (1.2%)	2 (0.7%)	8 (3.0%)
	2 (0.5%)	4 (0.3%)	0	0	0

2 (0.5%)	7 (0.5%)	0	0	0
4 (1.0%)	11 (0.9%)	0	0	0
40 (9.6%)	93 (7.4%)	5 (2.0%)	4 (1.5%)	12 (4.5%)
7 (1.7%)	22 (1.8%)	0	1 (0.4%)	4 (1.5%)
0	5 (0.4%)	0	0	0
0	3 (0.2%)	0	0	1 (0.4%)
0	3 (0.2%)	0	0	1 (0.4%)
72 (17.3%)	268 (21.3%)*	17 (6.7%)	17 (6.3%)	31 (11.7%)*
9 (2.2%)*	46 (3.8%)*	1 (0.4%)	0	1 (0.4%)
27 (6.5%)*	105 (8.4%)*	1 (0.4%)	3 (1.1%)	5 (1.9%)
49 (11.8%)*	129 (10.3%)*	16 (6.3%)	7 (2.6%)	19 (7.2%)
19 (4.6%)*	56 (4.5%)*	5 (2.0%)	1 (0.4%)	4 (1.5%)

sses

Year 1			Year 2 ^a		
int 2 mg 414	Taranabant 4 mg N = 415	Taranabant 6 mg (fixed and titrated) ^b N = 1256	Placebo N = 244	Taranabant 2 mg N = 264	Taranabant 4 mg N = 260
4%)	8 (1.9%)	57 (4.5%)	7 (2.9%)	12 (4.5%)	13 (5.0%)
1.8%)	194 (46.7%)	578 (46.0%)	52 (21.3%)	43 (16.3%)	54 (20.8%)
2.3%)	57 (13.7%)	174 (13.9%)	16 (6.6%)	9 (3.4%)	15 (5.8%)
5.7%)	89 (21.4%)	301 (24.0%)	5 (2.0%)	6 (2.3%)	11 (4.2%)
6%)	35 (8.4%)	91 (7.2%)	3 (1.2%)	3 (1.1%)	5 (1.9%)
5.7%)	90 (21.7%)	242 (19.3%)	20 (8.2%)	20 (7.6%)	23 (8.8%)
3%)	36 (8.7%)	85 (6.8%)	1 (0.4%)	0	3 (1.2%)
9.8%)	221 (53.3%)	623 (49.6%)	122 (50.0%)	109 (41.3%)	119 (45.8%)
2%)	25 (6.0%)	81 (6.4%)	10 (4.1%)	9 (3.4%)	15 (5.8%)
1.8%)	74 (17.8%)	198 (15.8%)	34 (13.9%)	36 (13.6%)	33 (12.7%)
3%)	22 (5.3%)	66 (5.3%)	13 (5.3%)	11 (4.2%)	13 (5.0%)
4.3%)	48 (11.6%)	140 (11.1%)	32 (13.1%)	39 (14.8%)	31 (11.9%)
5.2%)	70 (16.9%)	177 (14.1%)	37 (15.2%)	31 (11.7%)	30 (11.5%)
6%)	22 (5.3%)	72 (5.7%)	6 (2.5%)	7 (2.7%)	9 (3.5%)
3.3%)	121 (29.2%)	333 (26.5%)	58 (23.8%)	70 (26.5%)	65 (25.0%)
0%)	24 (5.8%)	83 (6.6%)	12 (4.9%)	16 (6.1%)	16 (6.2%)
2%)	41 (9.9%)	84 (6.7%)	15 (6.1%)	14 (5.3%)	14 (5.4%)
8%)	14 (3.4%)	46 (3.7%)	12 (4.9%)	13 (4.9%)	9 (3.5%)
2.7%)	138 (33.3%)	356 (28.3%)	29 (11.9%)	33 (12.5%)	42 (16.2%)
6%)	43 (10.4%)	109 (8.7%)	7 (2.9%)	8 (3.0%)	6 (2.3%)
2%)	48 (11.6%)	117 (9.3%)	15 (6.1%)	15 (5.7%)	18 (6.9%)
5.1%)	149 (35.9%)	444 (35.4%)	14 (5.7%)	16 (6.1%)	25 (9.6%)
9%)	41 (9.9%)	105 (8.4%)	2 (0.8%)	2 (0.8%)	6 (2.3%)
1%)	21 (4.8%)	65 (5.2%)	3 (1.2%)	4 (1.5%)	4 (1.5%)
5%)	40 (9.6%)	91 (7.3%)	5 (2.0%)	3 (1.1%)	11 (4.2%)
7%)	32 (7.7%)	67 (5.3%)	4 (1.6%)	1 (0.4%)	6 (2.3%)
0%)	28 (6.7%)	88 (7.0%)	4 (1.6%)	8 (3.0%)	13 (5.0%)
5.0%)	69 (16.6%)	211 (16.8%)	25 (10.2%)	19 (7.2%)	29 (11.2%)
7.4%)	72 (17.3%)	269 (21.4%)	15 (6.1%)	14 (5.3%)	22 (8.5%)
6%)	24 (5.8%)	95 (7.6%)	0	2 (0.8%)	2 (0.8%)
8%)	49 (11.8%)	129 (10.3%)	16 (6.6%)	6 (2.3%)	17 (6.5%)
2%)	22 (5.3%)	49 (3.9%)	9 (3.7%)	5 (1.9%)	9 (3.5%)
2.5%)	110 (26.5%)	339 (27.0%)	4 (1.6%)	7 (2.7%)	12 (4.6%)
8%)	29 (7.0%)	87 (6.9%)	2 (0.8%)	2 (0.8%)	1 (0.4%)
7%)	11 (2.7%)	14 (1.1%)	0	0	0
1.8%)	68 (16.4%)	224 (17.8%)	0	3 (1.1%)	5 (1.9%)
2%)	17 (4.1%)	39 (3.1%)	0	2 (0.8%)	2 (0.8%)
8%)	47 (11.3%)	133 (10.6%)	3 (1.2%)	2 (0.8%)	3 (1.2%)
7%)	7 (1.7)	31 (2.5%)	1 (0.4%)	2 (0.8%)	0
9%)	27 (6.5%)	60 (4.8%)	1 (0.4%)	0	2 (0.8%)
4%)	8 (1.9%)	38 (3.0%)	0	2 (0.8%)	1 (0.4%)
2%)	60 (14.5%)	154 (12.3%)	4 (1.6%)	1 (0.4%)	9 (3.5%)
4%)	21 (5.1%)	49 (3.9%)	2 (0.8%)	0	0
4%)	15 (3.6%)	47 (3.7%)	0	1 (0.4%)	3 (1.2%)

138 (33.3%)	356 (28.3%)	29 (11.9%)	33 (12.5%)	42 (16.2%)
43 (10.4%)	109 (8.7%)	7 (2.9%)	8 (3.0%)	6 (2.3%)
48 (11.6%)	117 (9.3%)	15 (6.1%)	15 (5.7%)	18 (6.9%)
149 (35.9%)	444 (35.4%)	14 (5.7%)	16 (6.1%)	25 (9.6%)
41 (9.9%)	105 (8.4%)	2 (0.8%)	2 (0.8%)	6 (2.3%)
21 (4.8%)	65 (5.2%)	3 (1.2%)	4 (1.5%)	4 (1.5%)
40 (9.6%)	91 (7.3%)	5 (2.0%)	3 (1.1%)	11 (4.2%)