


BNT327 (Biotheus / BioNTech)	
VEGF-A	
PD-L1	
None	
1L NSCLC:	47% ORR
1L TNBC:	79% ORR
2L SCLC:	61% ORR




PD-1xVEGF bispecific ivonesc superior outcomes when added


- Ivonescimab (PD-1xVEGF bispecific) + chemo outperforms chemo alone in 2L EGFRm+ NSCLC
 - Setting where Keytruda® failed*
- **Approved in 2L EGFRm NSCLC in China**

PD-L1xVEGF bispecific BNT301: compelling evidence of activity


Phase 1/2 study (NCT05918445):
efficacy signals in 1L NSCLC
Wu, C. et al., ASCO 2024, Poster #8533



Phase 2 (NCT05879066):
in combination with paclitaxel
Ying Cheng et al., ESMO 2023, Poster #100




Ivonescimab has demonstrated superiority to standard of care




27 has demonstrated ty in multiple solid tumors

38): clinical activity
clitaxel in 2L SCLC
Poster#1992P

ITT (n=36)	IO-naïve (n=22)	IO-treated (n=14)
61.1	72.7	42.9
86.1	81.8	92.9



Phase 2 (NCT05918133): clinical activity in
combination with nab-paclitaxel in 1L TNBC
Jiong Wu et al., SABCS 2023, Poster #PS08-06



ak11
phase III

H.

M

P

O

D





ARMONi-2 ITT (n=398); <i>Median Follow-up: 8.67 mos</i>	Ivonescimab (n=198)	
Median PFS	11.14 mos (95% CI: 7.33, NE)	
FS Stratified HR		0.51 (95% CI: 0.38, 0.6)
ORR	50.0% (95% CI: 42.8%, 57.2%)	
CR	89.9% (95% CI: 84.8%, 93.7%)	

HARMONi-2 Subgroup Analyses	Ivonescimab
PD-L1 High (PD-L1 TPS $\geq$ 50%) PFS stratified HR <i>Ivonescimab n=83; Pembrolizumab n=85</i>	(95% CI: 0.38, 0.6)
PD-L1 Low (PD-L1 TPS 1-49%) PFS stratified HR <i>Ivonescimab n=115; Pembrolizumab n=115</i>	(95% CI: 0.38, 0.6)
Squamous histology PFS stratified HR <i>Ivonescimab n=90; Pembrolizumab n=91</i>	(95% CI: 0.38, 0.6)
Non-Squamous histology PFS stratified HR <i>Ivonescimab n=108; Pembrolizumab n=109</i>	(95% CI: 0.38, 0.6)

syn 2510

Pembrolizumab (n=200)
5.82 mos (95% CI: 5.03, 8.21)
9; p<0.0001)
38.5% (95% CI: 31.7%, 45.6%)
70.5% (95% CI: 63.7%, 76.7%)

imab vs. Pembrolizumab
0.46 95% CI: 0.28, 0.75)
0.54 95% CI: 0.37, 0.79)
0.48 95% CI: 0.31, 0.74)
0.54 95% CI: 0.36, 0.82)



Treatment of d

Paclitaxel albumin-binding
type+Carboplatin

Docetaxel+Punebrine/Placebo


MW11 (PD-1 mAb)

IMM2510 (3mg/kg)

Note: Compassionate Use/Expanded Acc

Thymus


Treatment of d





Squamous NSCLC

Duration for anti-tumor therapies and best of response



Treatment of duration

Toripalimab (PD-1 mAb)+Lenvatinib

Docetaxel+Cisplatin

BAT6005 (TIGIT mAb)


QBH196 (multitarget kinase inhibitor)

IMM2510 (10mg/kg)

Note: Termination of treatment

adenosquamous carcinoma


Duration for anti-tumor therapies and best of response



SYN
escalate
with a
and n

amous NSCLC

ion for anti-tumor therapies and best of response



N-2510 completed dose
 ation up to 20 mg/kg Q2W
 manageable safety profile
 no observed dose-limiting
 toxicities (DLT)