

citation	wavelength range (nm)
[1]	6.2 - 26.2
[2]	26.2 - 66.2
[3]	66.2 - 76.2
[4]	76.2 - 96.2
[5]	96.2 - 106.2
[6]	106.2 - 116.2
[7]	116.2 - 146.2
[8]	146.2 - 196.2
[9]	196.2 - 206.2
[10]	206.2 - 226.2
[11]	226.2 - 236.3




Figure 1: C2H2

References

- [1] Glyn Cooper, Gordon R. Burton, and C.E. Brion. Absolute UV and soft x-ray photoabsorption of acetylene by high resolution dipole (e,e) spectroscopy. *Journal of Electron Spectroscopy and Related Phenomena*, 73(2):139–148, jul 1995.
- [2] C. Y. Robert Wu and D. L. Judge. Photoabsorption cross section of acetylene in the EUV region. *The Journal of Chemical Physics*, 82(10):4495–4499, may 1985.
- [3] P. H. Metzger and G. R. Cook. On the continuous absorption, photoionization, and fluorescence of h₂o, NH₃, CH₄, c₂h₂, c₂h₄, and c₂h₆ in the 600-to-1000-Å region. *The Journal of Chemical Physics*, 41(3):642–648, aug 1964.
- [4] J. C. Han, Chao Ye, Masako Suto, and L. C. Lee. Fluorescence from photoexcitation of c₂h₂ at 50–106 nm. *The Journal of Chemical Physics*, 90(8):4000–4007, apr 1989.
- [5] W. C. Walker and G. L. Weissler. Preliminary data on photoionization efficiencies and cross sections in c₂h₄ and c₂h₂. *The Journal of Chemical Physics*, 23(8):1547–1548, aug 1955.
- [6] Toshio Nakayama and K. Watanabe. Absorption and photoionization coefficients of acetylene, propyne, and 1-butyne. *The Journal of Chemical Physics*, 40(2):558–561, jan 1964.
- [7] Bing-Ming Cheng, Hui-Fen Chen, Hsiao-Chi Lu, Hong-Kai Chen, M. S. Alam, Sheng-Lung Chou, and Meng-Yeh Lin. ABSORPTION CROSS SECTION OF GASEOUS ACETYLENE

AT 85 k IN THE WAVELENGTH RANGE 110-155 nm. *The Astrophysical Journal Supplement Series*, 196(1):3, aug 2011.

- [8] Peter L. Smith, Kouichi Yoshino, W. H. Parkinson, Kenji Ito, and Glenn Stark. High-resolution, VUV (147–201 nm) photoabsorption cross sections for c2h2at 195 and 295 k. *Journal of Geophysical Research*, 96(E2):17529, 1991.
- [9] C. Y. Robert Wu, T. S. Chien, G. S. Liu, D. L. Judge, and J. J. Caldwell. Photoabsorption and direct dissociation cross sections of c2h2 in the 1530–1930 {aa region: A temperature dependent study. *The Journal of Chemical Physics*, 91(1):272–280, jul 1989.
- [10] Fengzhong Chen, D. L. Judge, C. Y. Robert Wu, J. Caldwell, H. Peter White, and R. Wager. High-resolution, low-temperature photoabsorption cross sections of c2h2, PH3, AsH3, and GeH4, with application to saturn's atmosphere. *Journal of Geophysical Research*, 96(E2):17519, 1991.
- [11] Y. Bénilan, N. Smith, A. Jolly, and F. Raulin. The long wavelength range temperature variations of the mid-UV acetylene absorption coefficient. *Planetary and Space Science*, 48(5):463–471, apr 2000.