| citation | wavelength range (nm) |
|----------|-----------------------|
| [1]      | 6.2 - 26.2            |
| [2]      | 26.2 - 66.2           |
| [3]      | 66.2 - 76.2           |
| [4]      | 76.2 - 96.2           |
| [5]      | 96.2 - 106.2          |
| [6]      | 106.2 - 116.2         |
| [7]      | 116.2 - 146.2         |
| [8]      | 146.2 - 196.2         |
| [9]      | 196.2 - 206.2         |
| [10]     | 206.2 - 226.2         |
| [11]     | 226.2 - 236.3         |




Figure 1: C2H2

## References

- [1] Glyn Cooper, Gordon R. Burton, and C.E. Brion. Absolute UV and soft x-ray photoabsorption of acetylene by high resolution dipole (e,e) spectroscopy. *Journal of Electron Spectroscopy and Related Phenomena*, 73(2):139–148, jul 1995.
- [2] C. Y. Robert Wu and D. L. Judge. Photoabsorption cross section of acetylene in the EUV region. *The Journal of Chemical Physics*, 82(10):4495–4499, may 1985.
- [3] P. H. Metzger and G. R. Cook. On the continuous absorption, photoionization, and fluorescence of h2o, NH3, CH4, c2h2, c2h4, and c2h6in the 600-to-1000-{aa region. *The Journal of Chemical Physics*, 41(3):642–648, aug 1964.
- [4] J. C. Han, Chao Ye, Masako Suto, and L. C. Lee. Fluorescence from photoexcitation of c2h2at 50–106 nm. *The Journal of Chemical Physics*, 90(8):4000–4007, apr 1989.
- [5] W. C. Walker and G. L. Weissler. Preliminary data on photoionization efficiencies and cross sections in c2h4 and c2h2. *The Journal of Chemical Physics*, 23(8):1547–1548, aug 1955.
- [6] Toshio Nakayama and K. Watanabe. Absorption and photoionization coefficients of acetylene, propyne, and 1-butyne. *The Journal of Chemical Physics*, 40(2):558–561, jan 1964.
- [7] Bing-Ming Cheng, Hui-Fen Chen, Hsiao-Chi Lu, Hong-Kai Chen, M. S. Alam, Sheng-Lung Chou, and Meng-Yeh Lin. ABSORPTION CROSS SECTION OF GASEOUS ACETYLENE

- AT 85 k IN THE WAVELENGTH RANGE 110-155 nm. The Astrophysical Journal Supplement Series, 196(1):3, aug 2011.
- [8] Peter L. Smith, Kouichi Yoshino, W. H. Parkinson, Kenji Ito, and Glenn Stark. High-resolution, VUV (147–201 nm) photoabsorption cross sections for c2h2at 195 and 295 k. *Journal of Geophysical Research*, 96(E2):17529, 1991.
- [9] C. Y. Robert Wu, T. S. Chien, G. S. Liu, D. L. Judge, and J. J. Caldwell. Photoabsorption and direct dissociation cross sections of c2h2 in the 1530–1930 {aa region: A temperature dependent study. *The Journal of Chemical Physics*, 91(1):272–280, jul 1989.
- [10] Fengzhong Chen, D. L. Judge, C. Y. Robert Wu, J. Caldwell, H. Peter White, and R. Wagener. High-resolution, low-temperature photoabsorption cross sections of c2h2, PH3, AsH3, and GeH4, with application to saturn's atmosphere. *Journal of Geophysical Research*, 96(E2):17519, 1991.
- [11] Y. Bénilan, N. Smith, A. Jolly, and F. Raulin. The long wavelength range temperature variations of the mid-UV acetylene absorption coefficient. *Planetary and Space Science*, 48(5):463–471, apr 2000.