

FIGURE 4–3a. ASSESSING ASTHMA CONTROL AND ADJUSTING THERAPY IN CHILDREN 0–4 YEARS OF AGE

Components of Control		Classification of Asthma Control (0–4 years of age)		
		Well Controlled	Not Well Controlled	Very Poorly Controlled
Impairment	Symptoms	≤2 days/week	>2 days/week	Throughout the day
	Nighttime awakenings	≤1x/month	>1x/month	>1x/week
	Interference with normal activity	None	Some limitation	Extremely limited
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week	Several times per day
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	2–3/year	>3/year
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.		
Recommended Action for Treatment (See figure 4–1a for treatment steps.)		<ul style="list-style-type: none"> • Maintain current treatment. • Regular followup every 1–6 months. • Consider step down if well controlled for at least 3 months. 	<ul style="list-style-type: none"> • Step up (1 step) and Reevaluate in 2–6 weeks. • If no clear benefit in 4–6 weeks, consider alternative diagnoses or adjusting therapy. • For side effects, consider alternative treatment options. 	<ul style="list-style-type: none"> • Consider short course of oral systemic corticosteroids. • Step up (1–2 steps), and Reevaluate in 2 weeks. • If no clear benefit in 4–6 weeks, consider alternative diagnoses or adjusting therapy. • For side effects, consider alternative treatment options.

Key: EIB, exercise-induced bronchospasm

Notes:

- The stepwise approach is meant to assist, not replace, the clinical decisionmaking required to meet individual patient needs.
- The level of control is based on the most severe impairment or risk category. Assess impairment domain by caregiver's recall of previous 2–4 weeks. Symptom assessment for longer periods should reflect a global assessment such as inquiring whether the patient's asthma is better or worse since the last visit.
- At present, there are inadequate data to correspond frequencies of exacerbations with different levels of asthma control. In general, more frequent and intense exacerbations (e.g., requiring urgent, unscheduled care, hospitalization, or ICU admission) indicate poorer disease control. For treatment purposes, patients who had ≥2 exacerbations requiring oral systemic corticosteroids in the past year may be considered the same as patients who have not-well-controlled asthma, even in the absence of impairment levels consistent with not-well-controlled asthma.
- Before step up in therapy:
 - Review adherence to medications, inhaler technique, and environmental control.
 - If alternative treatment option was used in a step, discontinue it and use preferred treatment for that step.