

COMP2022: Formal Languages and Logic

2017, Semester 1, Week 11

Joseph Godbehere

Adapted from slides by A/Prof Kalina Yacef

May 23, 2017



THE UNIVERSITY OF
SYDNEY

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING

This material has been reproduced and communicated to you by or on behalf of the University of Sydney pursuant to part VB of the Copyright Act 1968 (**the Act**).

The material in this communication may be subject to copyright under the Act. Any further copying or communication of this material by you may be subject of copyright protect under the Act.

Do not remove this notice.

ANNOUNCEMENTS

Assignment 3:

- ▶ Due on Thursday of week **12**
- ▶ May be completed individually or in pairs
- ▶ Proofs on the Logic Tutor
- ▶ 4 marks for correctness
- ▶ 1 mark based on proof length

OUTLINE

Predicate Logic (continued)

- ▶ Substitution of variables
- ▶ Semantics in predicate logic
- ▶ Validity, Satisfiability
- ▶ “Free to replace”
- ▶ Equivalence Laws involving quantifiers
- ▶ Inference Rules involving quantifiers
- ▶ Tautologies

SUBSTITUTION FOR FREE VARIABLES

Suppose W is a wff, x is a free variable in W , and t is a term

$W[t/x]$ is the wff obtained by substituting t for all *free* occurrences of x

i.e. replacing all *free* occurrences of x with t

Examples:

- ▶ If $W = \forall xP(x, y)$ then
 - ▶ $W[t/x] = \forall xP(x, y) = W$ (x was already bound)
 - ▶ $W[t/y] = \forall xP(x, t) \neq W$
- ▶ If $Z = (P(x, y) \vee \exists yQ(x, y))$ then
 - ▶ $Z[t/x] = (P(t, y) \vee \exists yQ(t, y)) \neq Z$
 - ▶ $Z[t/y] = (P(x, t) \vee \exists yQ(x, y)) \neq Z$

MEANING (SEMANTICS) OF A WFF

The meaning of a wff with respect to an interpretation with domain D , is the truth value obtained by applying the following rules:

- ▶ If the wff has no quantifier, then the truth value of the proposition is obtained by applying the interpretation to the wff
- ▶ $\forall xW$ is true if $W[d/x]$ is true for every $d \in D$. Otherwise it is false.
- ▶ $\exists xW$ is true if $W[d/x]$ is true for some $d \in D$. Otherwise it is false.

EXAMPLE

Existential quantification on the domain of interpretation

$$D = \{a_1, a_2, \dots, a_n\}$$

$$\exists x P(x) \equiv P(a_1) \vee P(a_2) \vee \dots \vee P(a_n)$$

e.g. suppose

- ▶ $D = \{a, b, c, d, e\}$
- ▶ $P(a), P(b), P(c), Q(a), Q(b), R(c), R(d)$ are true, all others false.

Then is $\exists x (P(x) \wedge (\neg Q(x) \rightarrow R(x)))$ true or false?

EXAMPLE

Existential quantification on the domain of interpretation

$$D = \{a_1, a_2, \dots, a_n\}$$

$$\exists x P(x) \equiv P(a_1) \vee P(a_2) \vee \dots \vee P(a_n)$$

e.g. suppose

- ▶ $D = \{a, b, c, d, e\}$
- ▶ $P(a), P(b), P(c), Q(a), Q(b), R(c), R(d)$ are true, all others false.

Then is $\exists x (P(x) \wedge (\neg Q(x) \rightarrow R(x)))$ true or false?

- ▶ Let $W = (P(x) \wedge (\neg Q(x) \rightarrow R(x)))$
- ▶ Notice that $W[c/x] = (P(c) \wedge (\neg Q(c) \rightarrow R(c)))$ is true
- ▶ Therefore $\exists x W$ is true

EXAMPLE

Universal quantification on the domain of interpretation

$$D = \{a_1, a_2, \dots, a_n\}$$

$$\forall x P(x) \equiv P(a_1) \wedge P(a_2) \wedge \dots \wedge P(a_n)$$

e.g. suppose

- ▶ $D = \{a, b, c, d, e\}$
- ▶ $P(a), P(b), P(c), Q(a), Q(b), R(c), R(d)$ are true, all others false.

Then is $\forall x (P(x) \rightarrow (Q(x) \vee R(x)))$ true or false?

EXAMPLE

Universal quantification on the domain of interpretation

$$D = \{a_1, a_2, \dots, a_n\}$$

$$\forall x P(x) \equiv P(a_1) \wedge P(a_2) \wedge \dots \wedge P(a_n)$$

e.g. suppose

- ▶ $D = \{a, b, c, d, e\}$
- ▶ $P(a), P(b), P(c), Q(a), Q(b), R(c), R(d)$ are true, all others false.

Then is $\forall x (P(x) \rightarrow (Q(x) \vee R(x)))$ true or false?

- ▶ Let $W = (P(x) \rightarrow (Q(x) \vee R(x)))$
- ▶ $W[a/x] = (P(a) \rightarrow (Q(a) \vee R(a))) = T \rightarrow (T \vee F) = T$
- ▶ $W[b/x] = (P(b) \rightarrow (Q(b) \vee R(b))) = T \rightarrow (T \vee F) = T$
- ▶ $W[c/x] = (P(c) \rightarrow (Q(c) \vee R(c))) = T \rightarrow (F \vee T) = T$
- ▶ $W[d/x] = (P(d) \rightarrow (Q(d) \vee R(d))) = F \rightarrow (F \vee T) = T$
- ▶ $W[e/x] = (P(e) \rightarrow (Q(e) \vee R(e))) = F \rightarrow (F \vee F) = T$
- ▶ so $\forall x W \equiv W[a/x] \wedge W[b/x] \wedge W[c/x] \wedge W[d/x] \wedge W[e/x]$
is true

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	
$\forall a P(a, \text{Andy})$	
$\forall a \exists c P(a, c)$	
$\forall c \exists a P(a, c)$	
$\exists a \forall c P(a, c)$	
$\exists c \forall a P(a, c)$	
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	
$\forall a \exists c P(a, c)$	
$\forall c \exists a P(a, c)$	
$\exists a \forall c P(a, c)$	
$\exists c \forall a P(a, c)$	
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	
$\forall c \exists a P(a, c)$	
$\exists a \forall c P(a, c)$	
$\exists c \forall a P(a, c)$	
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	
$\exists a \forall c P(a, c)$	
$\exists c \forall a P(a, c)$	
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	
$\exists c \forall a P(a, c)$	
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	false
$\exists c \forall a P(a, c)$	
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	false
$\exists c \forall a P(a, c)$	false
$\exists c \forall a \neg P(a, c)$	
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	false
$\exists c \forall a P(a, c)$	false
$\exists c \forall a \neg P(a, c)$	false
$\exists c \exists a P(a, c)$	
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	false
$\exists c \forall a P(a, c)$	false
$\exists c \forall a \neg P(a, c)$	false
$\exists c \exists a P(a, c)$	true
$\exists a \exists c P(a, c)$	
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	false
$\exists c \forall a P(a, c)$	false
$\exists c \forall a \neg P(a, c)$	false
$\exists c \exists a P(a, c)$	true
$\exists a \exists c P(a, c)$	true
$\forall a \forall c P(a, c)$	

EXAMPLES

$P(a, c)$ “ a is a parent of c ”, Domain = humans

Suppose Jim is a parent of Andy, but Fred is not a parent of anyone

$P(\text{Jim}, \text{Andy})$	true
$P(\text{Fred}, \text{Andy})$	false
$\exists a P(a, \text{Andy})$	true (e.g. $a = \text{Jim}$)
$\forall a P(a, \text{Andy})$	false (e.g. $a = \text{Fred}$)
$\forall a \exists c P(a, c)$	false (e.g. $a = \text{Fred}$)
$\forall c \exists a P(a, c)$	true
$\exists a \forall c P(a, c)$	false
$\exists c \forall a P(a, c)$	false
$\exists c \forall a \neg P(a, c)$	false
$\exists c \exists a P(a, c)$	true
$\exists a \exists c P(a, c)$	true
$\forall a \forall c P(a, c)$	false

DIFFERENT INTERPRETATIONS LEAD TO DIFFERENT SEMANTICS

$$W = \exists x \forall y (P(y) \rightarrow Q(x, y))$$

$Q(x, y)$: “ x is equal to y ”

- ▶ Let $D = \{1\}$ and $P(1) = \text{true}$. Then W is true.
- ▶ Let $D = \{1\}$ and $P(1) = \text{false}$. Then W is true.
- ▶ Let $D = \{1, 2\}$ and $P(1) = P(2) = \text{true}$. Then W is false.
 - ▶ Let $x = 1$, then
 - ▶ Let $y = 1$, then $(P(1) \rightarrow Q(1, 1)) = \text{true}$
 - ▶ Let $y = 2$, then $(P(1) \rightarrow Q(1, 2)) = \text{false}$
 - ▶ $\text{true} \wedge \text{false} = \text{false}$, so $\forall y (P(y) \rightarrow Q(1, y))$ is false
 - ▶ Let $x = 2$, then
 - ▶ Let $y = 1$, then $(P(2) \rightarrow Q(2, 1)) = \text{false}$
 - ▶ Let $y = 2$, then $(P(2) \rightarrow Q(2, 2)) = \text{true}$
 - ▶ $\text{false} \wedge \text{true} = \text{false}$, so $\forall y (P(y) \rightarrow Q(2, y))$ is false
 - ▶ $\text{false} \vee \text{false} = \text{false}$, so $\exists x \forall y (P(y) \rightarrow Q(x, y))$ is false

VALIDITY AND SATISFIABILITY

A wff is *valid* if it is *true for all* possible interpretations, otherwise it is *invalid*

A wff is *unsatisfiable* if it is *false for all* possible interpretations, otherwise it is *satisfiable*

- ▶ Valid (and satisfiable) \rightarrow tautology
- ▶ (Invalid and) unsatisfiable \rightarrow contradiction
- ▶ Invalid but satisfiable \rightarrow contingency

VALIDITY AND SATISFIABILITY

- ▶ $\exists x \forall y (P(y) \rightarrow Q(x, y))$ is satisfiable and invalid
 - ▶ Satisfiable because there is some interpretation for which it is not false
 - ▶ Invalid because it is not true for all interpretations
- ▶ $\forall x (P(x) \vee \neg P(x))$ is a tautology
- ▶ $\forall x (P(x) \wedge \neg P(x))$ is a contradiction

LOGICAL EQUIVALENCES/IMPLICATIONS

Propositions are predicates of arity 0, so all the tautologies for propositional logic also hold in predicate logic

- | | |
|---|---|
| ▶ $(A \vee \neg A)$ | ▶ $A \rightarrow B, A \vdash B$ |
| ▶ $(A(x) \vee \neg A(x))$ | ▶ $A(x) \rightarrow B(y), A(x) \vdash B(y)$ |
| ▶ $(\forall A(x) \vee \neg \forall A(x))$ | ▶ $\forall x A(x) \rightarrow \forall y B(y), \forall x A(x) \vdash \forall y B(y)$ |
| ▶ ... | ▶ ... |

LOGICAL EQUIVALENCES/IMPLICATIONS

Propositions are predicates of arity 0, so all the tautologies for propositional logic also hold in predicate logic

- ▶ $(A \vee \neg A)$
- ▶ $(A(x) \vee \neg A(x))$
- ▶ $(\forall A(x) \vee \neg \forall A(x))$
- ▶ ...
- ▶ $A \rightarrow B, A \vdash B$
- ▶ $A(x) \rightarrow B(y), A(x) \vdash B(y)$
- ▶ $\forall x A(x) \rightarrow \forall y B(y), \forall x A(x) \vdash \forall y B(y)$
- ▶ ...

Therefore all the laws of equivalence and rules of inferences seen in propositional logic can also be for predicate logic, when each proposition is replaced by a wff for predicate logic.

LOGICAL EQUIVALENCES/IMPLICATIONS

Propositions are predicates of arity 0, so all the tautologies for propositional logic also hold in predicate logic

- ▶ $(A \vee \neg A)$
- ▶ $(A(x) \vee \neg A(x))$
- ▶ $(\forall A(x) \vee \neg \forall A(x))$
- ▶ ...
- ▶ $A \rightarrow B, A \vdash B$
- ▶ $A(x) \rightarrow B(y), A(x) \vdash B(y)$
- ▶ $\forall x A(x) \rightarrow \forall y B(y), \forall x A(x) \vdash \forall y B(y)$
- ▶ ...

Therefore all the laws of equivalence and rules of inferences seen in propositional logic can also be for predicate logic, when each proposition is replaced by a wff for predicate logic.

We also need specific laws and rules involving quantifiers:

- ▶ Laws of equivalence involving quantifiers
- ▶ Rules of inference involving quantifiers (adding and removing quantifiers)



NATURAL DEDUCTION IN PREDICATE LOGIC

Simple example which does not involve reasoning with quantifiers:

Premises as below $\vdash (P(f(x)) \rightarrow \forall y P(y))$

Premises	Line	Formula	Just.	Refs.
1	1	$(\forall x P(x) \rightarrow Q(f(x), x))$	P	
2	2	$\neg Q(f(x), x)$	P	
3	3	$(\neg Q(f(x), x) \rightarrow (\exists y Q(x, y) \rightarrow \forall y P(y)))$	P	
4	4	$(\forall x P(x) \vee (P(f(x)) \rightarrow \exists y Q(x, y)))$	P	
2,3	5	$(\exists y Q(x, y) \rightarrow \forall y P(y))$	MP	2,3
1,2	6	$\neg \forall x P(x)$	MT	1,2
1,2,4	7	$(P(f(x)) \rightarrow \exists y Q(x, y))$	DS	4,6
1,2,3,4	8	$(P(f(x)) \rightarrow \forall y P(y))$	HS	5,7

NATURAL DEDUCTION IN PREDICATE LOGIC

Simple example which does not involve reasoning with quantifiers:

Premises as below $\vdash (P(f(x)) \rightarrow \forall yP(y))$

Let $A = P(f(x))$, $B = \forall yP(y)$, $C = \exists yQ(x, y)$, $D = Q(f(x), x)$, $E = \forall xP(x)$

Premises	Line	Formula	Just.	Refs.
1	1	$(E \rightarrow D)$	P	
2	2	$\neg D$	P	
3	3	$(\neg D \rightarrow (C \rightarrow B))$	P	
4	4	$(E \vee (A \rightarrow C))$	P	
2,3	5	$(C \rightarrow B)$	MP	2,3
1,2	6	$\neg E$	MT	1,2
1,2,4	7	$(A \rightarrow C)$	DS	4,6
1,2,3,4	8	$(A \rightarrow B)$	HS	5,7

LAWS OF EQUIVALENCE INVOLVING QUANTIFIERS

Let F and G be arbitrary wffs

Quantifiers and Negation (QNeg)

$$\neg \forall x F \equiv \exists x \neg F$$

$$\neg \exists x F \equiv \forall x \neg F$$

Interchanging of quantifiers of same type (QInter)

LAWS OF EQUIVALENCE INVOLVING QUANTIFIERS

Let F and G be arbitrary wffs

Quantifiers and Negation (QNeg)

$$\neg \forall x F \equiv \exists x \neg F$$

$$\neg \exists x F \equiv \forall x \neg F$$

Distribution of quantifiers (QDistr)

$$(\forall x F \wedge \forall x G) \equiv \forall x (F \wedge G)$$

$$(\exists x F \vee \exists x G) \equiv \exists x (F \vee G)$$

Interchanging of quantifiers of same type (QInter)

LAWS OF EQUIVALENCE INVOLVING QUANTIFIERS

Let F and G be arbitrary wffs

Quantifiers and Negation (QNeg)

$$\neg \forall x F \equiv \exists x \neg F$$

$$\neg \exists x F \equiv \forall x \neg F$$

Distribution of quantifiers (QDistr)

$$(\forall x F \wedge \forall x G) \equiv \forall x (F \wedge G)$$

$$(\exists x F \vee \exists x G) \equiv \exists x (F \vee G)$$

Interchanging of quantifiers of same type (QInter)

$$\forall x \forall y F \equiv \forall y \forall x F$$

$$\exists x \exists y F \equiv \exists y \exists x F$$

LAWS OF EQUIVALENCE INVOLVING QUANTIFIERS

Let F and G be arbitrary wffs

Extraction of quantifiers (QExtr)

If x does not occur in G :

$$(\forall x F \wedge G) \equiv \forall x (F \wedge G)$$

$$(\forall x F \vee G) \equiv \forall x (F \vee G)$$

$$(\exists x F \wedge G) \equiv \exists x (F \wedge G)$$

$$(\exists x F \vee G) \equiv \exists x (F \vee G)$$

LAWS OF EQUIVALENCE INVOLVING QUANTIFIERS

Let F and G be arbitrary wffs

Extraction of quantifiers (QExtr)

If x does not occur in G :

similarly:

$$(\forall x F \wedge G) \equiv \forall x (F \wedge G)$$

$$(G \wedge \forall x F) \equiv \forall x (G \wedge F)$$

$$(\forall x F \vee G) \equiv \forall x (F \vee G)$$

$$(G \vee \forall x F) \equiv \forall x (G \vee F)$$

$$(\exists x F \wedge G) \equiv \exists x (F \wedge G)$$

$$(G \wedge \exists x F) \equiv \exists x (G \wedge F)$$

$$(\exists x F \vee G) \equiv \exists x (F \vee G)$$

$$(G \vee \exists x F) \equiv \exists x (G \vee F)$$

EXAMPLE

Suppose we want to derive the following equivalence:

$$\neg(\exists xP(x, y) \vee \forall z\neg R(z)) \equiv \forall x\exists z(\neg P(x, y) \wedge R(z))$$

$$\neg(\exists xP(x, y) \vee \forall z\neg R(z))$$

EXAMPLE

Suppose we want to derive the following equivalence:

$$\neg(\exists xP(x, y) \vee \forall z\neg R(z)) \equiv \forall x\exists z(\neg P(x, y) \wedge R(z))$$

$$\begin{aligned} & \neg(\exists xP(x, y) \vee \forall z\neg R(z)) \\ \equiv & (\neg\exists xP(x, y) \wedge \neg\forall z\neg R(z)) \end{aligned} \quad \text{DeMorgan's Laws}$$

EXAMPLE

Suppose we want to derive the following equivalence:

$$\neg(\exists xP(x, y) \vee \forall z\neg R(z)) \equiv \forall x\exists z(\neg P(x, y) \wedge R(z))$$

$$\begin{aligned} & \neg(\exists xP(x, y) \vee \forall z\neg R(z)) \\ \equiv & (\neg\exists xP(x, y) \wedge \neg\forall z\neg R(z)) && \text{DeMorgan's Laws} \\ \equiv & (\forall x\neg P(x, y) \wedge \exists z\neg\neg R(z)) && \text{Quantifier Negation} \end{aligned}$$

EXAMPLE

Suppose we want to derive the following equivalence:

$$\neg(\exists xP(x, y) \vee \forall z\neg R(z)) \equiv \forall x\exists z(\neg P(x, y) \wedge R(z))$$

$$\begin{aligned} & \neg(\exists xP(x, y) \vee \forall z\neg R(z)) \\ \equiv & (\neg\exists xP(x, y) \wedge \neg\forall z\neg R(z)) && \text{DeMorgan's Laws} \\ \equiv & (\forall x\neg P(x, y) \wedge \exists z\neg\neg R(z)) && \text{Quantifier Negation} \\ \equiv & (\forall x\neg P(x, y) \wedge \exists zR(z)) && \text{Double Negation} \end{aligned}$$

EXAMPLE

Suppose we want to derive the following equivalence:

$$\neg(\exists x P(x, y) \vee \forall z \neg R(z)) \equiv \forall x \exists z (\neg P(x, y) \wedge R(z))$$

$$\begin{aligned} & \neg(\exists x P(x, y) \vee \forall z \neg R(z)) \\ \equiv & (\neg \exists x P(x, y) \wedge \neg \forall z \neg R(z)) && \text{DeMorgan's Laws} \\ \equiv & (\forall x \neg P(x, y) \wedge \exists z \neg \neg R(z)) && \text{Quantifier Negation} \\ \equiv & (\forall x \neg P(x, y) \wedge \exists z R(z)) && \text{Double Negation} \\ \equiv & \forall x (\neg P(x, y) \wedge \exists z R(z)) && \text{Quantifier Extraction} \end{aligned}$$

EXAMPLE

Suppose we want to derive the following equivalence:

$$\neg(\exists xP(x, y) \vee \forall z\neg R(z)) \equiv \forall x\exists z(\neg P(x, y) \wedge R(z))$$

$$\begin{aligned} & \neg(\exists xP(x, y) \vee \forall z\neg R(z)) \\ \equiv & (\neg\exists xP(x, y) \wedge \neg\forall z\neg R(z)) && \text{DeMorgan's Laws} \\ \equiv & (\forall x\neg P(x, y) \wedge \exists z\neg\neg R(z)) && \text{Quantifier Negation} \\ \equiv & (\forall x\neg P(x, y) \wedge \exists zR(z)) && \text{Double Negation} \\ \equiv & \forall x(\neg P(x, y) \wedge \exists zR(z)) && \text{Quantifier Extraction} \\ \equiv & \forall x\exists z(\neg P(x, y) \wedge R(z)) && \text{Quantifier Extraction} \end{aligned}$$

REASONING WITH QUANTIFIERS

Simple example: if a property holds for everything then it holds for a particular thing.

- ▶ So, it seems reasonable to deduce $W(a)$ from $\forall x W(x)$
 - ▶ Similarly ($W(b)$, $W(x)$, $W(f(x))$, ... etc.)

REASONING WITH QUANTIFIERS

Simple example: if a property holds for everything then it holds for a particular thing.

- ▶ So, it seems reasonable to deduce $W(a)$ from $\forall x W(x)$
 - ▶ Similarly ($W(b)$, $W(x)$, $W(f(x))$, ... etc.)
- ▶ *But not always!*

Consider $\forall x \exists y P(x, y)$

- ▶ Can we deduce $\exists y P(a, y)$?
- ▶ Can we deduce $\exists y P(x, y)$?
- ▶ Can we deduce $\exists y P(y, y)$?

REASONING WITH QUANTIFIERS

Simple example: if a property holds for everything then it holds for a particular thing.

- ▶ So, it seems reasonable to deduce $W(a)$ from $\forall x W(x)$
 - ▶ Similarly ($W(b)$, $W(x)$, $W(f(x))$, ... etc.)
- ▶ *But not always!*

Consider $\forall x \exists y P(x, y)$

- ▶ Can we deduce $\exists y P(a, y)$? YES
- ▶ Can we deduce $\exists y P(x, y)$?
- ▶ Can we deduce $\exists y P(y, y)$?

REASONING WITH QUANTIFIERS

Simple example: if a property holds for everything then it holds for a particular thing.

- ▶ So, it seems reasonable to deduce $W(a)$ from $\forall x W(x)$
 - ▶ Similarly ($W(b)$, $W(x)$, $W(f(x))$, ... etc.)
- ▶ *But not always!*

Consider $\forall x \exists y P(x, y)$

- ▶ Can we deduce $\exists y P(a, y)$? YES
- ▶ Can we deduce $\exists y P(x, y)$? YES
- ▶ Can we deduce $\exists y P(y, y)$?

REASONING WITH QUANTIFIERS

Simple example: if a property holds for everything then it holds for a particular thing.

- ▶ So, it seems reasonable to deduce $W(a)$ from $\forall x W(x)$
 - ▶ Similarly ($W(b)$, $W(x)$, $W(f(x))$, ... etc.)
- ▶ *But not always!*

Consider $\forall x \exists y P(x, y)$

- ▶ Can we deduce $\exists y P(a, y)$? YES
- ▶ Can we deduce $\exists y P(x, y)$? YES
- ▶ Can we deduce $\exists y P(y, y)$? NO

REASONING WITH QUANTIFIERS

e.g. If $P(x,y)$ means x is the child of y then:

- ▶ $\forall x \exists y P(x, y)$ means
- ▶ $\exists y P(x, y)$ means
- ▶ $\exists y P(y, y)$ means

REASONING WITH QUANTIFIERS

e.g. If $P(x,y)$ means x is the child of y then:

- ▶ $\forall x \exists y P(x, y)$ means everyone is the child of someone
- ▶ $\exists y P(x, y)$ means
- ▶ $\exists y P(y, y)$ means

REASONING WITH QUANTIFIERS

e.g. If $P(x,y)$ means x is the child of y then:

- ▶ $\forall x \exists y P(x, y)$ means everyone is the child of someone
- ▶ $\exists y P(x, y)$ means x is the child of someone
- ▶ $\exists y P(y, y)$ means

REASONING WITH QUANTIFIERS

e.g. If $P(x,y)$ means x is the child of y then:

- ▶ $\forall x \exists y P(x, y)$ means everyone is the child of someone
- ▶ $\exists y P(x, y)$ means x is the child of someone
- ▶ $\exists y P(y, y)$ means someone is their own child (!?)

REASONING WITH QUANTIFIERS

e.g. If $P(x,y)$ means x is the child of y then:

- ▶ $\forall x \exists y P(x, y)$ means everyone is the child of someone
- ▶ $\exists y P(x, y)$ means x is the child of someone
- ▶ $\exists y P(y, y)$ means someone is their own child (!?)
(Beware the time travel paradox quokka!)

Trouble arises when we try to infer $W(t)$ from $\forall x W(x)$ where t :

1. Contains an occurrence of a quantified variable, and
2. x occurs free within the scope of that quantifier

DEFINITION: FREE TO REPLACE

Suppose t is a term and x is a free variable in F

t is *free to replace* x in F if either:

1. no variable in t occurs bound to a quantifier in F
2. or, x does not occur free within the scope of a quantifier in F

i.e. both F and $F[t/x]$ have the *same* bound occurrences of variables.

(Recall that $F[t/x]$ is the wff in obtained from F by substituting t for all free occurrences of x in F)

EXAMPLES: FREE TO REPLACE

t is *free to replace* x in F if either:

1. no variable in t occurs bound to a quantifier in F
2. or, x does not occur free within the scope of a quantifier in F

Example: $\forall y(P(y) \rightarrow G(y, x))$

- Is $f(y)$ free to replace x ?
- Is $f(x, y)$ free to replace x ?
- Is $f(x)$ free to replace x ?

EXAMPLES: FREE TO REPLACE

t is *free to replace* x in F if either:

1. no variable in t occurs bound to a quantifier in F
2. or, x does not occur free within the scope of a quantifier in F

Example: $\forall y(P(y) \rightarrow G(y, x))$

- Is $f(y)$ free to replace x ? NO, because a variable of $f(y)$ occurs bound and x occurs free within the scope of a quantifier.
- Is $f(x, y)$ free to replace x ?
- Is $f(x)$ free to replace x ?

EXAMPLES: FREE TO REPLACE

t is *free to replace* x in F if either:

1. no variable in t occurs bound to a quantifier in F
2. or, x does not occur free within the scope of a quantifier in F

Example: $\forall y(P(y) \rightarrow G(y, x))$

- ▶ Is $f(y)$ free to replace x ? NO, because a variable of $f(y)$ occurs bound and x occurs free within the scope of a quantifier.
- ▶ Is $f(x, y)$ free to replace x ? NO, because a variable of $f(x, y)$ occurs bound and x occurs free within the scope of a quantifier.
- ▶ Is $f(x)$ free to replace x ?

EXAMPLES: FREE TO REPLACE

t is *free to replace* x in F if either:

1. no variable in t occurs bound to a quantifier in F
2. or, x does not occur free within the scope of a quantifier in F

Example: $\forall y(P(y) \rightarrow G(y, x))$

- ▶ Is $f(y)$ free to replace x ? NO, because a variable of $f(y)$ occurs bound and x occurs free within the scope of a quantifier.
- ▶ Is $f(x, y)$ free to replace x ? NO, because a variable of $f(x, y)$ occurs bound and x occurs free within the scope of a quantifier.
- ▶ Is $f(x)$ free to replace x ? YES, because no variable of $f(x)$ occurs bound in the resulting formula

RULES OF INFERENCE INVOLVING QUANTIFIERS

\exists -elimination (Existential instantiation)

$$\frac{S \vdash \exists x F}{S \vdash F[c/x]} \quad \text{where } c \text{ is a new constant}$$

\forall -elimination (Universal Instantiation)

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

\exists -introduction (Existential generalisation)

$$\frac{S \vdash F}{S \vdash \exists x F[x/t]} \quad \text{where } x \text{ is free to replace } t \text{ in } F$$

\forall -introduction (Universal generalisation)

$$\frac{S \vdash F}{S \vdash \forall x F} \quad \begin{array}{l} x \text{ is not free in any of the premises used} \\ \text{to deduce } F, \text{ and } x \text{ is not free in any wff} \\ \text{constructed by } \exists\text{-elimination} \end{array}$$

\exists -ELIMINATION (EXISTENTIAL INSTANTIATION)

If a property holds for something, then it holds for a particular thing

$$\frac{S \vdash \exists x F}{S \vdash F[c/x]} \quad \text{where } c \text{ is a new constant}$$

i.e. c has not previously been defined (either earlier in the argument, or in the conclusion.)

∃-ELIMINATION (EXISTENTIAL INSTANTIATION)

If a property holds for something, then it holds for a particular thing

$$\frac{S \vdash \exists x F}{S \vdash F[c/x]} \quad \text{where } c \text{ is a new constant}$$

i.e. c has not previously been defined (either earlier in the argument, or in the conclusion.)

From $\exists x \forall y P(x, y)$ we can deduce $\forall y P(c, y)$ if c is a new constant.

\exists -ELIMINATION (EXISTENTIAL INSTANTIATION)

If a property holds for something, then it holds for a particular thing

$$\frac{S \vdash \exists x F}{S \vdash F[c/x]} \quad \text{where } c \text{ is a new constant}$$

i.e. c has not previously been defined (either earlier in the argument, or in the conclusion.)

From $\exists x \forall y P(x, y)$ we can deduce $\forall y P(c, y)$ if c is a new constant.

But these arguments don't work:

1	1	$\exists x P(x)$	P	
2	2	$\exists x Q(x)$	P	
1	3	$P(c)$	\exists -elim	1
2	4	$Q(c)$	\exists -elim	2

Line 4 is an error, as c was previously defined on line 3.

Attempt to prove $\exists x P(x) \vdash P(c)$

1	1	$\exists x P(x)$	P	
1	2	$P(c)$	\exists -elim	1

Line 2 is an error, as c was previously defined in the conclusion.

\forall -ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

► $\exists y G(2, y, f(2))$

∀-ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

- ▶ $\exists y G(2, y, f(2))$ YES
- ▶ $\exists y G(x, y, f(x))$

∀-ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

- ▶ $\exists y G(2, y, f(2))$ YES
- ▶ $\exists y G(x, y, f(x))$ YES
- ▶ $\exists y G(f(x), y, f(f(x)))$

\forall -ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

- ▶ $\exists y G(2, y, f(2))$ YES
- ▶ $\exists y G(x, y, f(x))$ YES
- ▶ $\exists y G(f(x), y, f(f(x)))$ YES
- ▶ $\exists y G(y, y, f(y))$

∀-ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

- ▶ $\exists y G(2, y, f(2))$ YES
- ▶ $\exists y G(x, y, f(x))$ YES
- ▶ $\exists y G(f(x), y, f(f(x)))$ YES
- ▶ $\exists y G(y, y, f(y))$ NO. Why not?

∀-ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

- ▶ $\exists y G(2, y, f(2))$ YES
- ▶ $\exists y G(x, y, f(x))$ YES
- ▶ $\exists y G(f(x), y, f(f(x)))$ YES
- ▶ $\exists y G(y, y, f(y))$ NO. Why not?

Suppose we know $\forall x G(x, y, f(x))$. Can we deduce:

- ▶ $G(y, y, f(y))$

∀-ELIMINATION (UNIVERSAL INSTANTIATION)

If a property holds for everything, then it holds for any particular thing:

$$\frac{S \vdash \forall x F}{S \vdash F[t/x]} \quad \text{where } t \text{ is free to replace } x \text{ in } F$$

Suppose we know $F = \forall x \exists y G(x, y, f(x))$. Can we deduce:

- ▶ $\exists y G(2, y, f(2))$ YES
- ▶ $\exists y G(x, y, f(x))$ YES
- ▶ $\exists y G(f(x), y, f(f(x)))$ YES
- ▶ $\exists y G(y, y, f(y))$ NO. Why not?

Suppose we know $\forall x G(x, y, f(x))$. Can we deduce:

- ▶ $G(y, y, f(y))$ YES. Why?

∃-INTRODUCTION (EXISTENTIAL GENERALISATION)

If a property holds a particular thing, then it holds for something:

$$\frac{S \vdash F}{S \vdash \exists x F[x/t]} \quad \text{where } x \text{ is free to replace } t \text{ in } F$$

Examples:

\exists -INTRODUCTION (EXISTENTIAL GENERALISATION)

If a property holds a particular thing, then it holds for something:

$$\frac{S \vdash F}{S \vdash \exists x F[x/t]} \quad \text{where } x \text{ is free to replace } t \text{ in } F$$

Examples:

- From $\forall y P(c, y)$ we can deduce $\exists x \forall y P(x, y)$

∃-INTRODUCTION (EXISTENTIAL GENERALISATION)

If a property holds a particular thing, then it holds for something:

$$\frac{S \vdash F}{S \vdash \exists x F[x/t]} \quad \text{where } x \text{ is free to replace } t \text{ in } F$$

Examples:

- ▶ From $\forall y P(c, y)$ we can deduce $\exists x \forall y P(x, y)$
- ▶ From $\forall y P(2, y)$ we can deduce $\exists x \forall y P(x, y)$

∃-INTRODUCTION (EXISTENTIAL GENERALISATION)

If a property holds a particular thing, then it holds for something:

$$\frac{S \vdash F}{S \vdash \exists x F[x/t]} \quad \text{where } x \text{ is free to replace } t \text{ in } F$$

Examples:

- ▶ From $\forall y P(c, y)$ we can deduce $\exists x \forall y P(x, y)$
- ▶ From $\forall y P(2, y)$ we can deduce $\exists x \forall y P(x, y)$
- ▶ From $\forall y P(f(x, z), y)$ we can deduce any of these:
 - ▶ $\exists x \forall y P(x, y)$ (replaced term $f(x, z)$ with bound x)
 - ▶ $\exists x \forall y P(f(x, z), y)$
 - ▶ $\exists w \forall y P(f(w, z), y)$

\exists -INTRODUCTION (EXISTENTIAL GENERALISATION)

If a property holds a particular thing, then it holds for something:

$$\frac{S \vdash F}{S \vdash \exists x F[x/t]} \quad \text{where } x \text{ is free to replace } t \text{ in } F$$

Examples:

- ▶ From $\forall y P(c, y)$ we can deduce $\exists x \forall y P(x, y)$
- ▶ From $\forall y P(2, y)$ we can deduce $\exists x \forall y P(x, y)$
- ▶ From $\forall y P(f(x, z), y)$ we can deduce any of these:
 - ▶ $\exists x \forall y P(x, y)$ (replaced term $f(x, z)$ with bound x)
 - ▶ $\exists x \forall y P(f(x, z), y)$
 - ▶ $\exists w \forall y P(f(w, z), y)$
- ▶ But from $\forall y P(f(x, y), y)$ we cannot deduce $\exists x \forall y P(x, y)$
Why not?

\forall -INTRODUCTION (UNIVERSAL GENERALISATION)

If a property holds for an arbitrary thing, then it holds for all things:

$\frac{S \vdash F}{S \vdash \forall x F}$	x is not free in any of the premises used to deduce F , and x is not free in any wff constructed by \exists -elimination
---	--

We let x be an arbitrary but fixed element of the domain D . Next, we construct a proof that F is true for x . Then we can say that since x was arbitrarily chosen, it follows that F is true for any x of the domain.

However, if during our argument we made any assumptions about the value of x (i.e. if we used it in a formula inferred by \exists -elimination), then we can no longer claim x was arbitrarily chosen.)

\forall -INTRODUCTION (UNIVERSAL GENERALISATION)

Example of incorrect usage

Requirements:

- ▶ x is not free in any of the premises used to deduce F
- ▶ x is not free in any wff constructed by \exists -elimination

Premises	Line	Formula	Justification	References
1	1	$P(x)$	P	
1	2	$\forall xP(x)$	\forall -intro	1

x is free in premise 1, so we cannot justify line 2.

Example: Domain of natural numbers, $P(x) : x$ is prime

\forall -INTRODUCTION (UNIVERSAL GENERALISATION)

Example of incorrect usage

Requirements:

- ▶ x is not free in any of the premises used to deduce F
- ▶ x is not free in any wff constructed by \exists -elimination

Premises	Line	Formula	Justification	References
1	1	$\forall x \exists y P(x, y)$	P	
1	2	$\exists y P(x, y)$	\forall -elim	1
1	3	$P(x, c)$	\exists -elim	2
1	4	$\forall x P(x, c)$	\forall -intro	3

x was free in a wff constructed by \exists -elimination, therefore we cannot justify line 4.

Example: Domain of natural numbers, $P(x, y) : x < y$

FORMAL PROOF EXAMPLES (1)

All humans are mortal. Socrates is human. Hence, Socrates is mortal.

$$\forall x(H(x) \rightarrow M(x)), H(Soc) \vdash M(Soc)$$

Premises	Line	Formula	Justification	References
1	1	$\forall x(H(x) \rightarrow M(x))$	P	
2	2	$H(Soc)$	P	
1	3	$(H(Soc) \rightarrow M(Soc))$	\forall -elim	1
1,2	4	$M(Soc)$	MP	2,3

FORMAL PROOF EXAMPLES (2)

Prove $\forall xP(x), \exists xQ(x) \vdash \exists x(P(x) \wedge Q(x))$

P.	L.	Formula	Just.	Refs.
1	1	$\forall xP(x)$	P	
2	2	$\exists xQ(x)$	P	
2	3	$Q(c)$ (c is a new constant)	\exists -elim	2
1	4	$P(c)$ (c free to replace x)	\forall -elim	1
1,2	5	$(P(c) \wedge Q(c))$	Conj.	3,4
1,2	6	$\exists x(P(x) \wedge Q(x))$ (c free to replace x)	\exists -intro	5

The order of lines 3 and 4 is very important here. If we did the \forall -elim first, then we could not have used c in the \exists -elim, because it would not have been a new constant.

FORMAL PROOF EXAMPLES (3)

Prove

$$\forall x(A(x) \rightarrow B(x)), \forall x(B(x) \rightarrow C(x)) \vdash \forall x(A(x) \rightarrow C(x))$$

P.	L.	Formula	Just.	Refs.
1	1	$\forall x(A(x) \rightarrow B(x))$	P	
2	2	$\forall x(B(x) \rightarrow C(x))$	P	

FORMAL PROOF EXAMPLES (3)

Prove

$$\forall x(A(x) \rightarrow B(x)), \forall x(B(x) \rightarrow C(x)) \vdash \forall x(A(x) \rightarrow C(x))$$

P.	L.	Formula	Just.	Refs.
1	1	$\forall x(A(x) \rightarrow B(x))$	P	
2	2	$\forall x(B(x) \rightarrow C(x))$	P	
1	3	$(A(x) \rightarrow B(x))$	\forall -elim	1

FORMAL PROOF EXAMPLES (3)

Prove

$$\forall x(A(x) \rightarrow B(x)), \forall x(B(x) \rightarrow C(x)) \vdash \forall x(A(x) \rightarrow C(x))$$

P.	L.	Formula	Just.	Refs.
1	1	$\forall x(A(x) \rightarrow B(x))$	P	
2	2	$\forall x(B(x) \rightarrow C(x))$	P	
1	3	$(A(x) \rightarrow B(x))$	\forall -elim	1
2	4	$(B(x) \rightarrow C(x))$	\forall -elim	2

FORMAL PROOF EXAMPLES (3)

Prove

$$\forall x(A(x) \rightarrow B(x)), \forall x(B(x) \rightarrow C(x)) \vdash \forall x(A(x) \rightarrow C(x))$$

P.	L.	Formula	Just.	Refs.
1	1	$\forall x(A(x) \rightarrow B(x))$	P	
2	2	$\forall x(B(x) \rightarrow C(x))$	P	
1	3	$(A(x) \rightarrow B(x))$	\forall -elim	1
2	4	$(B(x) \rightarrow C(x))$	\forall -elim	2
1,2	5	$(A(x) \rightarrow C(x))$	HS	3,4

FORMAL PROOF EXAMPLES (3)

Prove

$$\forall x(A(x) \rightarrow B(x)), \forall x(B(x) \rightarrow C(x)) \vdash \forall x(A(x) \rightarrow C(x))$$

P.	L.	Formula	Just.	Refs.
1	1	$\forall x(A(x) \rightarrow B(x))$	P	
2	2	$\forall x(B(x) \rightarrow C(x))$	P	
1	3	$(A(x) \rightarrow B(x))$	\forall -elim	1
2	4	$(B(x) \rightarrow C(x))$	\forall -elim	2
1,2	5	$(A(x) \rightarrow C(x))$	HS	3,4
1,2	6	$\forall x(A(x) \rightarrow C(x))$	\forall -intro	5

- x was always free to replace x
- x was not free in any of the premises used to deduce $(A(x) \rightarrow C(x))$, nor was free in a wff constructed by \exists -elim.

FORMAL PROOF EXAMPLES (4)

Database, $S \vdash A(\text{Brown}, 66)$

P.	L.	Formula	Just.	Refs.
1	1	$CSG(2013, 1377660, 70)$	P	
2	2	$CSG(2013, 1377540, 66)$	P	
3	3	$CSG(2013, 1381660, 55)$	P	
4	4	$SN(1377540, \text{Brown})$	P	
5	5	$SN(1381660, \text{Liu})$	P	
6	6	$\forall s \forall g \forall n ((CSG(2013, s, g) \wedge SN(s, n)) \rightarrow A(n, g))$	P	
6	7	$\forall g \forall n ((CSG(2013, 1377540, g) \wedge SN(1377540, n)) \rightarrow A(n, g))$	\forall -elim	6
6	8	$\forall n ((CSG(2013, 1377540, 66) \wedge SN(1377540, n)) \rightarrow A(n, 66))$	\forall -elim	6
6	9	$((CSG(2013, 1377540, 66) \wedge SN(1377540, \text{Brown})) \rightarrow A(\text{Brown}, 66))$	\forall -elim	6
2,4	10	$(CSG(2013, 1377540, 66) \wedge SN(1377540, \text{Brown}))$	Conj	2,4
2,4,6	11	$A(\text{Brown}, 66)$	MP	9,10

FORMAL PROOF EXAMPLES (5)

Lewis Carroll's logic:

Babies are illogical. Nobody is despised who can manage a crocodile. Illogical persons are despised. Therefore, babies cannot manage crocodiles.

P.	L.	Formula	Just.	Refs.
1	1	$\forall x(B(x) \rightarrow I(x))$	P	
2	2	$\forall x(C(x) \rightarrow \neg D(x))$	P	
3	3	$\forall x(I(x) \rightarrow D(x))$	P	

TAUTOLOGIES

Tautologies of propositional logic are also tautologies in predicate logic, when the atomic formula is replaced with a wff of predicate logic. e.g.

- ▶ $(A \vee \neg A)$ is a tautology, so is $(\forall xP(x) \vee \neg \forall xP(x))$ (replacing A with $\forall xP(x)$)
- ▶ $(\neg P \vee (Q \rightarrow P))$ is a tautology, so is $(\neg \forall xP(x) \vee (\exists yQ(y) \rightarrow \forall xP(x)))$ (replacing P with $\forall xP(x)$ and Q with $\exists yQ(y)$)

TAUTOLOGIES

Tautologies of propositional logic are also tautologies in predicate logic, when the atomic formula is replaced with a wff of predicate logic. e.g.

- ▶ $(A \vee \neg A)$ is a tautology, so is $(\forall xP(x) \vee \neg \forall xP(x))$ (replacing A with $\forall xP(x)$)
- ▶ $(\neg P \vee (Q \rightarrow P))$ is a tautology, so is $(\neg \forall xP(x) \vee (\exists yQ(y) \rightarrow \forall xP(x)))$ (replacing P with $\forall xP(x)$ and Q with $\exists yQ(y)$)

Other tautologies involve manipulating quantifiers, e.g.

- ▶ $(\forall xP(x) \rightarrow \exists xP(x))$
- ▶ $(\forall xP(x) \rightarrow \neg \exists x \neg P(x))$
- ▶ $\forall x(P(x) \vee \neg P(x))$
- ▶ $(P(x, y) \rightarrow (\exists xP(x, y) \wedge \exists yP(x, y)))$

EXAMPLE

Prove $(\forall xP(x) \rightarrow \exists xP(x))$

Premises	Line	Formula	Justification	References
1	1		P	
	2			
	3			
	4			
	5			
	6			

EXAMPLE

Prove $(\forall xP(x) \rightarrow \neg\exists x\neg P(x))$

Premises	Line	Formula	Justification	References
1	1		P	
	2			
	3			
	4			
	5			
	6			

EXAMPLE

Prove $\forall x(P(x) \vee \neg P(x))$

Premises	Line	Formula	Justification	References
1	1		P	
	2			
	3			
	4			
	5			
	6			

EXAMPLE

Prove $(P(x, y) \rightarrow (\exists x P(x, y) \wedge \exists y P(x, y)))$

Premises	Line	Formula	Justification	References
1	1		P	
	2			
	3			
	4			
	5			
	6			