

Methods for Accelerated Convergence of the Electron Density in Density Functional Theory

Nick Woods

Supervisor: Dr. Phil Hasnip

March 2, 2017

Kohn-Sham (KS) DFT solves the following eigenequation to obtain occupied single particle KS orbitals (ψ_i)

$$H^{KS}[\rho]\psi_i = \epsilon_i\psi_i$$
$$\rho(\mathbf{r}) = \sum_{i \in \text{occupied}} |\psi_i|^2$$

The solutions $\{\psi_i\}$ depend on the input H^{KS} through $\rho(\mathbf{r})$

Introduction

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

$$F : \left\{ \begin{array}{c} \rho^{\text{in}} \rightarrow H^{KS}[\rho^{\text{in}}] \\ \downarrow \\ \text{Solve KS equation} \\ \downarrow \\ \rho^{\text{out}} \sim \sum |\psi|^2 \end{array} \right.$$

$$F[\rho^{\text{in}}] = \rho^{\text{out}}$$

$$\rho^{\text{in}} \neq \rho^{\text{out}}$$

Achieving Convergence

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Methods to achieve convergence:

Density Mixing – Find optimal form of

$$\rho_{n+1}^{\text{in}} = f(\{\rho_n^{\text{in}}, \rho_n^{\text{out}}\}) \quad n \in [1, \text{iteration history length}]$$

Preconditioners – Apply a preconditioning matrix P specific to problem to assist f

Density Mixing

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Simplest update one can imagine

$$\rho_{n+1}^{\text{in}} = F[\rho_n^{\text{in}}] = \rho_n^{\text{out}}$$

Fixed-point iteration

Analysis of this update highlights difficulties of the SCF process

Density Mixing

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

If $\rho = \rho^* + \delta\rho$ then the fixed-point iteration formula can be linearised,

$$\begin{aligned}\delta\rho_{n+1}^{\text{in}} &= \frac{\delta F}{\delta\rho} \delta\rho_n^{\text{in}} \\ &= \frac{\rho_n^{\text{out}}}{\rho_n^{\text{in}}} \delta\rho_n^{\text{in}} \\ &= M \delta\rho_n^{\text{in}}\end{aligned}$$

linear response

Density Mixing

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Convergence: $\delta\rho_{n+1}^{\text{in}} \rightarrow 0$ as $n \rightarrow \infty$

$$\implies (M)^n \delta\rho_1^{\text{in}} \rightarrow 0$$

What conditions required for M^n to tend to zero?

Diagonalising M in its orthonormal eigenbasis gives

$$((M^d)^n)_{ii} = (\lambda_i)^n |\mathbf{e}_i\rangle\langle\mathbf{e}_i|$$

$$\implies M\delta\rho_n^{\text{in}} \rightarrow 0 \text{ iff } |\lambda_i| < 1 \ \forall \ i$$

Density Mixing

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

In general not true!

Could imagine scaling M with a parameter α which guarantees this – *linear mixing*

The eigenvalues in this case $\sim \alpha\{|\lambda_i|\}$.

What is the problem with this?

Density Mixing

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Need a more sophisticated scheme than linear mixing or fixed-point iterations

CASTEP uses (variants of) Pulay method, multisecant Broyden (quasi-Newton)

Both need to be executed using limited memory (can't store $N_{\text{gridpoints}} \times N_{\text{gridpoints}}$ matrix) and prefer convergence in a low number of SCF cycles (diagonalising H^{KS} expensive)

Density Mixing

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

My project: investigate possible improvements

Robust mixing for ab initio quantum mechanical calculations

L. D. Marks and D. R. Luke

Phys. Rev. B 78, 075114 2008

A multisecant Broyden technique – general idea is to treat history of iterates as sampling phase space rather than a path to the solution

Preconditioners

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Can now think about specific problem (KS DFT) to investigate what makes a good preconditioner

In KS DFT, perturbations in density (like those induced in density mixing) propagate through F as such

$$\delta V_h(\mathbf{r}) = \int \frac{\delta \rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}',$$
$$\delta \rho(\mathbf{r}) = \int \chi^{\text{DFT}}(\mathbf{r}, \mathbf{r}') \delta V_h(\mathbf{r}') d\mathbf{r}'.$$

Preconditioners

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

$$\delta \tilde{V}_h(\mathbf{G}) \sim \frac{\delta \tilde{\rho}(\mathbf{G})}{|\mathbf{G}|^2}$$

Charge sloshing – unique to KS DFT (from Hartree potential)

Continuous overcorrection of V_h and ρ

Preconditioners

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Preconditioner should weight low \mathbf{G} vectors less, to avoid sloshing

It turns out that the preconditioning matrix, M from earlier, is closely related to the DFT dielectric of system, ϵ

Use dielectric model to precondition $\delta\rho$

CASTEP uses Kerker dielectric model (Jellium) – $\epsilon \sim \delta_{GG'} \frac{1}{G^2}$

Preconditioners


Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results



Preconditioners

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

My project: investigate better preconditioners than Kerker

First, a better dielectric model

New model dielectric function and exchange-correlation potential for semiconductors and insulators Zachary H. Levine and Steven G. Louie Phys. Rev. B 25, 6310 1982

Still isotropic – include anisotropy?

Preconditioners


Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results



Preconditioners

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

Both models include two parameters

How do they map onto each other?

What are the optimal parameters?

Converging
the Ground
State Electron
Density

Nick Woods

Introduction

Theory

Results

To come...