

QUIZ3.extre me_pts_tu...

MATB61 TUT03

Nick Huang Quiz3 Extreme point and extreme point theoremMar.4 2021 Week 8

Instuction: You will have 30 minutes to finish quiz and will have 5 minutes to submit the quiz to Crowdmark after the quiz has finished. You will need to keep your cameras on during the quiz and submission times. One hand-held calculator is allowed. Electronic devices, online calculators, notes and other aids are not allowed. Violation of the instruction can be considered as an academic misconduct, and will be reported to the instructor and the department immediately.

Question

Consider the following linear programming problem.

max
$$z = x_1 + 2x_2 - x_3$$

subject to
 $2x_1 - 6x_2 + 3x_3 \le 12$
 $2x_2 + x_3 \le 2$
 $x_1, x_2, x_3 \ge 0$

Find all the extreme points and find the optimal solution. (X11X21X3) = (0,0,0)

• Note: In order to receive full marks, you will need to do the computation correctly and justify every steps. In particular, explain why your solution is the optimal solution.

O FEOGRAPHO REGION IS bonded and non-empty, because DEXIE 9 OEXZEI, OE X3 E Z, and by previocitating theorem, optimal

DEXZET,
$$0 \le x_3 \le 2$$
, and by extremethant thousands of the solution which is a basic feasitive solution

2) Convenient form, max $x = C^Tx$ where $x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$, $c = \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}$, $A = \begin{pmatrix} 2 - 6 & 3 & 10 \\ 0 & 2 & 10 & 1 \end{pmatrix}$, $b = \begin{pmatrix} 12 \\ 2 \\ 2 \end{pmatrix}$

Max exprises $x > 0$.

(i)
$$\begin{pmatrix} 0 & 2 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 1 \end{pmatrix} + \lambda = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 0 & 2 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 2 & 3 \end{pmatrix} \times \begin{pmatrix} 0 & 1 & 2 \\ 2$$

$$(7) \left(\frac{3}{10} \frac{1}{2} \right) \sim \left(\frac{10}{01} \frac{2}{6} \right), \ X = \left(\frac{9}{6} \right)$$

$$(8) \left(\frac{30}{11} \frac{12}{2} \right) \sim \left(\frac{10}{01} \frac{4}{-2} \right), \ X = \left(\frac{9}{4} \right)$$

$$(9) \begin{pmatrix} 0 & 12 \\ 0 & 2 \end{pmatrix}, \quad \chi = \begin{pmatrix} 0 \\ 0 \\ 12 \\ 2 \end{pmatrix}$$

$$(9) \begin{pmatrix} 10 & 12 \\ 2 \\ 2 \end{pmatrix}$$

$$(9) \begin{pmatrix} 10 & 12 \\ 2 \\ 2 \end{pmatrix}$$

$$(9) \begin{pmatrix} 10 & 12 \\ 2 \\ 2 \end{pmatrix}$$

@ Bosic feosible solution, check x 7,0

$$X = \begin{pmatrix} 9 \\ 0 \\ 0 \end{pmatrix} \text{ with } Z = 11 \text{ , } \qquad X = \begin{pmatrix} 3 \\ 2 \\ 0 \\ 0 \end{pmatrix} \text{ with } Z = 6$$

$$X = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \text{ with } Z = 2 \text{ , } \qquad X = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \text{ with } Z = 6$$

$$X = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \text{ with } Z = 2 \text{ , } \qquad X = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \text{ with } Z = 0$$

Therefore, optimal solution exists at x = () with

optimal value 2=11 to the other maximization protolem,

ty reasoning to 1.