



Example_m
atrix_repr...

MATB24 TUT5
Nick Huang Examples: Matrix representation and coordinate vector May.30 2021

Definitions

You should always check the lecture note and the textbook for the definition, and make sure the definitions are consistent.

1. We call $\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$ the coordinate vector of $v \in V$ with respect to the basis (v_1, \dots, v_n) of V if $v = a_1 v_1 + \dots + a_n v_n$ for some $a_1, \dots, a_n \in \mathbb{F}$. Denoted as $[v]_{v_1, \dots, v_n}$

2. Let V, W be vector spaces over \mathbb{F} . Let $T: V \rightarrow W$ be a linear transformation. Let $\alpha = (v_1, \dots, v_n)$ and $\beta = (w_1, \dots, w_m)$ be ordered basis of V and W respectively. Then the matrix representation of T with respect to α and β is

$$[T]_{w_1, \dots, w_m}^{\alpha} = \begin{bmatrix} [T(v_1)]_{w_1, \dots, w_m} & \dots & [T(v_n)]_{w_1, \dots, w_m} \end{bmatrix}$$

Remark. When the question asks you the find the matrix of a given linear transformation $T: V \rightarrow W$ with respect to some bases as we have seen in the assignment, you are NOT asked to find a matrix A , such that $T(v) = Av$. You are asked to find the matrix representation as defined. While sometimes it is true that the matrix representation, let's call it A , will result $T(v) = Av$ for all $v \in V$, this is not always true. The Av is a matrix multiplication and we have seen vectors that are not vectors in \mathbb{R}^n .

Examples: Coordinate vector

1. We will start with the familiar one \mathbb{R}^n . In next few questions, consider the vector space $V = \mathbb{R}_4$ with the usual operations.
- (a) Circle the elements which belong the vector space V
- $\textcircled{1} (1, 0, 0, 0)^T$ $\textcircled{2} (2, 9, 0, 5)^T$ $\textcircled{3} (1, i, 0, 0)^T$ $\textcircled{4} (3, 4, 0)^T$ $\textcircled{5} (1, 4, 0, 0)^T$
- $\textcircled{1} \rightarrow 1+x$ $\textcircled{2} \rightarrow (1, x, x^2, x^3)^T$ $\textcircled{3} \rightarrow 3+x^2+x^3$ $\textcircled{4} \rightarrow 4x+i$ $\textcircled{5} \rightarrow \begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix}$
- (b) What is the standard (ordered) basis for V ? Call it α .
- (c) We know that α spans the vector space because it is a basis. For the elements above that are in the vector space, write those as linear combinations of the vectors in the basis. Find $[v]_\alpha$
- (d) Known $\beta = ((1, 1, 1, 1)^T, (1, 1, 1, 0)^T, (1, 1, 0, 0)^T, (1, 0, 0, 0)^T)$ is an ordered basis of V . Repeat the previous subquestion with β .

(b) $\alpha = \left(\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \right)$

(c) $\textcircled{1}: \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = 1 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \right]_\alpha = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$

$\textcircled{2}: \begin{pmatrix} 2 \\ 9 \\ 0 \\ 5 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + 9 \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 2 \\ 9 \\ 0 \\ 5 \end{pmatrix} \right]_\alpha = \begin{pmatrix} 2 \\ 9 \\ 0 \\ 5 \end{pmatrix}$

$\textcircled{3}: \begin{pmatrix} 1 \\ i \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = 1 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 1 \\ i \\ 0 \\ 0 \end{pmatrix} \right]_\alpha = \begin{pmatrix} 1 \\ i \\ 0 \\ 0 \end{pmatrix}$

(d) $\beta = \left(\begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \right)$

$\textcircled{1}: \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} + 0 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + 1 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \right]_\beta = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$

$\textcircled{2}: \begin{pmatrix} 2 \\ 9 \\ 0 \\ 5 \end{pmatrix} = 5 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} + 4 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + (-1) \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 2 \\ 9 \\ 0 \\ 5 \end{pmatrix} \right]_\beta = \begin{pmatrix} 5 \\ 4 \\ 2 \\ -1 \end{pmatrix}$

$\textcircled{3}: \begin{pmatrix} 1 \\ i \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} + 0 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + 1 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 1 \\ i \\ 0 \\ 0 \end{pmatrix} \right]_\beta = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$

2. In the next few questions, consider the vector space $V = \mathbb{P}_3(\mathbb{R})$ with the usual operations.
- (a) Circle the elements which belong the vector space V
- $\textcircled{1} 1+x$ $\textcircled{2} (1, x, x^2, x^3)^T$ $\textcircled{3} 3+x^2+x^3$ $\textcircled{4} 4x+i$ $\textcircled{5} \begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix}$
- (b) What is the standard (ordered) basis for V ? Call it α .
- (c) We know that α spans the vector space because it is a basis. For the elements above that are in the vector space, write those as linear combinations of the vectors in the basis. Find $[v]_\alpha$
- (d) One can check that $\beta = (2, 1+x, 1+x+x^2, 1+x+x^2+x^3)$ is an ordered basis of V . Repeat the previous subquestion with β .

(b) $\alpha = (1, x, x^2, x^3)$

(c) $\textcircled{1}: 1+x = 1 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3, \text{ st } [1+x]_\alpha = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}$

$\textcircled{2}: 3+x^2+x^3 = 3 \cdot 1 + 0 \cdot x + 1 \cdot x^2 + 1 \cdot x^3, \text{ st } [3+x^2+x^3]_\alpha = \begin{pmatrix} 3 \\ 0 \\ 1 \\ 1 \end{pmatrix}$

(d) $\beta = (2, 1+x, 1+x+x^2, 1+x+x^2+x^3)$

$\textcircled{1}: 1+x = 0 \cdot (1+x) + 0 \cdot (1+x+x^2) + 0 \cdot (1+x+x^2+x^3) + 1 \cdot (2), \text{ st } [1+x]_\beta = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$

$\textcircled{2}: 3+x^2+x^3 = 1 \cdot (1+x) + 0 \cdot (1+x+x^2) + 0 \cdot (1+x+x^2+x^3) + 2 \cdot (2), \text{ st } [3+x^2+x^3]_\beta = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 2 \end{pmatrix}$

3. In the next few questions, consider the vector space $V = M_2(\mathbb{R})$ with the usual operations.
- (a) Circle the elements which belong the vector space V
- $\textcircled{1} 1+x$ $\textcircled{2} (1, x, x^2, x^3)^T$ $\textcircled{3} (1, i, 0, 0)^T$ $\textcircled{4} (3, 4, 0)^T$ $\textcircled{5} \begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix}$
- (b) What is the standard (ordered) basis for V ? Call it α .
- (c) We know that α spans the vector space because it is a basis. For the elements above that are in the vector space, write those as linear combinations of the vectors in the basis. Find $[v]_\alpha$
- (d) One can check that $\beta = \left(\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \right)$ is an ordered basis of V . Repeat the previous subquestion with β .

(b) $\alpha = \left(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right)$

(c) $\textcircled{1}: \begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix} = 1 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + 4 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + 0 \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + 1 \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix} \right]_\alpha = \begin{pmatrix} 1 \\ 4 \\ 0 \\ 1 \end{pmatrix}$

(d) $\begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix} = 1 \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} + 1 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + 4 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} - 3 \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \text{ st } \left[\begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix} \right]_\beta = \begin{pmatrix} -3 \\ 1 \\ 4 \\ 0 \end{pmatrix}$

Examples: Matrix representation of linear transformation

1. In the next few questions, consider the following linear transformation.

$T: V \rightarrow W$ by $T\left(\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}\right) = \begin{pmatrix} v_1 + 2v_2 \\ v_2 \end{pmatrix}$

where $V = \mathbb{R}^2, W = \mathbb{R}^3$

- (a) Compute $T\left(\begin{pmatrix} 2 \\ 3 \end{pmatrix}\right)$ $T\left(\begin{pmatrix} 2 \\ 3 \end{pmatrix}\right) = \begin{pmatrix} 2+2 \cdot 3 \\ 3 \end{pmatrix} = \begin{pmatrix} 8 \\ 3 \end{pmatrix}$
- (b) Let α and β be the standard bases of V and W respectively. Find the matrix representation $[T]_\beta^\alpha$ by our definition and notation.
- (c) Let $\gamma = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}\right)$ which is another ordered basis of V . Find the matrix representation $[T]_\beta^\gamma$ by our definition and notation.
- (d) Let $\delta = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}\right)$ which is another ordered basis of W . Find the matrix representation $[T]_\delta^\alpha$ by our definition and notation.

Notice $T\left(\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}\right) = \begin{pmatrix} v_1 + 2v_2 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$

(b) $\alpha = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}\right), \beta = \left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}\right)$

$[T]_\beta^\alpha = \begin{pmatrix} [T\left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right)]_\beta & [T\left(\begin{pmatrix} 0 \\ 1 \end{pmatrix}\right)]_\beta \end{pmatrix} = \begin{pmatrix} \left[\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right]_\beta & \left[\begin{pmatrix} 2 \\ 1 \end{pmatrix}\right]_\beta \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$

(c) $\gamma = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}\right)$ is an ordered basis of V

$[T]_\beta^\gamma = \begin{pmatrix} [T\left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right)]_\beta & [T\left(\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right)]_\beta \end{pmatrix} = \begin{pmatrix} \left[\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right]_\beta & \left[\begin{pmatrix} 3 \\ 1 \end{pmatrix}\right]_\beta \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}$

(d) $\delta = \left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}\right)$ is an ordered basis of W

$[T]_\delta^\alpha = \begin{pmatrix} [T\left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right)]_\delta & [T\left(\begin{pmatrix} 0 \\ 1 \end{pmatrix}\right)]_\delta \end{pmatrix} = \begin{pmatrix} \left[\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right]_\delta & \left[\begin{pmatrix} 2 \\ 1 \end{pmatrix}\right]_\delta \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$

2. In the next few questions, consider the following linear transformation.

$T: V \rightarrow W$ by $T(f) = \begin{pmatrix} f'(2) \\ f(0) \end{pmatrix}$

where $V = \mathbb{P}_2(\mathbb{R}), W = \mathbb{R}^2$

- (a) Compute $T(1+x)$ and $T(x^2+x^3)$ $T(1+x) = \begin{pmatrix} 1 \\ 1+x=0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, T(x^2+x^3) = \begin{pmatrix} 2x+3x^2 \\ x^2+x^3 \end{pmatrix}_{x=0} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$
- (b) Let α and β be the standard bases of V and W respectively. Find the matrix representation $[T]_\beta^\alpha$ by our definition and notation.
- (c) Let $\gamma = (1, 1+x, 1+x+x^2)$ which is another ordered basis of V . Find the matrix representation $[T]_\beta^\gamma$ by our definition and notation.
- (d) Let $\delta = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}\right)$ which is another ordered basis of W . Find the matrix representation $[T]_\delta^\alpha$ by our definition and notation.

(b) $\alpha = (1, x^2), \beta = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}\right)$

$[T]_\beta^\alpha = \begin{pmatrix} [T(1)]_\beta & [T(x^2)]_\beta \end{pmatrix} = \begin{pmatrix} \left[\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right]_\beta & \left[\begin{pmatrix} 0 \\ 0 \end{pmatrix}\right]_\beta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$

(c) $\gamma = (1, 1+x, 1+x+x^2)$

$[T]_\beta^\gamma = \begin{pmatrix} [T(1)]_\beta & [T(1+x)]_\beta & [T(1+x+x^2)]_\beta \end{pmatrix} = \begin{pmatrix} \left[\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right]_\beta & \left[\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right]_\beta & \left[\begin{pmatrix} 5 \\ 1 \end{pmatrix}\right]_\beta \end{pmatrix} = \begin{pmatrix} 1 & 1 & 5 \\ 0 & 0 & 1 \end{pmatrix}$

(d) $\delta = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}\right)$

$[T]_\delta^\alpha = \begin{pmatrix} [T(1)]_\delta & [T(1+x)]_\delta & [T(1+x+x^2)]_\delta \end{pmatrix} = \begin{pmatrix} \left[\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right]_\delta & \left[\begin{pmatrix} 1 \\ 1 \end{pmatrix}\right]_\delta & \left[\begin{pmatrix} 5 \\ 1 \end{pmatrix}\right]_\delta \end{pmatrix} = \begin{pmatrix} -1 & 0 & 4 \\ 1 & 1 & 1 \end{pmatrix}$