

MT5763_2_220021614

Nico Herrig

2022-10-15

```
library(tidyverse)

## -- Attaching packages ----- tidyverse 1.3.2 --
## v ggplot2 3.3.6      v purrr 0.3.4
## v tibble 3.1.8       v dplyr 1.0.10
## v tidyr 1.2.1        v stringr 1.4.1
## v readr 2.1.2        v forcats 0.5.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
```

Problem 1

Description: The problem presents two variables X and Y , where $X \sim N(\mu = 4, \sigma^2 = 10)$ and $Y \sim U(a = 2, b = 8)$.

Compute $Pr(X > Y)$ and use bootstrapping to derive the sampling distribution for your estimate of $Pr(X > Y)$. Show how the sample variance of this sampling distribution changes as a function of the number of Monte Carlo simulations.

For the underlying problem, a sample containing 10000 random deviates from $X \sim N(\mu = 4, \sigma^2 = 10)$ and $Y \sim U(a = 2, b = 8)$ is used. To simulate “real-world conditions”, the solution is obtained from only the below given vectors for X and Y .

```
set.seed(0911)

X <- rnorm(10000, mean = 4, sd = sqrt(10))
Y <- runif(10000, min = 2, max = 8)

Pr_hat <- sum(X > Y) / 10000

print(Pr_hat)
```

```
## [1] 0.3852
```

Calculating $\bar{Pr}(X > Y)$ from the initial sample without any further methods, we derive a value of 0.3852. A non-parametric bootstrapping is now used to simulate the distribution of $\bar{Pr}(X > Y)$ from the given sample.

```
bootstrap <- function(n_bootstraps, vec1 = X, vec2 = Y) {

  vector_prob <- rep(NA, times = n_bootstraps)

  for (i in 1 : n_bootstraps) {
    X_resampled <- vec1[sample(1 : length(vec1), length(vec1), replace = TRUE)]
    Y_resampled <- vec2[sample(1 : length(vec2), length(vec2), replace = TRUE)]


    vector_prob[i] <- sum(X_resampled > Y_resampled) / length(X)
  }
  return(vector_prob)
}
```

The bootstrap algorithm re-samples the vectors X and Y *with replacement* and calculates the resulting $\bar{Pr}_i(X > Y)$, and repeating this procedure n times. The function puts out a vector with n generated probabilities.

Using the bootstrap function, we now can evaluate the distribution for $Pr(X > Y)$

```
df_probabilities <- data.frame(p_hat =bootstrap(n_bootstraps = 10000))

df_probabilities %>%
  ggplot(mapping = aes(p_hat)) +
    geom_density()
```



```
MC_sim <- function(runs) {  
}
```