

Yuji Tanaka

Division of Exploratory Research, Institute of Medical Science, University of Tokyo, Tokyo, Japan

Tomohiro Sawa and Yoshinori Nakata

Medical Information and System Research Center, Teikyo University School of Medicine, Tokyo, Japan

ACKNOWLEDGMENT

We are grateful to Toshiteru Nobue (Maikai Solutions, Tokyo, Japan) for his technical support. Supported by Grant from Health and Labor Sciences Research Grants, the third-term comprehensive control for cancer (H18-012 to Dr Nakata) from the Ministry of Health Welfare and Labor of the Japanese Government, Tokyo, Japan.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The author(s) indicated no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

Conception and design: Hiroto Narimatsu, Masahiro Kami, Tomohiro Sawa, Yoshinori Nakata

Collection and assembly of data: Hiroto Narimatsu, Tomoko Matsumura, Tomohiro Morita, Yukiko Kishi, Koichiro Yuji, Tsunehiko Komatsu, Yuji Tanaka

Data analysis and interpretation: Hiroto Narimatsu, Masahiro Kami, Tomohiro Sawa, Yoshinori Nakata

Manuscript writing: Hiroto Narimatsu, Masahiro Kami, Yoshinori Nakata

Final approval of manuscript: Hiroto Narimatsu, Tomoko Matsumura, Tomohiro Morita, Yukiko Kishi, Koichiro Yuji, Masahiro Kami, Tsunehiko Komatsu, Yuji Tanaka, Tomohiro Sawa, Yoshinori Nakata

REFERENCES

1. Newnham GM, Burns WI, Snyder RD, et al: Information from the Internet: Attitudes of Australian oncology patients. *Intern Med J* 36:718-723, 2006
2. Overture. <http://inventory.jp.overture.com/d/searchinventory/suggestion/>
3. Yahoo! Japan. Homepage. <http://www.yahoo.co.jp/>
4. Nikkei BP Consulting Inc: Access analysis. <http://consult.nikkeibp.co.jp/consult/bizmark/feature.html>
5. Alper BS: Usefulness of online medical information. *Am Fam Physician* 74:482-485, 2006
6. Berland GK, Elliott MN, Morales LS, et al: Health information on the internet: Accessibility, quality, and readability in English and Spanish. *JAMA* 285:2612-2621, 2001
7. D'Alessandro DM, Kingsley P, Johnson-West J: The readability of pediatric patient education materials on the World Wide Web. *Arch Pediatr Adolesc Med* 155:807-812, 2001
8. Morita T, Narimatsu H, Matsumura T, et al: A study of cancer information for cancer patients on the internet. *Int J Clin Oncol* 12:440-447, 2007
9. MSN Japan: Homepage. <http://jp.msn.com/>
10. Google Japan: Homepage. <http://www.google.co.jp/>

DOI: 10.1200/JCO.2008.18.7468

Cetuximab Pharmacokinetics in End-Stage Kidney Disease Under Hemodialysis

TO THE EDITOR: Cetuximab, an anti-epidermal growth factor receptor chimeric mouse/human immunoglobulin 1 monoclonal antibody against the epidermal growth factor (Merck, Darmstadt, Germany), has been approved as a treatment for advanced head and neck cancer in combination with radiation therapy.^{1,2} However, there is very little data on cetuximab in patients undergoing chronic dialysis.³ The treatment of cancer in patients with impaired renal function is an emerging problem because the population is getting older and the rate of chronic dialysis increases by 5% yearly in Western countries. We report a pharmacokinetic study of cetuximab in a patient with renal insufficiency requiring hemodialysis. Cetuximab was instituted at a dose of 250 mg/kg weekly for a 55-year-old patient with head and neck cancer.

We characterized the pharmacokinetics and efficacy of cetuximab at conventional efficacious dose levels in combination with radiation therapy in a hemodialyzed patient with head and neck cancer. The aim of the study was to determine whether conventional doses of cetuximab in combination with radiotherapy were appropriate for hemodialyzed patients.

Cetuximab serum concentration was measured by a validated enzyme-linked immunosorbent assay. The enzyme-linked immunosorbent assay method used a recombinant human epidermal growth factor receptor (extracellular domain) adsorbed onto microtiter plates to capture cetuximab in serum. The captured cetuximab was detected using a peroxidase-conjugated goat antihuman F(ab')₂ specific for Fc

fragment (horseradish peroxidase anti-human immunoglobulin G). Lower limit of quantitation and upper limit of quantitation were 0.75 and 15 µg/mL, respectively. The limit of detection was 0.012 µg/mL. Concentrations higher than the upper limit of quantification were diluted 1:10 or 1:100, deviation and variability of this procedure being lower than 4.5%. Serum samples were used to estimate cetuximab pharmacokinetics, assuming no time-dependence, with WINNonlin (Scientific Consultant, Apex, NC; Pharsight Corporation). One- and two-compartment models with first order distribution and elimination constants were tested. The best model was selected using the usual methods, which included the analysis of plots of observed versus predicted concentrations and the Akaike information criteria. The model that best fitted the observed data was a two-compartment model with first-order elimination from the central compartment (Fig 1). Clearance from central compartment was 0.025 L/h, central compartment volume was 3.8 L, and terminal elimination half-life was 11.9 days (Table 1).

Although analyses of cetuximab pharmacokinetics were previously reported, the results obtained in our patient cannot be readily compared with these publications. Tan et al⁴ did not use a formal compartment model. In the studies of Baselga et al⁵ and Delbaldo et al,⁶ cetuximab pharmacokinetics were described by a one-compartment model. However, a two-compartment model has previously been shown to be the best to describe the pharmacokinetics of immunoglobulin 1 monoclonal antibodies, including trastuzumab,⁷ inolimomab,⁸ rituximab,⁹ basiliximab,¹⁰ clenoliximab,¹¹ alemtuzumab,¹² and adalimumab.¹³ Dirks et al¹⁴ used a two-compartment model but with a Michaelis-Menten type of elimination. This last approach necessitates a large number of patients and the study of different dose regimens, and could not be applied to our patient.




Fig 1. Observed (blue dots) and predicted (continuous lines) cetuximab concentrations over time. The correlation between predicted and observed concentrations are presented in the insert.

The longer half-life estimated in our patient, as compared with the studies of Baselga et al⁵ and Delbaldo et al,⁶ may be explained by their use of a one-compartment model, which is known to underestimate the elimination half-life when a two-compartment model is needed. The concentrations measured in our patient were clinically effective, because tumor response was more than 70%. In contrast, his concentrations did not lead to a significant dose-dependent toxicity because only xeroderma and grade 1 acneiform rash were observed. Clinical response to cetuximab was previously reported to increase with the degree of cutaneous toxicity and a relationship was found between cetuximab trough concentrations and clinical response.¹⁵

Although, as indicated earlier, our results cannot be directly compared with the literature, the pharmacokinetic parameters estimated in our hemodialyzed patient are similar to those reported for other monoclonal antibodies in patients with normal kidney function. A lack of influence of hemodialysis was described for other monoclonal antibodies, such as bevacizumab.¹⁶ Inauen et al³ observed no effect of hemodialysis on cetuximab concentrations but the small number of blood samples did not allow them to estimate cetuximab pharmacokinetic parameters. These preliminary data indicate that treating dialyzed patients with monoclonal antibodies as a cancer therapy may be done safely. Further prospective studies are warranted to elucidate the pharmacokinetics and pharmacodynamics of monoclonal antibodies in patients with end-stage kidney disease. This report suggests that previously used pharmacokinetics models may not be appropriate.

Another important implication for the practicing oncologist is that hemodialyzed patients, who now have prolonged survival, should not be denied optimal treatment with monoclonal antibodies on the sole basis on their renal replacement. Translational research is needed to elucidate the pharmacokinetics and pharmacodynamics at the level of the normal and damaged kidney.

Juliette Thariat

Centre Lacassagne, Centre National de la Recherche Scientifique-UMR6543, Nice, France

Nicolas Azzopardi

Centre Hospitalier Régional Universitaire, Centre National de la Recherche Scientifique-UMR6239, Tours, France

Frederic Peyrade

Centre Lacassagne, Centre National de la Recherche Scientifique-UMR6543, Nice, France

Vincent Launay-Vacher

Centre Hospitalier Régional Universitaire, Pitié Salpêtrière, Paris, France

José Santini

Centre Hospitalier Universitaire de Pasteur, Nice, France

Thierry Lecomte

Centre Hospitalier Régional Universitaire, Centre National de la Recherche Scientifique-UMR6239, Tours, France

Marie-Christine Etienne-Grimaldi

Centre Lacassagne, Centre National de la Recherche Scientifique-UMR6543, Nice, France

Gilles Paintaud

Centre Hospitalier Régional Universitaire, Centre National de la Recherche Scientifique-UMR6239, Tours, France

Gerard Milano

Centre Lacassagne, Centre National de la Recherche Scientifique-UMR6543, Nice, France

ACKNOWLEDGMENT

We thank Sophie Dubbiosi, Merck, France, for independent logistic support.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The author(s) indicated no potential conflicts of interest.

Table 1. Estimated Pharmacokinetic Parameters

Parameter	Abbreviation	Estimated Value
Volume of central compartment, L	V1	3.8
Clearance from the central compartment, L/hour	CL	0.025
Volume of peripheral compartment, L	V2	3.77
Intercompartment clearance, L/hour	CLD2	0.022
Distribution half-life, days	$T_{1/2\alpha}$	1.84
Elimination half-life, days	$T_{1/2\beta}$	11.90
Mean residence time, days	MRT	12.62
Volume of distribution at steady state, L	Vss	7.56

REFERENCES

1. Bonner JA, Harari PM, Giralt J, et al: Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *N Engl J Med* 354:567-578, 2006
2. Thariat J, Milas L, Ang KK: Integrating radiotherapy with epidermal growth factor receptor antagonists and other molecular therapeutics for the treatment of head and neck cancer. *Int J Radiat Oncol Biol Phys* 69:974-984, 2007
3. Inauen R, Cathomas R, Boehm T, et al: Feasibility of using cetuximab and bevacizumab in a patient with colorectal cancer and terminal renal failure. *Oncology* 72:209-210, 2007
4. Tan AR, Moore DF, Hidalgo M, et al: Pharmacokinetics of cetuximab after administration of escalating single dosing and weekly fixed dosing in patients with solid tumors. *Clin Cancer Res* 12:6517-6522, 2006
5. Baselga J, Pfister D, Cooper MR, et al: Phase I studies of anti-epidermal growth factor receptor chimeric antibody C225 alone and in combination with cisplatin. *J Clin Oncol* 18:904-914, 2000
6. Delbaldo C, Pierga JY, Dieras V, et al: Pharmacokinetic profile of cetuximab (Erbix) alone and in combination with irinotecan in patients with advanced EGFR-positive adenocarcinoma. *Eur J Cancer* 41:1739-1745, 2005
7. Bruno R, Washington CB, Lu JF, et al: Population pharmacokinetics of trastuzumab in patients with HER2+ metastatic breast cancer. *Cancer Chemother Pharmacol* 56:361-369, 2005
8. Dartois C, Freyer G, Michallet M, et al: Exposure-effect population model of inolimomab, a monoclonal antibody administered in first-line treatment for acute graft-versus-host disease. *Clin Pharmacokinet* 46:417-432, 2007
9. Regazzi MB, Iacona I, Avanzini MA, et al: Pharmacokinetic behavior of rituximab: A study of different schedules of administration for heterogeneous clinical settings. *Ther Drug Monit* 27:785-792, 2005
10. Kovarik JM, Nashan B, Neuhaus P, et al: A population pharmacokinetic screen to identify demographic-clinical covariates of basiliximab in liver transplantation. *Clin Pharmacol Ther* 69:201-209, 2001
11. Mould DR, Davis CB, Minthorn EA, et al: A population pharmacokinetic-pharmacodynamic analysis of single doses of clenoliximab in patients with rheumatoid arthritis. *Clin Pharmacol Ther* 66:246-257, 1999
12. Mould DR, Baumann A, Kuhlmann J, et al: Population pharmacokinetics-pharmacodynamics of alemtuzumab (Campath) in patients with chronic lymphocytic leukaemia and its link to treatment response. *Br J Clin Pharmacol* 64:278-291, 2007
13. Weisman MH, Moreland LW, Furst DE, et al: Efficacy, pharmacokinetic, and safety assessment of adalimumab, a fully human anti-tumor necrosis factor- α monoclonal antibody, in adults with rheumatoid arthritis receiving concomitant methotrexate: A pilot study. *Clin Ther* 25:1700-1721, 2003
14. Dirks NL, Nolting A, Kovar A, et al: Population pharmacokinetics of cetuximab in patients with squamous cell carcinoma of the head and neck. *J Clin Pharmacol* 48:267-278, 2008
15. Fracasso PM, Burris III H, Arquette MA, et al: A phase 1 escalating single-dose and weekly fixed-dose study of cetuximab: Pharmacokinetic and pharmacodynamic rationale for dosing. *Clin Cancer Res* 13:986-993, 2007
16. Garnier-Viougat N, Rixe O, Paintaud G, et al: Pharmacokinetics of bevacizumab in haemodialysis. *Nephrol Dial Transplant* 22:975, 2007

DOI: 10.1200/JCO.2008.18.7674

Pharmacologic Intervention for Cancer-Related Dyspnea

TO THE EDITOR: We have read with great interest the work carried out by Ben-Aharon et al¹, who conducted a systematic review of randomized controlled trials assessing all pharmacologic and non-pharmacologic intervention for dyspnea palliation in patients with cancer. We would like to make some comments.

It is surprising that in the search terms for PubMed, chlorpromazine was not included (decreases breathlessness without affecting ventilation or producing sedation in healthy subjects).² In an open-labeled trial, McIver et al³ found chlorpromazine effective for relief of dyspnea in advanced cancer.

The authors do not explain, within the study population, if those patients with cancer and other comorbidity (mainly chronic obstructive pulmonary disease) are excluded. We believe that this information is important, because if there is comorbidity, we can not discern if the cause of breathlessness was cancer or chronic obstructive pulmonary disease. In the longitudinal study conducted by our group,⁴ there were two or more causes to explain the etiology of dyspnea at the end of life (both the equal importance).

As the authors note, the single trial that assessed the role of midazolam as adjunct therapy⁵ to morphine, demonstrated that the beneficial effects of morphine in controlling baseline levels of dyspnea could be improved by the addition of midazolam, without increased somnolence. However, no other benzodiazepines were assessed in randomized controlled trials. Moreover, there are studies in non-oncological terminal respiratory illness, which suggest the possibility of an improvement in terminal breathlessness treated with other benzodiazepines.⁶

We think it is important to emphasize in the final discussion, one aspect that is essential for clinicians, and reflecting a common practice

in our midst: our terminally ill patients with cancer who are already receiving regular doses of opioids have almost equal benefit from a supplemental equivalent of 25% to 50% of their 4-hourly oral or suncutaneous regular opioid dose, to improve their breathlessness.⁷

At present, our group is conducting a phase II randomized controlled trial to determine the effectiveness of oral transmucosal fentanyl citrate for the exertion dyspnea treatment in seriously ill patients with cancer and to contribute to drug security data in that population.

Miguel Ángel Cuervo Pinna, Rafael Mota Vargas, María José Redondo Moralo, and Miguel Ángel Sánchez Correás

Regional Palliative Care Program of Extremadura, Palliative Care Team of Badajoz, Spain

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The author(s) indicated no potential conflicts of interest.

REFERENCES

1. Ben-Aharon I, Gafter-Gvili A, Paul M, et al: Interventions for alleviating cancer-related dyspnea: A systematic review. *J Clin Oncol* 26:2396-2404, 2008
2. O'Neil PA, Morton PB, Stara RD: Chlorpromazine: A specific effect on breathlessness? *Br J Clin Pharmacol* 19:793-797, 1985
3. McIver B, Walsh D, Nelson K: The use of chlorpromazine for symptom control in dying cancer patients. *J Pain Symptom Manage* 9:341-345, 1994
4. Cuervo Pinna MA, Mota Vargas R, Redondo Moralo MJ, et al: Etiology factors of the dyspnea at the end of life. *Med Pal* 15:143-148, 2008
5. Navigante AH, Cerchietti LC, Castro MA, et al: Midazolam as adjunct therapy to morphine in the alleviation of severe dyspnea perception in patients with advanced cancer. *J Pain Symptom Manage* 31:38-47, 2006
6. Man GC, Hsu K, Sproule BJ: Effects of alprazolam on exercise and dyspnea in patients with chronic obstructive pulmonary disease. *Chest* 90:832-836, 1986
7. Allard P, Lamontagne C, Bernard P, et al: How effective are supplementary doses of opioids for dyspnea in terminally ill cancer patients? A randomized continuous sequential clinical trial. *J Pain Symptom Manage* 17:256-265, 1999

DOI: 10.1200/JCO.2008.18.3566