

## UNIVERSAL PRIMERS AMPLIFY A 23S rDNA PLASTID MARKER IN EUKARYOTIC ALGAE AND CYANOBACTERIA<sup>1</sup>

Alison R. Sherwood<sup>2</sup>

Department of Botany, 3190 Maile Way, University of Hawai'i, Honolulu, HI 96822, USA

and

Gernot G. Presting

Department of Molecular Biosciences and Bioengineering, University of Hawai'i, Honolulu, HI 96822, USA

The challenge in the development of universal algal primers lies in the genetic diversity contained within the vast array of evolutionary lineages present in this informally named group of organisms. A comparative genomics approach was used previously to identify conserved primers flanking a region of the plastid genome. Our present research illustrates the feasibility of amplifying and sequencing this marker across multiple algal lineages. We present a preliminary framework of 107 novel sequences of this region from 62 red algae, 19 green algae, 14 brown algae, 8 cyanobacteria, 2 diatoms, 1 xanthophyte, and 1 euglenoid, and illustrate levels of divergence of the marker for well-represented groups in a neighbor-joining analysis. **This ~410 nt region distinguishes most species included in the analysis.** The remarkable universality of these primers suggests **potential for their use in assays of environmental samples in which they could be used to simultaneously detect a number of different algal lineages.**

**Key index words:** biodiversity; cyanobacteria; environmental sampling; plastid 23S rRNA; sequence diversity; universal primer

**Abbreviations:** COI, cytochrome oxidase I; cox, cytochrome oxidase; K2P, Kimura-2-parameter; nt, nucleotide; *rbcL*, large subunit of RUBISCO

To date, a number of commonly employed sets of primers have been introduced in the phycozoology literature for PCR-based analyses of algal samples. However, most have limited utility beyond the algal lineage for which they were designed, and their use has been directed primarily at specific phylogenetic or phylogeographic inquiry rather than the general assessment of large numbers of individuals from different algal lineages. For example, Saunders and

Kraft (1994) and Harper and Saunders (2001) have published several sets of primers that are widely employed in phylogenetic analyses of the nuclear ribosomal SSU and LSU genes, respectively. Zuccarello et al. (1999) introduced a set of degenerate primers flanking the mitochondrial *cox 2-3* spacer region that have been widely used in phylogeographic studies of red algae (Provan et al. 2005), and Saunders (2005) recently published a pair of mitochondrial COI primers.

A number of more broadly amplifying “universal primers” have also appeared in the literature. A set of 16S rRNA cyanobacterial primers have been employed for phylogenetic analyses and species identification for almost a decade (Nübel et al. 1997) and have recently been investigated in greater detail for diversity analyses of microbial communities (Boutte et al. 2006). The 18S rRNA gene of green algae has been frequently analyzed through amplification and sequencing using universal eukaryotic primers from Hamby et al. (1988). In 2004, Provan et al. released a suite of universal primers for plastid DNA in either green algae (~5 kb of *rps11-rpl2* gene cluster) or red algae (13–17 kb of *rps10-dnaK* ribosomal protein gene cluster). However, they approached the design of their primers with a priori separation of the two lineages, stating that “it is unlikely that truly universal primers could be developed that would amplify in both Rhodophyta and Chlorophyta” (Provan et al. 2004, p. 47). It has been demonstrated that this is not the case: Stiller and McClanahan (2005) recently reported a primer pair that amplifies both cyanobacterial and plastid 16S rDNA, to the exclusion of bacteria.

Comparative genomics provides a powerful approach for the determination of conserved regions across available genomes, and such comparisons are becoming increasingly valuable now that genomic data are becoming more representative of known lineage diversity. Universal eukaryotic plastid primers were recently designed following analysis of 37 plastid genomes from a variety of algal lineages and land plants (Presting 2006). These primers

<sup>1</sup>Received 9 October 2006. Accepted 18 January 2007.

<sup>2</sup>Author for correspondence: e-mail asherwoo@hawaii.edu.




Fig. 1. Neighbor-joining distance tree based on the plastid 23S Domain V plastid rDNA sequence data obtained with a single universal primer pair. The data include 107 algal representatives from the red, green, and brown algae, as well as the diatoms, euglenoids, xanthophytes, and cyanobacteria. GenBank accession numbers and species authorities are provided in Supplementary Table S1.

flank Domain V of the 23S plastid rRNA gene (Harris et al. 1994). Here we demonstrate the ease of using a single pair of universal primers for amplification and sequencing of this plastid marker for multiple eukaryotic algal and cyanobacterial groups, present a preliminary DNA sequence framework based on these data, and illustrate the taxonomic level at which it distinguishes taxa for at least some algal lineages.

The primer pair p23SrV\_f1 (5' GGA CAG AAA GAC CCT ATG AA 3') and p23SrV\_r1 (5' TCA GCC TGT TAT CCC TAG AG 3') was designed from these conserved flanking sequences. These exact priming sequences are present only in cyanobacteria and plastids (Presting 2006). A phylogenetically diverse range of samples was obtained for the red (62), green (19), and brown algal lineages (14) and the cyanobacteria (8), with smaller representation from other groups (2 diatoms, 1 euglenoid, and 1 xanthophyte; sample collection information can be obtained from the authors). A culture of the coral symbiont dinoflagellate, *Symbiodinium* sp., was also included in the study, but it failed to amplify and sequence cleanly—a result that is not surprising given the unique 23S rDNA plastid sequence and structure known for members of this genus (Santos et al. 2002).

The DNA was extracted using the Qiagen DNeasy Plant Mini Kit (Qiagen, Valencia, CA, USA) or a phenol–chloroform protocol (Saunders 1993). The primer pair was used to generate a ~410 nt amplicon from the plastid genome for the 107 accessions. The PCR amplification was performed in an Eppendorf Mastercycler ep gradient S thermalcycler (Eppendorf, Hamburg, Germany) using the following cycling conditions: initial denaturation at 94°C for 2 min, followed by 35 cycles of 94°C for 20 s, 55°C for 30 s, and 72°C for 30 s, with a final extension of 72°C for 10 min. Initial gradient PCR reactions indicated strong amplification with annealing temperatures between 45°C and 62°C for the templates tested (one green and one red alga). The PCR reactions totaled 50 µL in volume and consisted of 5.0 µL of 10× PCR buffer (Promega), 4.0 µL of 2.0 mM MgCl<sub>2</sub> (Promega), 3.0 µL of 1.0% BSA solution, 2.0 µL of each primer (0.4 mM), 2.0 µL (20 mM) of each dNTP, 0.2 µL of Promega Taq, 26.0 µL of nanopure water, and 1.0 µL of total genomic DNA. The PCR products were purified (using the Qiagen PCR Purification kit) and sequenced on an ABI 3730XL capillary (Applied Biosystems, Foster City, CA, USA) or ABI 377XL DNA sequencer. Chromatograms were examined, and consensus sequences assembled from forward and reverse reads using ContigExpress of the Vector NTI Suite (InforMax, Frederick, MD, USA) or Sequencher (GeneCodes Corp., Ann Arbor, MI, USA). Sequences were aligned using the Windows version of Clustal X (Thompson et al. 1997). The software package MEGA2 (Kumar et al. 2001) was used to calculate

a neighbor-joining distance tree based on K2P comparisons. Sequences have been submitted to GenBank and can be retrieved with accession numbers EF426566–EF426672. The final consensus sequences contained very few insertions and deletions and varied in length from 407 to 414 nt, including primer sequences. Alignment was greatly simplified by the low number of indels, and a final alignment of 377 nt was constructed from the 107 sequences (excluding primers).

The breadth of cyanobacterial and eukaryotic algal diversity analyzed for this study is presented in a neighbor-joining tree of sequences (Fig. 1). With the exception of the single culture of *Symbiodinium* sp., the universal primer pair was successfully used for amplification and sequencing of red, green, and brown algae; diatoms; euglenoids; xanthophytes; and cyanobacteria. With a few exceptions, sequences of closely related species are distinct.

The remarkable universality of the plastid 23S rRNA primers has both advantages and disadvantages. Not surprisingly, contaminant algal sequences are occasionally recovered with the primer pair, and this issue becomes more serious when multispecies collections are used (e.g., turf algal communities) rather than clean macroalgal thalli or cultured isolates. This same universality, however, may be exploited in the development of protocols for quantitatively assaying environmental samples.

Numerous applications exist for algal universal primers, and more are certain to be developed as molecular assessment of algal diversity plays an increasing role in control and detection of invasive species, species identifications for environmental monitoring purposes, testing of commercial algal products, and characterization of herbarium collections. The results presented here demonstrate the amplification in, and provide sequences for, representatives from six eukaryotic algal lineages as well as the cyanobacteria and establish a preliminary framework for the development of such protocols using the 23S rDNA plastid marker.

This work was supported by the National Science Foundation (DEB 0542608 to ARS and GGP) and the University of Hawai'i. Most collections of the Nemaliales were collected and identified by Dr. Isabella Abbott (University of Hawai'i) and/or Dr. John Huisman (Murdoch University), and *Prasiola/Rosenvingiella* collections were isolated and identified by Dr. Fabio Rindi (University of Alabama). We also thank Jennifer Anwar, Connie Brown, Kerry Chesser, Kristopher Coontz, Nick Kalodimos, Daniel O'Doherty, Sarah Perry, Matthew Ross, Thomas Sauvage, Heather Spalding, Cheryl Squair, Roy Tsuda, and Morgan Vis for collecting or providing samples and Tara Chandrasekharan for laboratory assistance.

Boutte, C., Grubisic, S., Balthasart, P. & Wilmotte, A. 2006. Testing of primers for the study of cyanobacterial molecular diversity by DGGE. *J. Microbiol. Methods* 65:542–50.

Hamby, R. K., Sims, L. E., Issle, L. E. & Zimmer, E. A. 1988. Direct RNA sequencing: optimization of extraction and sequencing techniques for work with higher plants. *Plant Mol. Biol. Rep.* 6:179–97.

- Harper, J. T. & Saunders, G. W. 2001. The application of sequences of the ribosomal cistron to the systematics and classification of the florideophyte red algae (Florideophyceae, Rhodophyta). *Cah. Biol. Mar.* 42:25–38.
- Harris, E. H., Boynton, J. E. & Gillham, N. W. 1994. Chloroplast ribosomes and protein synthesis. *Microbiol. Rev.* 58:700–54.
- Kumar, S., Koichiro, T., Jakobsen, I. B. & Nei, M. 2001. *MEGA2: Molecular Evolutionary Genetics Analysis Software*. Arizona State University, Tempe, Arizona, USA.
- Nübel, U., Garcia-Pichel, F. & Muyzer, G. 1997. PCR primers to amplify 16S rRNA genes from cyanobacteria. *Appl. Environ. Microbiol.* 63:3327–32.
- Presting, G. G. 2006. Identification of conserved regions in the plastid genome—implications for DNA barcoding and biological function. *Can. J. Bot.* 84:1434–43.
- Provan, J., Murphy, S. & Maggs, C. A. 2004. Universal plastid primers for Chlorophyta and Rhodophyta. *Eur. J. Phycol.* 39:43–50.
- Provan, J., Wattier, R. A. & Maggs, C. A. 2005. Phylogeographic analysis of the red seaweed *Palmaria palmata* reveals a Pleistocene marine glacial refugium in the English Channel. *Mol. Ecol.* 14:793–803.
- Santos, S. R., Taylor, D. J., Kinzie, R. A. III, Sakai, K. & Coffroth, M. A. 2002. Evolution of length variation and heteroplasmy in the chloroplast rDNA of symbiotic dinoflagellates (Symbiodinium, Dinophyta) and a novel insertion in the universal core region of the large subunit rDNA. *Phycologia* 41:311–8.
- Saunders, G. W. 1993. Gel purification of red algal genomic DNA: an inexpensive and rapid method for the isolation of polymerase chain reaction-friendly DNA. *J. Phycol.* 29:251–4.
- Saunders, G. W. 2005. Applying DNA barcoding to red macroalgae: a preliminary appraisal holds promise for future application. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 360:1879–88.
- Saunders, G. W. & Kraft, G. T. 1994. Small-subunit rRNA gene sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta). 1. Evidence for the Plocamiales ord. nov. *Can. J. Bot.* 72:1250–63.
- Stiller, J. W. & McClanahan, A. 2005. Phyto-specific 16S rDNA PCR primers for recovering algal and plant sequences from mixed samples. *Mol. Ecol. Notes* 5:1–3.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 25:4876–82.
- Zuccarello, G. C., Burger, G., West, J. A. & King, R. J. 1999. A mitochondrial marker for red algal intraspecific relationships. *Mol. Ecol.* 8:1443–7.

### Supplementary Material

The following supplemental material is available for this article:

**Table S1.** List of taxa, collection information, and associated voucher information for specimens included in analysis of the 23S rDNA plastid marker.

This material is available as part of the online article from: <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1529-8817.2007.00341.x> (This link will take you to the article abstract).

Please note: Blackwell Publishing is not responsible for the content or functionality of any supplementary materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.