

# Exercises on Non Negative Matrix Factorization

M-L. Cauwet

December 2, 2018

The following exercises are inspired from [1, 2, 3].

## Rank(s).

Consider  $X$  a  $n \times p$  non-negative matrix (i.e., every coefficient of  $X$  is non-negative) with at least one positive coefficient. The **non-negative rank** of a  $n \times p$  non-negative matrix  $X$  is the smallest integer  $k$  such that  $X = AB$  where  $A$  is a  $n \times k$  non-negative matrix and  $B$  is a  $k \times p$  non-negative matrix. The non-negative rank will be denoted by  $\text{rank}_+(X)$ . In the following,  $X_{i,j}$  denotes the coefficient on the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column of  $X$ . We denote by  $\text{rank}(X)$  the rank of  $X$ .

1. Proof that  $\text{rank}(X) \leq \text{rank}_+(X) \leq \min(n, p)$ .

2. We say that  $X_{i,j}$  and  $X_{k,\ell}$  are *independent* if

$$X_{i,j}X_{k,\ell} > 0 \text{ and } X_{i,\ell}X_{k,j} = 0.$$

Show that if  $X_{i,j}$  and  $X_{k,\ell}$  are independent, then  $\text{rank}_+(X) \geq 2$ . *Hint: write the system satisfied by the coefficients of  $A$  and  $B$  if  $\text{rank}_+(X) = 1$ .*

More generally, we could show that, if  $X$  contains a set of  $q$  pairwise independent coefficients, then  $\text{rank}_+(X) \geq q$ . (Result admitted).

3. Consider:

$$X = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

- (a) Give the rank of  $X$ .
- (b) We can show that  $X_{1,1}$ ,  $X_{2,3}$ ,  $X_{3,2}$  and  $X_{4,4}$  are pairwise independent. Deduce the non-negative rank of  $X$ .

## Exact factorization: particular case of the symmetric factorization

Assume that  $X$  is a  $n \times n$  matrix which is:

- non-negative;
- symmetric (i.e.  $X = X^T$ , where  $X^T$  is the transpose of  $X$ );
- positive semidefinite (i.e.  $v^T X v \geq 0$  for all vector  $v$ );
- $\text{rank}(X) = 2$ .

In this particular case, there exists a  $n \times 2$  non-negative matrix  $W$  such that  $X = WW^T$ . The purpose of this exercise is to build such  $W$ .

1. Show that  $X$  has exactly 2 non-zero (and positive) eigenvalues that we will denote  $\lambda_1$  and  $\lambda_2$  with  $\lambda_1 \geq \lambda_2$ . We denote  $\mathbf{v}_1$  and  $\mathbf{v}_2$  their corresponding orthonormal eigenvectors (column vectors).
2. Let  $D$  be the diagonal matrix  $D = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$  and  $P$  the  $n \times 2$  matrix  $P = (\mathbf{v}_1 \quad \mathbf{v}_2)$ . Show that  $X = PDP^T$ .
3. We define  $V$  by  $V = P\sqrt{D}$ , where  $\sqrt{D} = \begin{pmatrix} \sqrt{\lambda_1} & 0 \\ 0 & \sqrt{\lambda_2} \end{pmatrix}$ . Show that for any rotation matrix  $R$ ,  $X = WW^T$ , with  $W = VR$ . We recall that a rotation matrix  $R$  is defined by  $RR^T = R^T R = I$  and  $\det(R) = 1$ .
4.  $R = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}$ . Graphically, how to choose  $\theta$  in order to have  $W$  non-negative?

# On the equivalence of Non-Negative Matrix Factorization and K-means

We consider  $n$  points  $\mathbf{x}_1, \dots, \mathbf{x}_n$  (columns vectors). In the following,  $\|\cdot\|$  stands for the Euclidean norm. A K-means clustering over these  $n$  points consists into finding  $k$  clusters  $\mathcal{C}_1, \dots, \mathcal{C}_k$  that minimize the inertia  $J_K$ , defined by:

$$J_K = \sum_{k=1}^K \sum_{i \in \mathcal{C}_k} \|\mathbf{x}_i - \mathbf{m}_k\|^2, \quad (1)$$

where  $\mathbf{m}_k$  is the centroid of  $\mathcal{C}_k$ :  $\mathbf{m}_k = \frac{1}{n_k} \sum_{i \in \mathcal{C}_k} \mathbf{x}_i$  and  $n_k = \text{Card}(\mathcal{C}_k)$ .

1. Show that Eq. 1 can be rewritten:

$$J_K = \sum_i \|\mathbf{x}_i\|^2 - \sum_{k=1}^K \frac{1}{n_k} \sum_{i,j \in \mathcal{C}_k} \mathbf{x}_i^T \mathbf{x}_j. \quad (2)$$

2. The rescaled indicator vectors  $\delta_k$  represent a given clustering as follows:

$$\begin{cases} \delta_k^i = \frac{1}{\sqrt{n_k}} & \text{if } \mathbf{x}_i \in \mathcal{C}_k \\ 0 & \text{otherwise} \end{cases}$$

where  $\delta_k^i$  is the  $i^{\text{th}}$  component of column vector  $\delta_k$ .

We denote  $X = (\mathbf{x}_1 | \dots | \mathbf{x}_n)$  the  $p \times n$  matrix containing the data and  $\Delta = (\delta_1 | \dots | \delta_K)$  the  $n \times K$  matrix containing the indicator vectors.

Show that Eq. 2 can be rewritten:

$$J_K = \text{Tr}(X^T X) - \text{Tr}(\Delta^T X^T X \Delta), \quad (3)$$

where  $\text{Tr}(A)$  is the trace of matrix  $A$ , i.e. the sum of the elements of its diagonal.

3. Deduce from question 2 that minimizing  $J_K$  boils down to solve:

$$\max_{\Delta^T \Delta = I_K, \Delta \geq 0} \text{Tr}(\Delta^T W \Delta), \text{ where } W = X^T X. \quad (4)$$

4. Prove that Eq. 4 can be rewritten as:

$$\min_{\Delta^T \Delta = I, \Delta \geq 0} \|W - \Delta \Delta^T\|_F^2 \quad (5)$$

where  $\|\cdot\|_F^2$  is the Frobenius norm, i.e. the sum of the squared coefficient of the matrix. In particular,  $\|A\|_F^2 = \text{Tr}(AA^T)$ .

5. Show that  $\Delta \Delta^T$  is a diagonal bloc matrix of the form:

$$\Delta \Delta^T = \begin{pmatrix} \boxed{\mathbf{1}_{n_1}} & 0 & \dots & 0 \\ 0 & \boxed{\mathbf{1}_{n_2}} & & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \dots & 0 & \boxed{\mathbf{1}_{n_K}} \end{pmatrix}$$

where  $\mathbf{1}_{n_k}$  is a  $k \times k$  matrix with every coefficients equal to 1.

6. In question 4, we proved that K-means clustering is a non-negative matrix factorization with the additional constraints of:

- symmetry (i.e.  $W = AB$  with  $B = A^T$ );
- orthogonality (i.e.  $\Delta^T \Delta = I$ ).

What is this impact on the clusters if the orthogonality constraint is relaxed?

## References

- [1] Joel Cohen and Uriel G. Rothblum. Nonnegative ranks, decompositions, and factorizations of nonnegative matrices. 190:149–168, 09 1993.
- [2] Vassilis Kalofolias and Efstratios Gallopoulos. Computing symmetric nonnegative rank factorizations. *Linear Algebra and its Applications*, 436(EPFL-ARTICLE-198764):421–435, 2012.

- [3] Chris Ding, Xiaofeng He, and Horst D Simon. On the equivalence of nonnegative matrix factorization and spectral clustering. In *Proceedings of the 2005 SIAM International Conference on Data Mining*, pages 606–610. SIAM, 2005.