

# COMPUTING CUP-PRODUCTS IN INTEGER COHOMOLOGY OF HILBERT SCHEMES OF POINTS ON K3 SURFACES

SIMON KAPFER

ABSTRACT. We study the images of cup products in integer cohomology of the Hilbert scheme of  $n$  points on a K3 surface.

## 1. PRELIMINARIES

**Definition 1.1.** Let  $n$  be a natural number. A partition of  $n$  is a sequence  $\lambda = (\lambda_1 \geq \dots \geq \lambda_k > 0)$  of natural numbers such that  $\sum_i \lambda_i = n$ . It is convenient to write  $\lambda = (1^{m_1}, 2^{m_2}, \dots)$  as a sequence of multiplicities. We define the weight  $\|\lambda\| := \sum m_i i = n$  and the length  $|\lambda| := \sum_i m_i = k$ . We also define  $z_\lambda := \prod_i i^{m_i} m_i!$ .

**Definition 1.2.** Let  $\Lambda$  be the ring of symmetric functions. Let  $m_\lambda$  and  $p_\lambda$  denote the monomial and the power sum symmetric functions. They are both indexed by partitions and form a basis of  $\Lambda$ . They are linearly related by  $p_\lambda = \sum_\mu \psi_{\lambda\mu} m_\mu$ , the sum being over partitions with the same weight as  $\lambda$ , hence finite. The base change matrix  $(\psi_{\lambda\mu})$  has integral entries, but its inverse  $(\psi_{\lambda\mu}^{-1})$  has not. For example,  $p_{(2,1,1)} = 2m_{(2,1,1)} + 2m_{(2,2)} + 2m_{(3,1)} + m_{(4)}$  but  $m_{(2,1,1)} = \frac{1}{2}p_{(2,1,1)} - \frac{1}{2}m_{(2,2)} - p_{(3,1)} + p_{(4)}$ . A method to determine the coefficients  $(\psi_{\lambda\mu})$  is given in [2, Sect. 3.7].

**Definition 1.3.** Let  $S$  be a projective K3 surface. We fix integral bases  $1$  of  $H^0(S, \mathbb{Z})$ ,  $x$  of  $H^4(S, \mathbb{Z})$  and  $\alpha_1, \dots, \alpha_{22}$  of  $H^2(S, \mathbb{Z})$ . The cup product induces a symmetric bilinear form  $B_{H^2}$  on  $H^2(S, \mathbb{Z})$  and thus the structure of a lattice. It turns out, that  $H^2(S, \mathbb{Z})$  is isomorphic to  $U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$ , *i.e.* three times the hyperbolic lattice and two times the negative  $E_8$  lattice. We may extend  $B_{H^2}$  to a symmetric non-degenerate bilinear form  $B$  on  $H^*(S, \mathbb{Z})$  by setting  $B(1, 1) = 0$ ,  $B(1, \alpha_i) = 0$ ,  $B(1, x) = 1$ ,  $B(x, x) = 0$ .

**Definition 1.4.**  $B$  induces a form  $B \otimes B$  on  $\text{Sym}^2 H^*(S, \mathbb{Z})$ . So the cup-product

$$\mu : \text{Sym}^2 H^*(S, \mathbb{Z}) \longrightarrow H^*(S, \mathbb{Z})$$

has an adjoint comultiplication  $\Delta$  that is coassociative, given by:

$$\Delta : H^*(S, \mathbb{Z}) \longrightarrow \text{Sym}^2 H^*(S, \mathbb{Z}), \quad \Delta = (B \otimes B)^{-1} \mu^T B$$

The image of  $1$  under the composite map  $\mu(\Delta(1)) = B(\Delta(1), \Delta(1)) = 24x$ , denoted by  $e$  is called the Euler Class.

We denote by  $S^{[n]}$  the Hilbert scheme of  $n$  points on  $S$ , *i.e.* the classifying space of all zero-dimensional closed subschemes of length  $n$ , which is smooth. A classical result by Nakajima gives an explicit description of  $H^*(S^{[n]}, \mathbb{Q})$  in terms of creation operators  $\mathfrak{q}_l(\beta)$ ,  $\beta \in H^*(S, \mathbb{Q})$ , acting on the direct sum  $\bigoplus_n H^*(S^{[n]}, \mathbb{Q})$ .

---

*Date:* September 29, 2014.

An integral basis for  $H^*(S^{[n]}, \mathbb{Z})$  in terms of Nakajima's operators was given by Qin–Wang:

**Theorem 1.5.** [7, Thm. 5.4.] *Let  $\mathbf{m}_{\nu, \alpha} := \sum_{\rho} \psi_{\nu\rho}^{-1} \mathbf{q}(\alpha)$ , with coefficients as in Definition 1.2. The classes*

$$\frac{1}{z_{\lambda}} \mathbf{q}_{\lambda}(1) \mathbf{q}_{\mu}(x) \mathbf{m}_{\nu^1, \alpha_1} \dots \mathbf{m}_{\nu^{22}, \alpha_{22}} |0\rangle, \quad \|\lambda\| + \|\mu\| + \sum_{i=1}^{22} \|\nu^i\| = n$$

*form an integral basis for  $H^*(S^{[n]}, \mathbb{Z})$ . Here,  $\lambda, \mu, \nu^i$  are partitions.*

**Notation 1.6.** To enumerate the basis of  $H^*(S^{[n]}, \mathbb{Z})$ , we introduce the following abbreviation:

$$1^{\lambda} \alpha_1^{\nu^1} \dots \alpha_{22}^{\nu^{22}} x^{\mu} := \frac{1}{z_{\tilde{\lambda}}} \mathbf{q}_{\tilde{\lambda}}(1) \mathbf{q}_{\mu}(x) \mathbf{m}_{\nu^1, \alpha_1} \dots \mathbf{m}_{\nu^{22}, \alpha_{22}} |0\rangle$$

where the partition  $\tilde{\lambda}$  is built from  $\lambda$  by appending sufficiently many Ones, such that  $\|\tilde{\lambda}\| + \|\mu\| + \sum \|\nu^i\| = n$ . If  $\|\lambda\| + \|\mu\| + \sum \|\nu^i\| > n$ , we put  $1^{\lambda} \alpha_1^{\nu^1} \dots \alpha_{22}^{\nu^{22}} x^{\mu} = 0$ .

The ring structure of  $H^*(S^{[n]}, \mathbb{Q})$  has been studied by Lehn and Sorger in [3], where an explicit algebraic model is constructed, which we recall briefly:

**Definition 1.7.** [3, Sect. 2] Let  $\pi$  be a permutation of  $n$  letters, written as a sum of disjoint cycles. To each cycle we may associate an element of  $A := H^*(S, \mathbb{Q})$ . This defines an element in  $A^{\otimes m}$ ,  $m$  being the number of cycles. So these mappings span a vector space over  $\mathbb{Q}$ . The space obtained by taking the direct sum over all  $\pi \in S_n$  will be denoted by  $A\{S_n\}$ .

To define a ring structure, take two permutations  $\pi, \tau$ , together with mappings. The result of the multiplication will be the permutation  $\pi\tau$ , together with a mapping of cycles. To construct the mappings to  $A$ , look first at the orbit space of the group of permutations  $\langle \pi, \tau \rangle$ , generated by  $\pi$  and  $\tau$ . For each cycle of  $\pi, \tau$  contained in one orbit  $B$  of  $\langle \pi, \tau \rangle$ , multiply with the associated element of  $A$ . Also multiply with a certain power of the Euler class  $e^g$ . Afterwards, apply the comultiplication  $\Delta$  repeatedly on the product to get a mapping from the cycles of  $\pi\tau$  contained in  $B$  to  $A$ . Here the "graph defect"  $g$  is defined as follows: Let  $u, v, w$  be the number of cycles contained in  $B$  of  $\pi, \tau, \pi\tau$ , respectively. Then  $g := \frac{1}{2}(|B| + 2 - u - v - w)$ . Now follow this procedure for each orbit  $B$ .

The symmetric group  $S_n$  acts on  $A\{S_n\}$  by conjugation. This action preserves the ring structure. Therefore the space of invariants  $A^{[n]} := (A\{S_n\})^{S_n}$  becomes a subring. The main theorem of [3] can now be stated:

**Theorem 1.8.** [3, Thm. 3.2.] *The following map is an isomorphism of rings:*

$$H^*(S^{[n]}, \mathbb{Q}) \longrightarrow A^{[n]} \\ \mathbf{q}_{n_1}(\beta_1) \dots \mathbf{q}_{n_k}(\beta_k) |0\rangle \longmapsto \sum_{\sigma \in S_n} \sigma a \sigma^{-1}$$

*with  $n_1 + \dots + n_k = n$  and  $a \in A\{S_n\}$  corresponds to an arbitrary permutation with  $k$  cycles of lengths  $n_1, \dots, n_k$  that are associated to the classes  $\beta_1, \dots, \beta_k \in H^*(S, \mathbb{Q})$ , respectively.*

Since  $H^{\text{odd}}(S^{[n]}, \mathbb{Z}) = 0$  and  $H^{\text{even}}(S^{[n]}, \mathbb{Z})$  is torsion-free by [4], we can apply these results to  $H^*(S^{[n]}, \mathbb{Z})$  to determine the multiplicative structure of cohomology with integer coefficients. It turns out, that it is somehow independent of  $n$ . More precisely, we have the following stability theorem due to Li, Qin and Wang:

**Theorem 1.9.** [7, Thm. 2.1] *Let  $Q_1, \dots, Q_s$  be products of creation operators, i.e.  $Q_i = \prod_j \mathfrak{q}_{\lambda_{i,j}}(\beta_{i,j})$  for some partitions  $\lambda_{i,j}$  and classes  $\beta_{i,j} \in H^*(S, \mathbb{Z})$ . Set  $n_i := \sum_j \|\lambda_{i,j}\|$ . Then the cup product  $\prod_{i=1}^s \left( \frac{1}{(n-n_i)!} \mathfrak{q}_{n-n_i}(1) Q_i |0\rangle \right)$  is equal to a finite linear combination of classes of the form  $\frac{1}{(n-m)!} \mathfrak{q}_{n-m}(1) \prod_j \mathfrak{q}_{\mu_j}(\gamma_j) |0\rangle$ , with  $\gamma \in H^*(S, \mathbb{Z})$ ,  $m = \sum_j \|\mu_j\|$ , whose coefficients are independent of  $n$ . We have the upper bound  $m \leq \sum_i n_i$ .*

**Corollary 1.10.** *The coefficients  $c_i$  of the cup product in  $H^*(S^{[n]}, \mathbb{Z})$*

$$1^{\lambda^0} \alpha_1^{\lambda^1} \dots \alpha_{22}^{\lambda^{22}} x^{\lambda^{23}} \smile 1^{\mu^0} \alpha_1^{\mu^1} \dots \alpha_{22}^{\mu^{22}} x^{\mu^{23}} = \sum_i c_i \cdot 1^{\nu_i^0} \alpha_1^{\nu_i^1} \dots \alpha_{22}^{\nu_i^{22}} x^{\nu_i^{23}}$$

*are polynomials in  $n$  of degree at most  $\sum_{j=0}^{23} \|\lambda^j\| + \|\mu^j\| - \|\nu_i^j\|$ .*

## 2. COMPUTATIONAL RESULTS

We now give some results in low degrees, obtained by computing multiplication matrices with respect to the integral basis and a then reducing to Smith normal form (both done by a computer).

*Remark 2.1.* Denote  $h^k(S^{[n]})$  the rank of  $H^k(S^{[n]}, \mathbb{Z})$ . We have:

- $h^2(S^{[n]}) = 23$  for  $n \geq 2$ .
- $h^4(S^{[n]}) = 276, 299, 300$  for  $n = 2, 3, \geq 4$  resp.
- $h^6(S^{[n]}) = 23, 2554, 2852, 2875, 2876$  for  $n = 2, 3, 4, 5, \geq 6$  resp.

The algebra generated by classes of degree 2 is an interesting object to study. For cohomology with complex coefficients, Verbitsky has proven in [8] that the cup product mapping from  $\text{Sym}^k H^2(S^{[n]}, \mathbb{C})$  to  $H^{2k}(S^{[n]}, \mathbb{C})$  is injective for  $k \leq n$ . One concludes that this also holds for integral coefficients.

**Proposition 2.2.** *Studying the image of  $\text{Sym}^2 H^2$  in  $H^4$ , we obtain:*

$$\frac{H^4(S^{[2]}, \mathbb{Z})}{\text{Sym}^2 H^2(S^{[2]}, \mathbb{Z})} \cong \left( \frac{\mathbb{Z}}{2\mathbb{Z}} \right)^{\oplus 23} \oplus \frac{\mathbb{Z}}{5\mathbb{Z}}$$

*This was already known to Boissière, Nieper-Wißkirchen and Sarti, [1, Prop. 3].*

$$\frac{H^4(S^{[3]}, \mathbb{Z})}{\text{Sym}^2 H^2(S^{[3]}, \mathbb{Z})} \cong \frac{\mathbb{Z}}{3\mathbb{Z}} \oplus \mathbb{Z}^{\oplus 23}$$

*The torsion part of the quotient is generated by the integral class  $1^{(3)}$ .*

$$\frac{H^4(S^{[n]}, \mathbb{Z})}{\text{Sym}^2 H^2(S^{[n]}, \mathbb{Z})} \cong \mathbb{Z}^{\oplus 24}, \quad \text{for } n \geq 4.$$

*This was already proven by Markman, [5, Thm. 1.10].*

**Proposition 2.3.** *Studying triple products of  $H^2(S^{[n]}, \mathbb{Z})$ , we get:*

$$\frac{H^6(S^{[2]}, \mathbb{Z})}{\text{Sym}^3 H^2(S^{[2]}, \mathbb{Z})} \cong \frac{\mathbb{Z}}{2\mathbb{Z}}$$

*The quotient is generated by the integral class  $1^{(2)}$ .*

$$\frac{H^6(S^{[3]}, \mathbb{Z})}{\text{Sym}^3 H^2(S^{[3]}, \mathbb{Z})} \cong \left(\frac{\mathbb{Z}}{2\mathbb{Z}}\right)^{\oplus 230} \oplus \left(\frac{\mathbb{Z}}{36\mathbb{Z}}\right)^{\oplus 22} \oplus \frac{\mathbb{Z}}{72\mathbb{Z}} \oplus \mathbb{Z}^{\oplus 507}$$

$$\frac{H^6(S^{[4]}, \mathbb{Z})}{\text{Sym}^3 H^2(S^{[4]}, \mathbb{Z})} \cong \frac{\mathbb{Z}}{2\mathbb{Z}} \oplus \mathbb{Z}^{\oplus 552}$$

*For  $n \geq 5$ , the quotient is free.*

We study now cup products between classes of degree 2 and 4. The case of  $S^{[3]}$  is of particular interest.

**Proposition 2.4.** *Comparing  $H^2(S^{[n]}, \mathbb{Z}) \smile H^4(S^{[n]}, \mathbb{Z})$  with  $H^6(S^{[n]}, \mathbb{Z})$ , we obtain:*

$$\begin{aligned} (1) \quad & \frac{H^6(S^{[2]}, \mathbb{Z})}{H^2(S^{[2]}, \mathbb{Z}) \smile H^4(S^{[2]}, \mathbb{Z})} = 0 \\ (2) \quad & \frac{H^6(S^{[3]}, \mathbb{Z})}{H^2(S^{[3]}, \mathbb{Z}) \smile H^4(S^{[3]}, \mathbb{Z})} \cong \left(\frac{\mathbb{Z}}{3\mathbb{Z}}\right)^{\oplus 22} \oplus \frac{\mathbb{Z}}{3\mathbb{Z}} \\ (3) \quad & \frac{H^6(S^{[4]}, \mathbb{Z})}{H^2(S^{[4]}, \mathbb{Z}) \smile H^4(S^{[4]}, \mathbb{Z})} \cong \left(\frac{\mathbb{Z}}{6\mathbb{Z}}\right)^{\oplus 22} \oplus \frac{\mathbb{Z}}{108\mathbb{Z}} \oplus \frac{\mathbb{Z}}{2\mathbb{Z}} \\ (4) \quad & \frac{H^6(S^{[5]}, \mathbb{Z})}{H^2(S^{[5]}, \mathbb{Z}) \smile H^4(S^{[5]}, \mathbb{Z})} \cong \mathbb{Z}^{\oplus 22} \oplus \mathbb{Z} \\ (5) \quad & \frac{H^6(S^{[n]}, \mathbb{Z})}{H^2(S^{[n]}, \mathbb{Z}) \smile H^4(S^{[n]}, \mathbb{Z})} \cong \mathbb{Z}^{\oplus 22} \oplus \mathbb{Z} \oplus \mathbb{Z}, \quad n \geq 6. \end{aligned}$$

*In each case, the first 22 parts of the quotient are generated by the integral classes*

$$\alpha_i^{(1,1,1)} - 3 \cdot \alpha_i^{(2,1)} + 3 \cdot \alpha_i^{(3)} + 3 \cdot 1^{(2)} \alpha_i^{(1,1)} - 6 \cdot 1^{(2)} \alpha_i^{(2)} + 6 \cdot 1^{(2,2)} \alpha_i^{(1)} - 3 \cdot 1^{(3)} \alpha_i^{(1)},$$

*for  $i = 1 \dots 22$ . Now define an integral class*

$$\begin{aligned} K := & \sum_{i \neq j} B(\alpha_i, \alpha_j) \left[ \alpha_i^{(1,1)} \alpha_j^{(1)} - 2 \cdot \alpha_i^{(2)} \alpha_j^{(1)} + \frac{3}{2} \cdot 1^{(2)} \alpha_i^{(1)} \alpha_j^{(1)} \right] + \\ & + \sum_i B(\alpha_i, \alpha_i) \left[ \alpha_i^{(1,1,1)} - 2 \cdot \alpha_i^{(2,1)} + \frac{3}{2} \cdot 1^{(2)} \alpha_i^{(1,1)} \right] + x^{(2)} - 1^{(2)} x^{(1)}. \end{aligned}$$

*In the case  $n = 3$ , the last part of the quotient is generated by  $K$ .*

*In the case  $n = 4$ , the class  $1^{(4)}$  generates the 2-torsion part and  $K + 38 \cdot 1^{(4)}$  generates the 108-torsion part.*

*In the case  $n = 5$ , the last part of the quotient is generated by  $K + 16 \cdot 1^{(4)} - 21 \cdot 1^{(3,2)}$ .*

*If  $n \geq 6$ , the two last parts of the quotient are generated by some multiples of  $K - \frac{4}{3}(45 - n)1^{(2,2,2)} + (48 - n)1^{(3,2)}$  and  $K - \frac{1}{2}(40 - n)1^{(2,2,2)} + \frac{1}{4}(48 - n)1^{(4)}$ .*

**Remark 2.5.** (1) The first statement of Prop. 2.4 is clear by Poincaré Duality.

(2) Observe that the generators of the quotients are independent of the choice of the base of  $H^2(S, \mathbb{Z})$ .

## REFERENCES

1. S. Boissière, M. Nieper-Wißkirchen, and A. Sarti, *Smith theory and Irreducible Holomorphic Symplectic Manifolds*, eprint arXiv:1204.4118 (2012).
2. A. Lascoux, *Symmetric functions*, Notes of the course given at Nankai University, 2001, <http://www.mat.univie.ac.at/slc/wpapers/s68vortrag/ALCoursSf2.pdf>.
3. M. Lehn and C. Sorger, *The cup product of Hilbert schemes for K3 surfaces*, Invent. Math. **152** (2003), no. 2, 305–329.
4. E. Markman, *Integral generators for the cohomology ring of moduli spaces of sheaves over Poisson surfaces*, Adv. Math. **208** (2007), no. 2, 622–646.
5. E. Markman, *Integral constraints on the monodromy group of the hyperKähler resolution of a symmetric product of a K3 surface*, Internat. J. Math. **21** (2010), no. 2, 169–223.
6. H. Nakajima, *Heisenberg algebra and Hilbert schemes of points on projective surfaces*, Ann. of Math. (2) **145** (1997), no. 2, 379–388.
7. Z. Qin and W. Wang, *Integral operators and integral cohomology classes of Hilbert schemes*, Math. Ann. **331** (2005), no. 3, 669–692.
8. M. Verbitsky, *Cohomology of compact hyper-Kähler manifolds and its applications*, Geom. Funct. Anal. **6** (1996), no. 4, 601–611.

SIMON KAPFER, LEHRSTUHL FÜR ALGEBRA UND ZAHLENTHEORIE, UNIVERSITÄTSSTRASSE 14,  
D-86159 AUGSBURG

*E-mail address:* `simon.kapfer@math.uni-augsburg.de`