

Petersson Inner Product of Theta Series

An experimental approach

Nicolas SIMARD

McGill University

December 1st, 2017

Stark's observation

Let $K = \mathbb{Q}(\sqrt{-23})$ and let H be the HCF of K . Let

$$\psi : \text{Gal}(H/K) \rightarrow \text{GL}_1(\mathbb{C})$$

be a non-trivial one-dimensional Artin representation and let

$$\rho = \text{Ind}_K^{\mathbb{Q}} \psi : \text{Gal}(H/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{C})$$

be the induced representation.

Stark's observation (cont.)

By Deligne-Serre, one has

$$L(\rho, s) = L(\theta_\psi, s),$$

where

$$\theta_\psi(q) = \eta(q)\eta(23q) = q \prod_{n \geq 1} (1 - q^n)(1 - q^{23n}) \in M_1(\Gamma_0(23), \chi_{-23}).$$

Then Stark proves that

$$\langle \theta_\psi, \theta_\psi \rangle = 3 \log \varepsilon,$$

where ε is the real root of

$$x^3 - x - 1.$$

Structure of the talk

Introduction

Motivation

Structure of the Talk

Petersson Inner Product of Weight One Theta Series

Explicit Formulas

Generalizing Stark's Observation

Petersson Inner Product of Higher Weight Theta Series

Spaces of Theta Series

Notation

Throughout this talks, let

- K be an imaginary quadratic field of discriminant D ,
- H be the Hilbert class field of K ,
- h_K be the class number of K ,
- w_K be the number of roots of unity in K and
- ClK be the class group of K .

Weight one theta series

Let ψ be a class character of K , i.e. a homomorphism

$$\psi : \text{Cl}_K \rightarrow \mathbb{C}^\times.$$

Then

$$\theta_\psi(q) = \sum_{\mathfrak{a}} \psi(\mathfrak{a}) q^{N(\mathfrak{a})} \in M_1(\Gamma_0(|D|), \chi_D).$$

Moreover, θ_ψ is an eigenform for all Hecke operators.

If $\psi^2 = 1$, θ_ψ is an Eisenstein series.

If $\psi^2 \neq 1$, θ_ψ is a cusp form (in fact, a newform).

Stark's example

Let

$$K = \mathbb{Q}(\sqrt{-23})$$

and let ψ be a non-trivial class character as above. Then

$$\text{Stark's } \theta_\psi = \text{our } \theta_\psi \in M_1(\Gamma_0(23), \chi_{-23}).$$

Note that if ψ' is the other non-trivial class character, then

$$\theta_\psi = \theta_{\psi'}.$$

Petersson inner product of weight one theta series

The Petersson inner product of any two $f, g \in S_k(\Gamma_0(N), \chi)$ is defined as

$$\langle f, g \rangle = \iint_{\Gamma_0(N) \backslash \mathcal{H}} f(x + iy) \overline{g(x + iy)} y^k d\mu.$$

Then

Proposition (S.)

Let ψ be a class character which is not a genus character. Then

$$\langle \theta_\psi, \theta_\psi \rangle = \frac{-h_K}{3w_K^2} \sum_{\mathcal{A} \in Cl_K} \psi^2(\mathcal{A}) \log N(\mathcal{A})^6 |\Delta(\mathcal{A})|.$$

Siegel units

Let \mathfrak{a} be a fractional ideal of K and define

$$|\delta_{\mathcal{A}}| = (N(\mathfrak{a})^6 |\Delta(\mathcal{O}_K)/\Delta(\mathfrak{a}^{-1})|)^{h_K},$$

where \mathfrak{a} is any ideal in the class \mathcal{A} . Then $|\delta_{\mathcal{A}}|$ is a unit in H . Since ψ^2 is not trivial, one sees that

$$\langle \theta_{\psi}, \theta_{\psi} \rangle = \frac{1}{3w_K^2} \sum_{\mathcal{A} \in \text{Cl}_K} \psi^2(\mathcal{A}) \log |\delta_{\mathcal{A}}|,$$

where $\{\mathfrak{a}_1, \dots, \mathfrak{a}_{h_K}\}$ is a set of class representatives for Cl_K .

What about Stark's observation?

One can write

$$\langle \theta_\psi, \theta_\psi \rangle = h_K \log \kappa_\psi,$$

where

$$\kappa_\psi = \prod_{j=1}^{h_K} \Phi(\mathfrak{a}_j)^{-\psi^2(\mathfrak{a}_j)}$$

with

$$\Phi(\mathfrak{a}) = \sqrt{N(\mathfrak{a})} |\Delta(\mathfrak{a})|^{1/12}.$$

Question

Is κ_ψ a unit in H ?

Calcs in class nbr 3, 4, 5, 6.

Generalizing Stark's Observation

Proposition (S.)

Let ψ be a class character such that ψ^2 is a non-trivial character with rational real part. Then κ_ψ is an algebraic integer which is a unit. Moreover, if ψ^2 is a non-trivial genus character corresponding to the factorisation $D = D_1 D_2$, with $D_1 > 0$ say, then

$$\kappa_\psi = \epsilon_{D_1}^{\frac{4h_{D_1} h_{D_2}}{w_K w_{D_2}}},$$

where ϵ_{D_1} is the fundamental unit of $\mathbb{Q}(\sqrt{D_1})$, h_{D_j} is the class number of $\mathbb{Q}(\sqrt{D_j})$ and w_{D_2} is the number of roots of unity in $\mathbb{Q}(\sqrt{D_2})$.

Examples

If $K = \mathbb{Q}(\sqrt{-23})$, the Proposition implies that κ_ψ is a unit. But is it in the Hilbert class field?

Examples

If $K = \mathbb{Q}(\sqrt{-23})$, the Proposition implies that κ_{ψ} is a unit. But is it in the Hilbert class field?

If $K = \mathbb{Q}(\sqrt{-39})$, the Proposition implies

$$\kappa_{\psi} = \epsilon_{13}^{\frac{1}{3}},$$

which is *not* in the Hilbert class field.

Stark's observation: the final word?

Note that ψ^2 has rational real part if and only if its order divides 4 or 3.

Corollary

Suppose that K has class number divisible by 2 or 3. Then there exists a class character ψ such that

$$\kappa_\psi$$

is a unit.

Question

Is the converse true?

Thank you!

Presentation and notes available at:

<https://github.com/NicolasSimard/Notes>

Code available at : <https://github.com/NicolasSimard/ENT>

Or from my webpage: <http://www.math.mcgill.ca/nsimard/>