

# Constructing the p-adic zeta function via cyclotomic units

Nicolas Simard

October 29, 2016

## Contents

<b>1</b>	<b>p-adic measures</b>	<b>1</b>
1.1	p-adic measures, distributions and Iwasawa algebras . . . . .	1
1.1.1	Distributions and Iwasawa algebras . . . . .	2
1.1.2	Distributions and step functions . . . . .	2
1.1.3	p-adic measures and continuous functions . . . . .	3
1.1.4	The Iwasawa algebra $\Lambda(\mathbb{Z}_p)$ and Mahler's transform . . . . .	5
1.1.5	The Iwasawa algebra $\Lambda(\mathbb{Z}_p^\times)$ . . . . .	6
1.1.6	The Iwasawa algebras $\Lambda(\mathcal{G})$ and $\Lambda(G)$ . . . . .	6
1.2	Operators on p-adic measures . . . . .	7
1.2.1	Operators on $\mathbb{R}$ . . . . .	7
1.2.2	Operators on $\Lambda(\mathbb{Z}_p)$ . . . . .	7
1.2.3	Restriction of measures from $\mathbb{Z}_p$ to $\mathbb{Z}_p^\times$ . . . . .	7
1.3	Moments of p-adic measures . . . . .	7
1.4	Summary . . . . .	7

## Introduction

### 1 p-adic measures

In this section, we first define p-adic measures and see how they are related to Iwasawa Algebras and power series rings. We then introduce operators on them and conclude with of few results on moments of measures.

#### 1.1 p-adic measures, distributions and Iwasawa algebras

Let  $\mathcal{G}$  be an abelian profinite group, let  $\mathfrak{B}_{\mathcal{G}}$  be the boolean algebra of open subsets of  $\mathcal{G}$ , let  $\mathfrak{T}_{\mathcal{G}} \subseteq \mathfrak{B}_{\mathcal{G}}$  be the set of open subgroups of  $\mathcal{G}$  and let  $A$  be any abelian group.

**Definition 1.** An  $A$ -valued distribution  $\lambda$  on  $\mathcal{G}$  is a finitely additive function

$$\lambda : \mathfrak{T}_{\mathcal{G}} \rightarrow A.$$

The set of distributions is denoted  $\mathcal{D}(\mathcal{G}, A)$ . If  $A \subseteq \mathbb{C}_p$ , the elements of  $\mathcal{D}(\mathcal{G}, A)$  are called p-adic distributions.

The set  $\mathcal{D}(\mathcal{G}, A)$  is naturally an abelian group. If  $A$  is a  $B$ -algebra for some ring  $B$ , the set  $\mathcal{D}(\mathcal{G}, A)$  is a  $B$ -algebra under convolution product, which we won't bother to define here!

### 1.1.1 Distributions and Iwasawa algebras

If  $\mathfrak{G}$  is finite,  $\mathfrak{B}_{\mathfrak{G}} = \{\{g\} | g \in \mathfrak{G}\}$  and we have an isomorphism of abelian groups

$$\lambda \mapsto \sum_{g \in \mathfrak{G}} \lambda(\{g\})g : \mathfrak{D}(\mathfrak{G}, A) \rightarrow A[\mathfrak{G}].$$

If  $A$  is a  $B$ -algebra for some ring  $B$ , so is  $A[\mathfrak{G}]$  and the isomorphism is an isomorphism of  $B$ -algebras. For  $\mathfrak{G}$  finite, we define

$$\Lambda(\mathfrak{G}, A) := A[\mathfrak{G}]$$

and call it an Iwasawa algebra.

For  $\mathfrak{G}$  not necessarily finite, we define

$$\Lambda(\mathfrak{G}, A) = \varprojlim \Lambda(\mathfrak{G}/\mathfrak{H}, A) = \varprojlim A[\mathfrak{G}/\mathfrak{H}],$$

where the limit is taken over all elements of  $\mathfrak{T}_{\mathfrak{G}}$ . Given  $\mathfrak{H}$  in  $\mathfrak{T}_{\mathfrak{G}}$ , we have a natural map

$$\lambda \mapsto \sum_{g \bmod \mathfrak{H}} \lambda(g)(g + \mathfrak{H}) : \mathfrak{D}(\mathfrak{G}, A) \rightarrow \Lambda(\mathfrak{G}/\mathfrak{H}, A).$$

Since distributions are finitely additive, we have then a natural map

$$\mathfrak{D}(\mathfrak{G}, A) \rightarrow \Lambda(\mathfrak{G}, A),$$

which is in fact an isomorphism. In a certain sense, the elements of the Iwasawa algebra  $\Lambda(\mathfrak{G}, A)$  are like the generating series of distributions.

**Example:** If  $A = \mathbb{Z}_p$ , one obtains the usual Iwasawa algebra

$$\Lambda(\mathfrak{G}) := \Lambda(\mathfrak{G}, \mathbb{Z}_p).$$

### 1.1.2 Distributions and step functions

From now on, suppose that  $A$  is a  $B$ -algebra for some ring  $B$ .

Recall that if  $s : \mathfrak{G} \rightarrow A$  is a locally constant function, also called a step function, there exists an open subgroup  $\mathfrak{H}$  such that  $s$  is well defined on  $\mathfrak{G}/\mathfrak{H}$ , i.e.

$$s(x) = \sum_{g \in \mathfrak{G}/\mathfrak{H}} s(g)\varepsilon_g(x),$$

where  $\varepsilon_g(x)$  is the characteristic function of the (open) coset  $g \in \mathfrak{G}/\mathfrak{H}$ . Note that such a representation of  $s$  as a finite linear combination of characteristic functions is not unique. The set of step functions from  $\mathfrak{G}$  to  $A$  is denoted

$$\text{Step}(\mathfrak{G}, A).$$

If  $A$  is a  $B$ -algebra,  $\text{Step}(\mathfrak{G}, A)$  is a  $B$ -algebra under the pointwise addition and multiplication of functions.

Let  $\lambda$  be an  $A$ -valued distribution on  $\mathfrak{G}$ , let  $s$  be a step function which is well-defined  $\bmod \mathfrak{H}$  as above and define

$$\int_{\mathfrak{G}} s d\lambda := \sum_{g \in \mathfrak{G}/\mathfrak{H}} s(g)\lambda(g).$$

This gives a well-defined  $B$ -linear map

$$\Lambda(\mathfrak{G}, A) \rightarrow \text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A).$$

For convenience, the value of an element  $\lambda \in \text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A)$  at a step function  $s(x)$  is denoted

$$\int_{\mathfrak{G}} s(x) d\lambda(x)$$

or simply

$$\int_{\mathfrak{G}} s d\lambda$$

when there is no risk of confusion. The  $B$ -module

$$\text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A)$$

can be equipped with a natural  $B$ -algebra structure via the convolution product which is defined as follows: for  $\lambda, \mu \in \text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A)$ , let  $\lambda * \mu$  be defined as

$$\int_{\mathfrak{G}} f d\lambda * \mu = \int_{\mathfrak{G}} (f(x+y) d\lambda(x)) d\mu(y).$$

The map above is then a  $B$ -algebra homomorphism. In fact, it is an isomorphism. Indeed, its inverse takes a homomorphism  $\lambda \in \text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A)$  to the distribution  $\lambda$  defined as

$$\lambda(U) = g(\varepsilon_U).$$

This sketches the proof of the following proposition.

**Proposition 1.** *There is a natural  $B$ -algebra isomorphism*

$$\Lambda(\mathfrak{G}, A) \rightarrow \text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A).$$

### 1.1.3 $p$ -adic measures and continuous functions

From now on, suppose that  $A$  is contained in  $\mathbb{C}_p$  (e.g.  $A = B = \mathbb{Z}_p$ ). Let

$$C(\mathfrak{G}, \mathbb{C}_p)$$

be the set of continuous functions from  $\mathfrak{G}$  to  $\mathbb{C}_p$ . This is a  $\mathbb{C}_p$ -Banach when equipped with the norm

$$\|f\| = \sup_{x \in \mathfrak{G}} |f(x)|_p.$$

The set  $\text{Step}(\mathfrak{G}, \mathbb{C}_p)$  is dense in  $C(\mathfrak{G}, \mathbb{C}_p)$ .

**Definition 2.** A  $p$ -adic distribution  $\lambda \in \mathfrak{D}(\mathfrak{G}, A)$  is called a  $p$ -adic measure if it is bounded (as a function from  $\mathfrak{T}_{\mathfrak{G}}$  to  $A \subseteq \mathbb{C}_p$ ). The set of  $p$ -adic measures is denoted  $\mathfrak{M}(\mathfrak{G}, A)$ .

Note that if  $A$  is bounded, which is the case if  $A = \mathbb{Z}_p$  for example, then  $\mathfrak{M}(\mathfrak{G}, A) = \mathfrak{D}(\mathfrak{G}, A)$ . The importance of introducing measures is the following proposition.

**Proposition 2.** *Let  $\lambda \in \mathfrak{M}(\mathfrak{G}, \mathbb{C}_p)$  be viewed as an element of*

$$\text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p)$$

*via the above isomorphism. Then  $\lambda$  extends uniquely to a continuous map*

$$\lambda : C(\mathfrak{G}, \mathbb{C}_p) \rightarrow \mathbb{C}_p.$$

*Proof.* Let  $\lambda$  be a  $p$ -adic measure and suppose that

$$|\lambda(U)|_p \leq M$$

for all  $U \in \mathfrak{B}_{\mathfrak{G}}$  and some  $M \in \mathbb{R}$ . By the density of  $\text{Step}(\mathfrak{G}, \mathbb{C}_p)$  in  $C(\mathfrak{G}, \mathbb{C}_p)$ , for any  $f \in C(\mathfrak{G}, \mathbb{C}_p)$  one can find a sequence of step functions  $s_n \in \text{Step}(\mathfrak{G}, \mathbb{C}_p)$  such that

$$f(x) = \lim_{n \rightarrow \infty} s_n(x).$$

Then it is easy to see that for any integers  $m$  and  $n$ ,

$$\lambda(s_n - s_m) \leq M \|s_n - s_m\|.$$

Since the sequence  $\{s_n\}$  is Cauchy, so is the sequence  $\{\lambda(s_n)\}$  and it makes sense to define

$$\lambda(f) = \lim_{n \rightarrow \infty} \lambda(s_n).$$

The uniqueness is clear. □

For  $\lambda \in \text{Hom}_{\text{cont}}(C(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p)$ , define

$$\|\lambda\| = \sup_{f \in C(\mathfrak{G}, \mathbb{C}_p)} \frac{|\lambda(f)|}{\|f\|},$$

which is a finite real number by the continuity of  $\lambda$ . Equipped with the convolution product, this set becomes a  $\mathbb{C}_p$ -Banach algebra.

In the case where  $A = \mathbb{Z}_p$ , we have  $\mathfrak{M}(\mathfrak{G}, \mathbb{Z}_p) = \mathfrak{D}(\mathfrak{G}, \mathbb{Z}_p)$  and we have the following proposition.

**Proposition 3.** *The image of  $\mathfrak{M}(\mathfrak{G}, \mathbb{Z}_p)$  under the injection of the previous proposition is the set of*

$$\lambda \in \text{Hom}_{\text{cont}}(C(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p)$$

such that

$$\|\lambda\| \leq 1 \quad \text{and} \quad \lambda(C(\mathfrak{G}, \mathbb{Q}_p)) \subseteq \mathbb{Q}_p.$$

*Proof.* Let  $\lambda \in \mathfrak{M}(\mathfrak{G}, \mathbb{Z}_p)$  and take

$$s \in \text{Step}(\mathfrak{G}, \mathbb{Q}_p).$$

Writing

$$s = \sum_{g \in \mathfrak{G}/\text{Hfrak}} s(g) \varepsilon_g,$$

we see that

$$\int_{\mathfrak{G}} s d\lambda = \sum_{g \in \mathfrak{G}/\text{Hfrak}} s(g) \lambda(g) \in \mathbb{Q}_p$$

and so

$$\left\| \int_{\mathfrak{G}} s d\lambda \right\|_p \leq \sup_{g \in \mathfrak{G}/\text{Hfrak}} |s(g)|_p |\lambda(g)|_p \leq \|s\|.$$

From the density of  $\text{Step}(\mathfrak{G}, \mathbb{C}_p)$  in  $C(\mathfrak{G}, \mathbb{C}_p)$  and the continuity of the norm function, it follows that

$$\|\lambda\| \leq 1 \quad \lambda(C(\mathfrak{G}, \mathbb{Q}_p)) \subseteq \mathbb{Q}_p.$$

Conversely, let

$$\lambda \in \text{Hom}_{\text{cont}}(C(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p)$$

be such that

$$\|\lambda\| \leq 1 \quad \text{and} \quad \lambda(C(\mathfrak{G}, \mathbb{Q}_p)) \subseteq \mathbb{Q}_p.$$

Then

$$\lambda(\varepsilon_U) \in \mathbb{Q}_p$$

for any  $U \in \mathfrak{B}_{\mathfrak{G}}$  since  $\varepsilon_U \in C(\mathfrak{G}, \mathbb{Q}_p)$ . Moreover,

$$\|\varepsilon_U\| = 1,$$

and  $\|\lambda\| \leq 1$ , so in fact

$$\lambda(\varepsilon_U) \in \mathbb{Z}_p.$$

This concludes the proof. □

If  $\rho : \mathfrak{G} \rightarrow \mathbb{C}_p^\times$  is a character, i.e. a group homomorphism, and  $\lambda, \mu \in \mathfrak{M}(\mathfrak{G}, \mathbb{C}_p)$  then

$$\int_{\mathfrak{G}} \rho d\lambda * \mu = \int_{\mathfrak{G}} \rho d\lambda \int_{\mathfrak{G}} \rho d\mu.$$

A *pseudo-measure* is an element  $\lambda$  of the total ring of fractions of  $\Lambda(\mathfrak{G})$ , i.e. a quotient  $\lambda = \mu/\nu$  of elements  $\mu, \nu \in \Lambda(\mathfrak{G}, \mathbb{Z}_p)$  where  $\nu$  is not a zero divisor, with the property that

$$(g - 1)\lambda \in \Lambda(\mathfrak{G}).$$

For any pseudo-measure  $\lambda$  as above and any non-trivial character  $\rho$  of  $\mathfrak{G}$ , define

$$\int_{\mathfrak{G}} \rho d\lambda := \frac{\int_{\mathfrak{G}} \rho d\mu}{\int_{\mathfrak{G}} \rho d(g - 1)} = \frac{\int_{\mathfrak{G}} \rho d\mu}{\rho(g) - 1},$$

where  $g$  is any element of  $\mathfrak{G}$  not in the kernel of  $\rho$ . This definition does not depend on this choice of  $g$ . Note that we used the fact that for any  $g \in \mathfrak{G}$ ,

$$\int_{\mathfrak{G}} f dg = f(g).$$

In other words, the elements of  $\mathfrak{G}$  correspond to Dirac measures.

#### 1.1.4 The Iwasawa algebra $\Lambda(\mathbb{Z}_p)$ and Mahler's transform

When  $\mathfrak{G} = \mathbb{Z}_p$ , one can say more about  $p$ -adic measures. This is because the  $\mathbb{C}_p$ -Banach algebra of continuous functions on  $\mathbb{Z}_p$  has a special *Mahler basis*.

For  $n \in \mathbb{Z}_{\geq 0}$ , define

$$e_n(x) := \binom{x}{n} = \frac{x(x-1)\dots(x-n+1)}{n!}.$$

**Theorem 1.** *Let  $f \in C(\mathbb{Z}_p, \mathbb{C}_p)$ . Then there exists a unique sequence  $\{a_n\}_{n \geq 0}$  of elements of  $\mathbb{C}_p$  such that*

$$\lim_{n \rightarrow \infty} a_n = 0$$

and

$$f(x) = \sum_{n=0}^{\infty} a_n \binom{x}{n}.$$

*This is called the Mahler expansion of  $f$ .*

*Proof.* This is Theorem 3.3.1 in [CS]. □

Knowing that elements of  $\Lambda(\mathbb{Z}_p)$  can be viewed as continuous linear functional on  $C(\mathbb{Z}_p, \mathbb{C}_p)$ , one could form their generating function with respect to the Mahler basis:

$$\mathcal{M}(\lambda) := \sum_{n=0}^{\infty} T^n \int_{\mathbb{Z}_p} \binom{x}{n} d\lambda.$$

This is called the *Mahler transform* of  $\lambda$ . Note that

$$\mathcal{M}(\lambda) \in \mathbb{Z}_p[[T]].$$

Intuitively, the Mahler transform should determine  $\lambda$  (because the  $e_n(x)$  form a basis of  $C(\mathbb{Z}_p, \mathbb{C}_p)$ ). In fact, much more is true.

**Theorem 2.** *The Mahler transform*

$$\mathcal{M} : \Lambda(\mathbb{Z}_p) \rightarrow \mathbb{Z}_p[[T]]$$

*is an isomorphism of  $\mathbb{Z}_p$ -algebras.*

*Proof.* This is Theorem 3.3.3 in [CS]. □

The inverse of  $\mathcal{M}$ , denoted  $\mathcal{Y}$  in [CS], is defined as follows. If a continuous function  $f$  has Mahler expansion

$$f(x) = \sum_{n=0}^{\infty} a_n e_n(x)$$

and

$$g(T) = \sum_{n=0}^{\infty} b_n T^n \in \mathbb{Z}_p[[T]],$$

we define

$$\int_{\mathbb{Z}_p} f d\mathcal{Y}(g) = \sum_{n=0}^{\infty} a_n b_n.$$

For convenience, we sometimes denote  $\mathcal{Y}(g)$  by  $\lambda_g$ .

**Example:** For any  $a \in \mathbb{Z}_p$ , viewed as a constant compatible sequence in  $\Lambda(\mathbb{Z}_p)$ , one has

$$\mathcal{M}(a) = (1 + T)^a,$$

so that the power series  $(1 + T)^a$  corresponds to the Dirac measures in  $\Lambda(\mathbb{Z}_p)$ .

### 1.1.5 The Iwasawa algebra $\Lambda(\mathbb{Z}_p^\times)$

Integration over  $\mathfrak{G} = \mathbb{Z}_p^\times$  is closely related to integration over  $\mathbb{Z}_p$ . Since  $\Lambda(\mathbb{Z}_p)$  has more structure, it is desirable to relate  $\Lambda(\mathbb{Z}_p^\times)$  to  $\Lambda(\mathbb{Z}_p)$ . Since  $\mathbb{Z}_p^\times$  is a subset of  $\mathbb{Z}_p$ , it is natural to define a map

$$\iota : \Lambda(\mathbb{Z}_p^\times) \rightarrow \Lambda(\mathbb{Z}_p)$$

as

$$\int_{\mathbb{Z}_p} f d\iota(\lambda) := \int_{\mathbb{Z}_p^\times} f|_{\mathbb{Z}_p^\times} d\lambda,$$

for all  $f \in C(\mathbb{Z}_p, \mathbb{C}_p)$ , where  $f|_{\mathbb{Z}_p^\times} \in C(\mathbb{Z}_p^\times, \mathbb{C}_p)$  is the restriction of  $f$  to  $\mathbb{Z}_p^\times$ . One can check that this map is well-defined, i.e. that the functional

$$f \mapsto \int_{\mathbb{Z}_p} f d\iota(\lambda)$$

is in the image of  $\Lambda(\mathbb{Z}_p)$  in  $\lambda \in \text{Hom}_{\text{cont}}(C(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p)$ .

The next step is to identify the image of  $\Lambda(\mathbb{Z}_p^\times)$  inside  $\Lambda(\mathbb{Z}_p)$ . This will be done in the next section, after we introduce the trace and restriction operators.

### 1.1.6 The Iwasawa algebras $\Lambda(\mathcal{G})$ and $\Lambda(G)$

Recall the following notation

$$\mathcal{F}_n = \mathbb{Q}(\mu_{p^{n+1}}) \quad \text{and} \quad F_n = \mathbb{Q}(\mu_{p^{n+1}})^+.$$

$$\mathcal{G} = \text{Gal}(\mathcal{F}_\infty/\mathbb{Q}) \quad \text{and} \quad G = \text{Gal}(F_\infty/\mathbb{Q}).$$

Fixing a generator  $(\zeta_{p^n})$  for the Tate module  $T_p(\mu)$ , we obtain an isomorphism

$$\chi : \mathcal{G} \rightarrow \mathbb{Z}_p^\times,$$

called the cyclotomic character. This induces an isomorphism

$$\tilde{\chi} : \Lambda(\mathcal{G}) \rightarrow \Lambda(\mathbb{Z}_p^\times).$$

But more is true. One can define a natural action of  $\mathcal{G}$  on  $\Lambda(\mathbb{Z}_p^\times)$  and  $\Lambda(\mathbb{Z}_p)$  via the cyclotomic character. Then  $\tilde{\chi}$  becomes a  $\mathcal{G}$ -isomorphism, i.e.  $\tilde{\chi}(\mathcal{G})$  is  $\mathcal{G}$ -equivariant.

For each  $n \geq 0$ , the CM field  $\mathcal{F}_n$  has complex conjugation action  $\iota_n$  and the fixed field of  $\{1, \iota_n\}$  is  $F_n$ . This extends to a complex conjugation action  $\iota$  in  $\mathcal{G}$  which fixes  $G$ . This makes  $\Lambda(\mathcal{G})$  into a  $\mathbb{Z}_p[\mathcal{J}]$ -module, where  $\mathcal{J} = \{1, \iota\}$ . For  $p$  odd decomposes naturally as

$$\Lambda(\mathcal{G}) = \Lambda(\mathcal{G})^+ \oplus \Lambda(\mathcal{G})^-,$$

where

$$\Lambda(\mathcal{G})^+ = \frac{1+\iota}{2}\Lambda(\mathcal{G}) \quad \text{and} \quad \Lambda(\mathcal{G})^- = \frac{1-\iota}{2}\Lambda(\mathcal{G}).$$

Finally, one has the following proposition.

**Proposition 4.** *The restriction to  $\Lambda(\mathcal{G})^+$  of the natural surjection from  $\Lambda(\mathcal{G})$  to  $\Lambda(G)$  induces an isomorphism*

$$\Lambda(\mathcal{G})^+ \simeq \Lambda(G).$$

*Proof.* This is Lemma 4.2.1 of [CS]. □

## 1.2 Operators on $p$ -adic measures

### 1.2.1 Operators on $\mathbb{R}$

### 1.2.2 Operators on $\Lambda(\mathbb{Z}_p)$

### 1.2.3 Restriction of measures from $\mathbb{Z}_p$ to $\mathbb{Z}_p^\times$

## 1.3 Moments of $p$ -adic measures

## 1.4 Summary

1. For a general abelian group, we have

$$\mathfrak{D}(\mathfrak{G}, A) \simeq \Lambda(\mathfrak{G}, A)$$

canonically as abelian groups. When  $A$  is a  $B$ -algebra, this is an isomorphism of  $B$ -algebras.

2. When  $A$  is a  $B$ -algebra, we have

$$\Lambda(\mathfrak{G}, A) \simeq \text{Hom}_{B\text{-alg}}(\text{Step}(\mathfrak{G}, A), A),$$

canonically as  $B$ -algebras, where the product on the right set is the convolution product.

3. When  $A$  is contained in  $\mathbb{C}_p$ , any  $p$ -adic measure  $\lambda$  extends uniquely to a continuous map

$$\lambda : C(\mathfrak{G}, \mathbb{C}_p) \rightarrow \mathbb{C}_p.$$

4. When  $A = \mathbb{Z}_p$ , the image of  $\mathfrak{M}(\mathfrak{G}, \mathbb{Z}_p)$  under the above injection is the set of

$$\lambda \in \text{Hom}_{\text{cont}}(C(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p)$$

such that

$$\|\lambda\| \leq 1 \quad \text{and} \quad \lambda(C(\mathfrak{G}, \mathbb{Q}_p)) \subseteq \mathbb{Q}_p.$$

5. When  $A = \mathbb{Z}_p$  and  $\mathfrak{G} = \mathbb{Z}_p$ , the Mahler transform  $\mathcal{M}$  establishes a  $\mathbb{Z}_p$ -algebra isomorphism

$$\mathcal{M} : \Lambda(\mathbb{Z}_p) \rightarrow \mathbb{Z}_p[[T]].$$

6. When  $A = \mathbb{Z}_p$  and  $\mathfrak{G} = \mathbb{Z}_p^\times$ , we have a natural injection

$$\iota : \Lambda(\mathbb{Z}_p^\times) \rightarrow \Lambda(\mathbb{Z}_p).$$

7. When  $A = \mathbb{Z}_p$  and  $\mathfrak{G} = \mathcal{G}$ , we have a  $\mathcal{G}$ -isomorphism  $\Lambda(\mathcal{G}) \approx \Lambda(\mathbb{Z}_p^\times)$ . Moreover,  $\Lambda(G)$  can be canonically identified as a  $\mathbb{Z}_p$ -submodule of  $\Lambda(\mathcal{G})$ .

Here are all the maps we introduced for  $\mathbb{Z}_p$ -valued measures:

$$\Lambda(G) \simeq \Lambda(\mathcal{G})^+ \hookrightarrow \Lambda(\mathcal{G}) \xrightarrow{\tilde{\iota}} \Lambda(\mathbb{Z}_p^\times) \xrightarrow{\iota} \Lambda(\mathbb{Z}_p) \simeq \mathfrak{D}(\mathbb{Z}_p, \mathbb{Z}_p) = \mathfrak{M}(\mathbb{Z}_p, \mathbb{Z}_p) \hookrightarrow \text{Hom}_{\text{cont}}(C(\mathfrak{G}, \mathbb{C}_p), \mathbb{C}_p).$$

## References

- [CS] Coates, J., Sujatha, R., *Cyclotomic Fields and zeta Values*, Springer Monographs in Mathematics, Springer, 2006.