

# **Notes of**

Wyant College of Optical Sciences  
University of Arizona

Nicolás Hernández Alegría

# Preface

---

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

# Contents

---

Preface

2

# List of Figures

---

# List of Tables

---



# Listings

---

This page is blank intentionally

## Formula sheet

### $z'$ and $m$ curves



### Useful formulas

$$\nu = \frac{V}{\lambda} = \frac{1}{T} = \frac{V}{\lambda}$$

$$\frac{1}{z'} = \frac{1}{z} + \frac{1}{f}$$

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$d = \frac{n-1}{n} t$$

$$(f > 0) |z| \gg f \Rightarrow z' \approx f \wedge m \approx f/z \wedge L = z' - z \approx -z$$

$$(f > 0) |z| \gg f \Rightarrow z \approx -f \wedge m \approx -z'/f \wedge L \approx z'$$

$$\text{Afocal } m = \frac{h'}{h} = \frac{-f_2}{f_1}$$

$$\phi = (n' - n)C$$

$$f = f_E = 1/\phi$$

$$\frac{n'}{z'} = \frac{n}{z} + \phi$$

$$\frac{1}{f} = (n-1)[1/R_1 + 1/R_2 - \frac{(n-1)}{R_1 R_2} \frac{t}{n}]$$

$$n = \frac{c}{V}$$

$$m = \frac{z'}{z} = \frac{h'}{h}$$

$$\theta_2 = -\theta_1$$

$$D \approx -t\theta \frac{n-1}{n}$$

$$\text{Keplerian } m < 0$$

$$C = 1/R$$

$$f_F = -nf_E$$

$$\frac{1}{f} = (n-1)(1/R_1 - 1/R_2)$$

$$\frac{1}{f} = (n-1)[1/R_1 + 1/R_2 - \frac{(n-1)}{R_1 R_2} \frac{t}{n}]$$

$$\text{OPL} = \int_a^b \mathbf{n}(s) \cdot d\mathbf{s}$$

$$m_{\text{total}} = \prod_i m_i$$

$$\theta_i > \theta_c = \sin^{-1} n_2/n_1$$

$$\tau = t/n$$


$$\text{Galilean } m > 0$$

$$n'u' = nu - y\phi$$

$$f'_R = n'f_E$$

$$\frac{1}{f} = (n-1)(1/R_1 - 1/R_2)$$

$$\frac{1}{f} = (n-1)[1/R_1 + 1/R_2 - \frac{(n-1)}{R_1 R_2} \frac{t}{n}]$$



### Key points

- $n$  tells us how much light slows down compared to the vacuum. Frequency doesn't change but wavelength does.
- Fermat's principle states that the path is given by  $\text{OPL}'(\text{path}) = 0$ .
- Reflection is a refraction with negative index  $n' = -n$ .



- Sign convention is: up-right, counter clockwise, vertex-radius of curvature.
- Parity change is preserved only for an **even** number of reflections. It is determined by looking backwards to the object.
- Wherever we have a roof mirror, denoted by a V, we must account for two reflections.
- Reduced thickness is the air-equivalent distance of a medium. All objects are therefore reduced.
- In negative lenses, the rear  $f'_R$  and front  $f_F$  focal points are reversed from positive lenses.
- The  $\text{FOV} = 2\text{HFOV}$  has several definitions, but all are related each other: solid arc can be measured.
- Newtonian equations measure the object and image distances from the focal planes, while Gaussian equations from the principal planes.
- Nodal points  $N$  and  $N'$  preserves magnification of 1.

This page is blank intentionally

