

**Assignment 7**  
**OPTI 570 Quantum Mechanics**  
**University of Arizona**

Nicolás Hernández Alegría

October 15, 2025  
Total time: 15 hours

**Problem I**

a) The evolution operator would be of the form

$$\hat{U}_E(t) = e^{-i\hat{H}_1 t/\hbar} = e^{-i\Omega(\hat{N}^2 - 1/2)t}.$$

The checking is as follows:

$$\begin{aligned}\hat{U}_E\left(\frac{2\pi}{\Omega}\right)|\varphi_n\rangle &= e^{-i\Omega(n^2 - 1/2)\frac{2\pi}{\Omega}}|\varphi_n\rangle \\ &= e^{-i2\pi(n^2 - 1/2)}|\varphi_n\rangle \\ &= (e^{-2\pi})^{n^2} e^{i\pi}|\varphi_n\rangle \\ \hat{U}_E\left(\frac{2\pi}{\Omega}\right)|\varphi_n\rangle &= -|\varphi_n\rangle.\end{aligned}$$

b) For  $\tau = \pi/2\Omega$ , the evolution is

$$\begin{aligned}\hat{U}_E(\tau)|\varphi_n\rangle &= e^{-i\Omega(n^2 - 1/2)\frac{\pi}{2\Omega}}|\varphi_n\rangle \\ &= e^{-i\frac{\pi}{2}n^2} e^{i\frac{\pi}{4}}|\varphi_n\rangle \\ &= (e^{-i\frac{\pi}{2}})^{n^2} e^{i\frac{\pi}{4}}|\varphi_n\rangle \\ &= (-i)^{n^2} e^{i\frac{\pi}{4}}|\varphi_n\rangle \\ \hat{U}_E(\tau)|\varphi_n\rangle &= \begin{cases} e^{i\frac{\pi}{4}} = \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}, & n \text{ even} \\ e^{-i\frac{\pi}{4}} = \frac{1}{\sqrt{2}} - \frac{i}{\sqrt{2}}, & n \text{ odd} \end{cases} |\varphi_n\rangle.\end{aligned}$$

c) We use the fact that in a coherent state, we can express it in terms of the energy eigenstates.

$$|\alpha_0\rangle = e^{-\frac{|\alpha_0|^2}{2}} \sum_{n=0}^{\infty} \frac{\alpha_0^n}{\sqrt{n!}} |n\rangle.$$

We have found that

$$\hat{U}_E(\tau) = \begin{cases} e^{i\frac{\pi}{4}} = \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}, & n \text{ even} \\ e^{-i\frac{\pi}{4}} = \frac{1}{\sqrt{2}} - \frac{i}{\sqrt{2}}, & n \text{ odd} \end{cases}.$$

We then, must split the  $|\alpha_0\rangle$  accordingly, in even and odd term so that the application of the evolution operator gives

$$|\psi_E(\tau)\rangle = e^{-\frac{|\alpha_0|^2}{2}} \left[ e^{i\frac{\pi}{4}} S_{\text{even}} + e^{-i\frac{\pi}{4}} S_{\text{odd}} \right],$$

where

$$S_{\text{even}} = \sum_{n \text{ even}}^{\infty} \frac{\alpha_0^n}{\sqrt{n!}} |n\rangle, \quad \text{and} \quad S_{\text{odd}} = \sum_{n \text{ odd}}^{\infty} \frac{\alpha_0^n}{\sqrt{n!}} |n\rangle. \quad (1)$$

We then have that

$$\left. \begin{aligned} |\alpha_0\rangle &= e^{-\frac{|\alpha_0|^2}{2}} (S_{\text{even}} + S_{\text{odd}}) \\ |-\alpha_0\rangle &= e^{-\frac{|\alpha_0|^2}{2}} (S_{\text{even}} - S_{\text{odd}}) \end{aligned} \right\} \longrightarrow \begin{aligned} S_{\text{even}} &= \frac{1}{2} e^{\frac{|\alpha_0|^2}{2}} (|\alpha_0\rangle + |-\alpha_0\rangle) \\ S_{\text{odd}} &= \frac{1}{2} e^{\frac{|\alpha_0|^2}{2}} (|\alpha_0\rangle - |-\alpha_0\rangle) \end{aligned}.$$

Substituting those in the evolution equation and rearranging:

$$|\psi_E(\tau)\rangle = \frac{1}{2} \left[ (e^{i\frac{\pi}{4}} + e^{-i\frac{\pi}{4}}) |\alpha_0\rangle + (e^{i\frac{\pi}{4}} - e^{-i\frac{\pi}{4}}) |-\alpha_0\rangle \right] = \frac{1}{\sqrt{2}} [\alpha_0 + i |-\alpha_0\rangle],$$

where

$$|\pm \alpha_0\rangle = e^{-\frac{|\alpha_0|^2}{2}} \sum_{n=0}^{\infty} \frac{(\pm \alpha_0)^n}{\sqrt{n!}} |n\rangle.$$

d) The transformation from the Interaction picture to the Schrodinger picture is

$$\begin{aligned} |\psi(\tau)\rangle &= \hat{U}_0(\tau) |\psi_E(\tau)\rangle, \quad \hat{U}_0(\tau) = e^{-iH_0\tau/\hbar} \\ &= e^{-i\omega\tau(\hat{N}+1/2)} |\psi_E(\tau)\rangle \\ &= \frac{1}{\sqrt{2}} e^{-i\omega\tau(\hat{N}+1/2)} [\alpha_0 + i |-\alpha_0\rangle] \\ &= \frac{1}{\sqrt{2}} e^{-\frac{|\alpha_0|^2}{2}} \sum_{n=0}^{\infty} \frac{1}{\sqrt{n!}} \left[ \alpha_0^n e^{-i\omega\tau(n+1/2)} + i(-\alpha_0)^n e^{-i\omega\tau(n+1/2)} \right] |n\rangle \\ &= \frac{1}{\sqrt{2}} e^{-i\frac{\omega}{2}\tau} e^{-\frac{|\alpha_0|^2}{2}} \sum_{n=0}^{\infty} \frac{1}{\sqrt{n!}} \left[ (\alpha_0 e^{-i\omega\tau})^n |n\rangle + i(-\alpha_0 e^{-i\omega\tau})^n |n\rangle \right] \\ |\psi(\tau)\rangle &= \frac{1}{\sqrt{2}} e^{-i\frac{\omega}{2}\tau} [\alpha_0 e^{-i\omega\tau} + i |-\alpha_0 e^{-i\omega\tau}\rangle]. \end{aligned}$$

e) asgagasga

## Problem II

## Problem III


a) sagasg

b) asgasg

- c) asgag
- d) gasgas
- e) asgagasga
- f) asgasg
- g) asgag
- h) asgasg
- i) asgasgasgasg
- j) asgasgasgasg
- k) asgag
- l) asgasg
- m) asfas

## Problem IV

- a) We plot the function  $\text{sech}(x)$  to verify its parity. We can see that it is **even**.



This fact will facilitate us when computing  $\Delta X$ , as we must integrate over  $|\phi(x)|^2$  which therefore, is also even. We then have,

$$\begin{aligned}\langle X \rangle &= \int_{-\infty}^{\infty} x |\phi(x)|^2 dx = \frac{1}{2\beta} \int_{-\infty}^{\infty} x \text{sech}(x/\beta) dx = 0 \\ \langle X^2 \rangle &= \int_{-\infty}^{\infty} x^2 |\phi(x)|^2 dx = \frac{1}{2\beta} \int_{-\infty}^{\infty} x^2 \text{sech}(x/\beta) dx = \frac{\beta^2}{2} \int_{-\infty}^{\infty} u^2 \text{sech}^2(u) du = \frac{\pi^2 \beta^2}{12}.\end{aligned}$$

The  $X$  uncertainty is

$$\Delta X = \sqrt{\langle X^2 \rangle - \langle X \rangle^2} = \frac{\pi\beta}{2\sqrt{3}}.$$

Similarly, for the Fourier transform we have:

$$\begin{aligned}\langle P \rangle &= \int_{-\infty}^{\infty} p |\bar{\phi}(p)|^2 dp = \frac{\pi\beta}{4\hbar} \int_{-\infty}^{\infty} p \text{sech}^2\left(\frac{\pi\beta p}{2\hbar}\right) dp = 0 \\ \langle P^2 \rangle &= \int_{-\infty}^{\infty} p^2 |\bar{\phi}(p)|^2 dp = \frac{\pi\beta}{4\hbar} \int_{-\infty}^{\infty} p^2 \text{sech}^2\left(\frac{\pi\beta p}{2\hbar}\right) dp = \frac{2\hbar^2}{\pi^2 \beta^2} \int_{-\infty}^{\infty} u^2 \text{sech}^2(u) du = \frac{\hbar^2}{\beta^2 3}.\end{aligned}$$

Thus

$$\Delta P = \sqrt{\langle P^2 \rangle - \langle P \rangle^2} = \frac{\hbar}{\beta\sqrt{3}}.$$

The uncertainty product is

$$\Delta X \Delta P = \frac{\pi\beta}{2\sqrt{3}} \frac{\hbar}{\beta\sqrt{3}} = \frac{\hbar\pi}{6}.$$

b) The evolution in  $\pi/2\omega$  gives a well-known quantity, a scaled Fourier transform of the wavefunction.

$$\Phi(x, \frac{\pi}{2\omega}) = U(\frac{\pi}{2\omega}, 0)\Phi(x, 0) = e^{-i\pi/4} \sqrt{\frac{\hbar}{\sigma^2}} \mathcal{F}\{\Phi(x, 0)\} \Big|_{p=\hbar x/\sigma^2}$$

We can see that the function to be computed its Fourier transform is spatially shifted by  $x_0$  so we could directly use the respective property of Fourier transform of a shifter function:

$$\mathcal{F}\{\Phi(x, 0)\} = \bar{\Phi}(p, 0) \implies \mathcal{F}\{\Phi(x - x_0, 0)\} = e^{-ipx_0/\hbar} \bar{\Phi}(p, 0).$$

So,

$$\Phi(x, \frac{\pi}{2\omega}) = -e^{-i\pi/4} \sqrt{\frac{\hbar}{\sigma^2}} \left[ e^{-ipx_0/\hbar} \bar{\Phi}(p, 0) \right] \Big|_{p=\hbar x/\sigma^2} = -\sqrt{\frac{\pi\beta}{4\sigma^2}} e^{-i\pi/4} e^{-i\frac{xx_0}{\sigma^2}} \operatorname{sech}\left(\frac{\pi\beta x}{2\sigma^2}\right).$$

c) To maintain the width  $\Delta X = \frac{\pi\beta}{2\sqrt{3}}$ , we compute  $\Delta X$  for  $\Phi(0, \pi/2\omega)$  and equate it to the uncertainty at  $t = 0$ :

$$\left. \begin{aligned} \langle X \rangle &= 0 \\ \langle X^2 \rangle &= \frac{\pi\beta}{4\sigma^2} \int_{-\infty}^{\infty} x^2 \operatorname{sech}^2\left(\frac{\pi\beta x}{2\sigma^2}\right) dx = \frac{\sigma^4}{3\beta^2}. \end{aligned} \right\} \Delta X = \sqrt{\langle X^2 \rangle} = \frac{\sigma^2}{\sqrt{3}\beta}.$$

Equating it with the uncertainty of the wavefunction at  $t = 0$ :

$$\frac{\pi\beta}{2\sqrt{3}} = \frac{\sigma^2}{\sqrt{3}\beta} \longrightarrow \beta = \sqrt{\frac{2\sigma^2}{\pi}}.$$

## Problem V