

Notes of Quantum Mechanics

Wyant College of Optical Sciences
University of Arizona

Nicolás Hernández Alegría

Preface

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Contents

Preface	2
1 Theory of angular momentum	8
1.1 Spin 1/2 particle: quantization of the angular momentum	9
1.2 Illustration of the postules in the case of a spin 1/2	10
1.3 Two-level systems	11

List of Figures

1.1	9
-----	-------	---

List of Tables

Listings

This page is blank intentionally

Chapter 1

Theory of angular momentum

1.1	Spin 1/2 particle: quantization of the angular momentum	9
1.2	Illustration of the postules in the case of a spin 1/2	10
1.3	Two-level systems	11

1.1 Spin 1/2 particle: quantization of the angular momentum

1.1.1 Experimental demonstration

We are going to describe and analyze the Stern-Gerlach experiment, which demonstrated the quantization of the components of the angular momentum.

The Stern-Gerlach apparatus

The experiment consists of studying the deflection of a beam of neutral paramagnetic atoms (in this case silver atoms). The beam leaves a furnace E through a small opening and propagates in a straight line in the high vacuum existing inside the apparatus. Then, the atomic beam traverses the electromagnet A and thus being deflected before reaching the plate P .

This B-field has a plane of symmetry yOz that contains the initial direction Oy of the atomic beam. The B-field has no components along Oy , and its largest component is along Oz ; it varies strongly with z . Since the B-field has a conserved flux $\nabla \cdot \mathbf{B} = 0$, it must also have a component along Ox which varies with the distance x from the plane of symmetry.




Figure 1.1

1.1.2 Classical calculations of the deflection

The neutral silver atoms possess a permanent magnetic moment μ (they are paramagnetic atoms); the resulting forces are derived from the potential energy:

$$W_B = -\mu \cdot \mathbf{B}. \quad (1.1)$$

For a given atomic level, the magnetic moment μ and the angular momentum \mathbf{J} are proportional:

$$\mu = \gamma \mathbf{J}, \quad (1.2)$$

where γ is the **gyromagnetic ratio** of the level.

1.2 Illustration of the postules in the case of a spin 1/2

1.2.1 Evolution of a spin 1/2 particle in a uniform magnetic field

1.2.2 The interaction Hamiltonian and the Schrodinger equation

Consider a silver atom in a unifrom magnetic field B_0 , and choose the Oz axis along B_0 . The classical potential energy of the magnetic moment $\mu = \gamma J$ of this atom is then:

$$W = -\mu \cdot B_0 = -\mu_z B_0 = \underbrace{-\gamma B_0}_{\omega_0} J_z. \quad (1.3)$$

Since we are quantizing onl the internal degrees of freedom of the particle, J_z must be replaced by the operator S_z , and the clasical energy above becomes an operator: it is the Hamiltonian H which describes the evolution of the spin of the atom in the field B_0 :

$$H = \omega_0 S_z. \quad (1.4)$$

Since H is time-independent, we solve the respective eigenequation. We see that the eigenvectors of H are tose of S_z :

$$H|\pm\rangle = \pm \frac{\hbar\omega_0}{2}|\pm\rangle = E_{\pm}|\pm\rangle. \quad (1.5)$$

There are thetherefore two energy levels, E_{\pm} . Their separation $\hbar\omega_0$ is proportional to the B-field; they define a single Bohr frequency:

$$\nu_{+-} = \frac{1}{\hbar}(E_+ - E_-) = \frac{\omega_0}{2\pi}. \quad (1.6)$$

- If B_0 is parallel to the unit vector \mathbf{u} , the Hmailtonian (1.4) msut be replaced by its general form:

$$\text{General form Hamiltonian} \quad H = \omega_0 \mathbf{S} \cdot \mathbf{u}. \quad (1.7)$$

- For silver atoms, $\gamma < 0$; ω_0 is therefore positive.

Larmor precession

Consider the spin at $t = 0$ in the state

$$|\psi(0)\rangle = \cos \frac{\theta}{2} e^{-i\phi/2} |+\rangle + \sin \frac{\theta}{2} e^{i\phi/2} |-\rangle. \quad (1.8)$$

We saw that any state can be put in this form. To calculate the state at $t > 0$, we apply the evolution operator:

$$|\psi(t)\rangle = \cos \frac{\theta}{2} e^{-i\phi/2} e^{-iE_+t/\hbar} |+\rangle + \sin \frac{\theta}{2} e^{i\phi/2} e^{-iE_-t/\hbar} |-\rangle = \cos \frac{\theta}{2} e^{-\frac{i(\phi+\omega_0 t)}{2}} |+\rangle + \sin \frac{\theta}{2} e^{\frac{i(\phi+\omega_0 t)}{2}} |-\rangle.$$

The presence of B_0 therefore introduces a phase shift between $|+\rangle$ and $|-\rangle$. The direction of $\mathbf{u}(t)$ along which the spin component if $+\hbar/2$ with certainty is defined by the polar angles:

$$\begin{aligned} \theta(t) &= \theta \\ \phi(t) &= \phi + \omega_0 t \end{aligned} \quad (1.9)$$

The angle between $\mathbf{u}(t)$ and Oz therefore remains constant, but $\mathbf{u}(t)$ revolves about Oz at an angular velocity of ω_0 . This effect is called the **Larmor precession**.

It can be verified from $|\psi(t)\rangle$ that the probabilities of obtaining $+\hbar/2$ or $-\hbar/2$ in a measurement of this observable are time-independent. These probabilities are equal, respectively, to $\cos^2 \theta/2$ and $\sin^2 \theta/2$. The mean value of S_z is also time-independent:

$$\langle \psi(t) | S_z | \psi(t) \rangle = \frac{\hbar}{2} \cos \theta. \quad (1.10)$$

Because S_x and S_y do not commute with H , we have that

$$\langle \psi(t) | S_x | \psi(t) \rangle = \frac{\hbar}{2} \sin \theta \cos(\phi + \omega_0 t), \quad \langle \psi(t) | S_y | \psi(t) \rangle = \frac{\hbar}{2} \sin \theta \sin(\phi + \omega_0 t). \quad (1.11)$$

We again find the Bohr frequencies $\omega_0/2\pi$ of the system. Moreover, the mean values above behave like the components of a classical AM of modulus $\hbar/2$ undergoing Larmor precession.

1.3 Two-level systems

This page is blank intentionally

