

QM of angular momentum

Two types of AM: • orbital AM \rightarrow classical analogue
• spin AM \rightarrow intrinsic
 \rightarrow QM, no classical analog

Both described by general theory in QM

Angular momentum in QM

\vec{L} : Orbital angular momentum (OAM)

\vec{S} : Spin angular momentum

\vec{I} : nuclear spin angular momentum

\vec{J} : sum of all \vec{L} OAM and SAM, $\vec{L} + \vec{S}$

\vec{F} : sum of all AM of atom: sum of $\vec{L}, \vec{S}, \vec{I}$
 $\vec{F} = \vec{L} + \vec{S} + \vec{I}$

for now \vec{J} : general angular momentum in QM

Angular momentum in QM:

$\vec{J} \equiv$ any ordered set of three observables $\vec{J} = (J_x, J_y, J_z)$ that satisfy:

$$[J_x, J_y] = i\hbar J_z$$

$$[J_y, J_z] = i\hbar J_x$$

$$[J_z, J_x] = i\hbar J_y$$

$$\vec{J}^2 = \vec{J} \cdot \vec{J} = J_x^2 + J_y^2 + J_z^2$$

Based on definition, J_x, J_y, J_z are not compatible observables

\vec{J} does not have eigenstates

$$[J^2, J_x] = 0$$

$$[J^2, J_y] = 0$$

$$[J^2, J_z] = 0$$

$$\left\{ \begin{array}{l} [J^2, J_u] = 0 \\ \hat{u} \text{ is any direction in space} \end{array} \right.$$

CSCO: $\{J^2, J_u\}$

J^2 : squared magnitude of AM in the system.

J_u : amount of AM about \hat{u}

Find the eigenstates and ass. eigenvalues of J^2 and J_u ?

CT ch VI

New operators:

$$J_+ \equiv J_x + iJ_y$$

$$J_- \equiv J_x - iJ_y$$

$$\rightarrow J_x = \frac{1}{2}(J_+ + J_-)$$

$$J_y = -\frac{i}{2}(J_+ - J_-)$$

Results: eigenstates + eigenvalues of $\{J^2, J_u\}$

$$J^2 |j, m_u\rangle = j(j+1)\hbar^2 |j, m_u\rangle$$

$$J_u |j, m_u\rangle = m_u \hbar |j, m_u\rangle$$

o AM or spin AM

spin only

$\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots$

j : any integer or half integer ≥ 0

m_u : for any given j , $m_u \in \{j, j-1, \dots, -j+1, -j\}$

magnetic quantum number

Convention $\hat{u} \equiv \hat{z}$. Choose a basis $\{|j, m_z\rangle$

magnitude of AM

magnetic quantum number about z

$$J_{\pm} |j, m_z\rangle = \hbar \sqrt{j(j+1) - m_z(m_z \pm 1)} |j, m_z \pm 1\rangle \text{ if } m_z \neq \{j, -j\}$$

$$J_- |j, m_z = -j\rangle = 0$$

$$J_+ |j, m_z = j\rangle = 0$$

$$\Sigma_j: \text{basis } \{|j, m_z\rangle\}$$

standard representation: order elements of basis

$$\text{from } m_z = j, m_z = j-1, \dots, m_z = -j$$

$$\{|j, m_z = j\rangle, |j, m_z = j-1\rangle, |j, m_z = j-2\rangle, \dots, |j, m_z = -j\rangle\}$$

$2j+1$ elements \Rightarrow

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \end{pmatrix} \quad \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \end{pmatrix} \quad \begin{pmatrix} 0 \\ 0 \\ 1 \\ \vdots \end{pmatrix} \quad \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}$$

Σ_x : $\Sigma_j = 3/2$: corresponds to nuclear of Rb^{87} atom

$$m_z \in \{3/2, 1/2, -1/2, -3/2\}$$

standard rep.: $\{|3/2, 3/2\rangle, |3/2, 1/2\rangle, |3/2, -1/2\rangle, |3/2, -3/2\rangle\}$
 $|j, m_z\rangle$

any $|\psi\rangle \in \Sigma_j = 3/2 \quad j = 3/2$

• meas J^2 what are pos. results? $\frac{3}{2} \cdot \frac{5}{2} \cdot \hbar^2 = \frac{15}{4} \hbar^2$

J magnitude of ang. momentum $\sqrt{\frac{15}{4}} \hbar$

• meas J_x - x component of AM, what are possible results?
 $\frac{3}{2} \hbar, \frac{1}{2} \hbar, -\frac{1}{2} \hbar, -\frac{3}{2} \hbar$

• meas J_z - z component of AM, — 11 — ?

$$\frac{3}{2}\hbar, \frac{1}{2}\hbar, -\frac{1}{2}\hbar, -\frac{3}{2}\hbar$$

• $|j = \frac{3}{2}, m_y = \frac{1}{2}\rangle$ - what do I know?
 $J^2 = \frac{15}{4}\hbar^2$

$$\frac{1}{2}\hbar \text{ about } y \text{ axis}$$

$$|j = \frac{3}{2}, m_y = \frac{1}{2}\rangle \neq |j = \frac{3}{2}, m_z = \frac{1}{2}\rangle$$

$$|j, m_x\rangle = \sum_{m_z = -j}^j C_{m_x, m_z} |j, m_z\rangle$$

$$\underline{\Sigma_x}: \quad \Sigma_{j=1}$$

2p state in Hydrogen.

One basis $\{ |j=1, m_z=1\rangle, |j=1, m_z=0\rangle, |j=1, m_z=-1\rangle \}$
 $\quad \quad \quad |z_+\rangle \quad \quad \quad |z_0\rangle \quad \quad \quad |z_-\rangle$

Another basis $\{ |j=1, m_x=1\rangle, |j=1, m_x=0\rangle, |j=1, m_x=-1\rangle \}$
 $\quad \quad \quad |x_+\rangle \quad \quad \quad |x_0\rangle \quad \quad \quad |x_-\rangle$

Consider $|j=1, m_z=0\rangle = |z_0\rangle$

$$\langle \bar{J} \rangle = (\langle \bar{J}_x \rangle, \langle \bar{J}_y \rangle, \langle \hat{J}_z \rangle)$$

$$\langle z_0 | \underline{\hat{J}_z} | z_0 \rangle = 0$$

$$\langle z_0 | \hat{J}_x | z_0 \rangle = \frac{1}{2} \langle z_0 | \hat{J}_+ + \hat{J}_- | z_0 \rangle = 0$$

$$\langle \hat{J}_y \rangle = 0$$

$$\langle \bar{J} \rangle = (0, 0, 0)$$

Consider $|z_+ \rangle = |j=1, m_z=1\rangle$


$$\langle J_z \rangle = \hbar$$

$$\langle J_x \rangle = 0$$

$$\langle J_y \rangle = 0$$

$$\langle \vec{J} \rangle = (0, 0, \hbar) = (\langle J_x \rangle, \langle J_y \rangle, \langle J_z \rangle)$$

$|z_+ \rangle$, $J_z = \hbar$, magnitude of $A\hbar$ is $\sqrt{j(j+1)}\hbar = \sqrt{2}\hbar$
for $j=1$



$$\sqrt{j(j+1)}\hbar^2 = \sqrt{2}\hbar$$