---

title: “RWorksheet_Deluna#3b” author: “Nicole De Luna” date: “2023-10-19” output: pdf_document

## R Markdown

This is an R Markdown document. Markdown is a simple formatting syntax for authoring HTML, PDF, and MS Word documents. For more details on using R Markdown see http://rmarkdown.rstudio.com.

When you click the **Knit** button a document will be generated that includes both content as well as the output of any embedded R code chunks within the document. You can embed an R code chunk like this:

```
summary(cars)
```

```
##      speed           dist
##  Min.   : 4.0   Min.   :  2.00
##  1st Qu.:12.0   1st Qu.: 26.00
##  Median :15.0   Median : 36.00
##  Mean   :15.4   Mean   : 42.98
##  3rd Qu.:19.0   3rd Qu.: 56.00
##  Max.   :25.0   Max.   :120.00
```

## Including Plots

You can also embed plots, for example:




Note that the `echo = FALSE` parameter was added to the code chunk to prevent printing of the R code that generated the plot.

```
#1.a
resp_no <- c(1:20)
sex <- c(2,2,1,2,2,2,2,2,2,2,1,2,2,2,2,2,2,2,1,2)
occ <- c(1,3,3,3,1,2,3,1,1,1,3,2,1,3,3,1,3,1,2,1)
pers_at_home <- c(5,7,3,8,5,9,6,7,8,4,7,5,4,7,8,8,3,11,7,6)
sibs <- c(6,4,4,1,2,1,5,3,1,2,3,2,5,5,2,1,2,5,3,2)
```

```r
h_type <- c(1,2,3,1,1,3,3,1,2,3,2,3,2,2,3,3,3,3,3,2)


household_data <- data.frame(
  Respondents = resp_no,
  Sex = sex,
  FatherOccupation = occ,
  PersonAtHome = pers_at_home,
  SiblingsAtSchool = sibs,
  HouseType = h_type
)
household_data
```

```
##    Respondents Sex FatherOccupation PersonAtHome SiblingsAtSchool HouseType
## 1            1   2                1            5                6         1
## 2            2   2                3            7                4         2
## 3            3   1                3            3                4         3
## 4            4   2                3            8                1         1
## 5            5   2                1            5                2         1
## 6            6   2                2            9                1         3
## 7            7   2                3            6                5         3
## 8            8   2                1            7                3         1
## 9            9   2                1            8                1         2
## 10          10   2                1            4                2         3
## 11          11   1                3            7                3         2
## 12          12   2                2            5                2         3
## 13          13   2                1            4                5         2
## 14          14   2                3            7                5         2
## 15          15   2                3            8                2         3
## 16          16   2                1            8                1         3
## 17          17   2                3            3                2         3
## 18          18   2                1           11                5         3
## 19          19   1                2            7                3         3
## 20          20   2                1            6                2         2
```

```r
#1.2

  str(household_data)
```

```
## 'data.frame':    20 obs. of  6 variables:
##  $ Respondents     : int  1 2 3 4 5 6 7 8 9 10 ...
##  $ Sex             : num  2 2 1 2 2 2 2 2 2 2 ...
##  $ FatherOccupation: num  1 3 3 3 1 2 3 1 1 1 ...
##  $ PersonAtHome    : num  5 7 3 8 5 9 6 7 8 4 ...
##  $ SiblingsAtSchool: num  6 4 4 1 2 1 5 3 1 2 ...
##  $ HouseType       : num  1 2 3 1 1 3 3 1 2 3 ...
```

```r
  summary(household_data)
```

```
##   Respondents         Sex       FatherOccupation  PersonAtHome
##  Min.   : 1.00   Min.   :1.00   Min.   :1.00     Min.   : 3.0
##  1st Qu.: 5.75   1st Qu.:2.00   1st Qu.:1.00     1st Qu.: 5.0
##  Median :10.50   Median :2.00   Median :2.00     Median : 7.0
##  Mean   :10.50   Mean   :1.85   Mean   :1.95     Mean   : 6.4
##  3rd Qu.:15.25   3rd Qu.:2.00   3rd Qu.:3.00     3rd Qu.: 8.0
```

```
##  Max.   :20.00   Max.   :2.00   Max.   :3.00     Max.   :11.0
##  SiblingsAtSchool   HouseType
##  Min.   :1.00     Min.   :1.0
##  1st Qu.:2.00     1st Qu.:2.0
##  Median :2.50     Median :2.5
##  Mean   :2.95     Mean   :2.3
##  3rd Qu.:4.25     3rd Qu.:3.0
##  Max.   :6.00     Max.   :3.0
```

```
# the data frame consists of 20 observations(rows) and 6 variables (columns)

# the variables are:
# respondents - which contains a numeric identifier for each respondent
# sex - represents the gender of the respondent (1 for male, 2 for female)
# father's occupation - indicates the father's occupation (1 for farmer, 2 for driver, 3 for others)
# persons at home - represents the number of people at home
# siblings at school - indicates the number of siblings attending school
# type of house - describes the type of house (1 for wood, 2 for semi-concrete, 3 for concrete)

#1.c
  sibs_mean <- mean(household_data$SiblingsAtSchool)

  sibs_mean
```

```
## [1] 2.95
```

```
# the mean of the number of siblings at school is 2.95, which is not 5

#1.d
  firstTwoRows <- household_data[1:2,]
  firstTwoRows
```

```
##   Respondents Sex FatherOccupation PersonAtHome SiblingsAtSchool HouseType
## 1           1   2                1            5                6         1
## 2           2   2                3            7                4         2
```

```
#1.e
  thirdAndFifthRows <- household_data[c(3,5),c(2,4)]
  thirdAndFifthRows
```

```
##   Sex PersonAtHome
## 3   1            3
## 5   2            5
```

```
#1.f
  types_houses <- household_data$HouseType
  types_houses
```

```
##  [1] 1 2 3 1 1 3 3 1 2 3 2 3 2 2 3 3 3 3 3 2
```

```
#1.g
  male_farmer <- household_data[household_data$Sex == 1 & household_data$FatherOccupation == 1,]

  male_farmer
```

```
## [1] Respondents      Sex              FatherOccupation PersonAtHome
## [5] SiblingsAtSchool HouseType
## <0 rows> (or 0-length row.names)
```

```r
# there are no observations

#1.h

  female_resp <- household_data[household_data$SiblingsAtSchool >= 5,]

  female_resp
```

```
##    Respondents Sex FatherOccupation PersonAtHome SiblingsAtSchool HouseType
## 1            1   2                1            5                6         1
## 7            7   2                3            6                5         3
## 13          13   2                1            4                5         2
## 14          14   2                3            7                5         2
## 18          18   2                1           11                5         3
```

```r
# there are five observations

# -------------------------------------

# 2
  df = data.frame(Ints=integer(),
  Doubles=double(), Characters=character(),
  Logicals=logical(),
  Factors=factor(),
  stringsAsFactors=FALSE
)

  print("Structure of the empty dataframe:")
```

```
## [1] "Structure of the empty dataframe:"
```

```r
  print(str(df))
```

```
## 'data.frame':    0 obs. of  5 variables:
##  $ Ints      : int
##  $ Doubles   : num
##  $ Characters: chr
##  $ Logicals  : logi
##  $ Factors   : Factor w/ 0 levels:
## NULL
```

```r
# df is an empty data frame created with 0 rows and 5 columns
# the columns has the following data type:
# ints = integer
# doubles = double
# characters = character
# logicals = logical
# factors  = factor (0 levels which means empty)

# serves as a template and can be populated with data

# ------------------------------------------------

# 3

  new_resp <- c(1:10)
```

```
  new_sex <- c("Male", "Female", "Female", "Male", "Male", "Female", "Female", "Male", "Female", "Male"
  new_occ <- c(1,2,3,3,1,2,2,3,1,3)
  new_personsAtHome <- c(5,7,3,8,6,4,4,2,11,6)
  new_sibs <- c(2,3,0,5,2,3,1,2,6,2)
  new_type <- c("Wood", "Congrete", "Congrete", "Wood", "Semi-congrete", "Semi-congrete", "Wood", "Semi

  HouseholdData <- data.frame(
  Respondents = new_resp,
  Sex = new_sex,
  FatherOccupation = new_occ,
  PersonAtHome = new_personsAtHome,
  SiblingsAtSchool = new_sibs,
  HouseType = new_type
)

  write.csv(HouseholdData, file = "HouseholdData.csv")

#3a

  imported <-  read.csv("HouseholdData.csv")
  imported
```

```
##     X Respondents    Sex FatherOccupation PersonAtHome SiblingsAtSchool
## 1   1           1   Male                1           5                2
## 2   2           2 Female                2           7                3
## 3   3           3 Female                3           3                0
## 4   4           4   Male                3           8                5
## 5   5           5   Male                1           6                2
## 6   6           6 Female                2           4                3
## 7   7           7 Female                2           4                1
## 8   8           8   Male                3           2                2
## 9   9           9 Female                1          11                6
## 10 10          10   Male                3           6                2
##        HouseType
## 1           Wood
## 2       Congrete
## 3       Congrete
## 4           Wood
## 5  Semi-congrete
## 6  Semi-congrete
## 7           Wood
## 8  Semi-congrete
## 9  Semi-congrete
## 10      Congrete
```

```
#3b

  imported$Sex <- factor(imported$Sex, levels = c("Male", "Female"))
  imported$Sex <- as.integer(imported$Sex)

  imported$Sex
```

```
##  [1] 1 2 2 1 1 2 2 1 2 1
```

```r
#3c

  imported$HouseType <- factor(imported$HouseType, levels = c("Wood", "Congrete", "Semi-congrete"))
  imported$HouseType <- as.integer(imported$HouseType)

  imported$HouseType
```

```
##  [1] 1 2 2 1 3 3 1 3 3 2
```

```r
#3d

  imported$FatherOccupation <- factor(imported$FatherOccupation, levels = c(1,2,3), labels = c("Farmer"
  imported$FatherOccupation
```

```
##  [1] Farmer Driver Others Others Farmer Driver Driver Others Farmer Others
## Levels: Farmer Driver Others
```

```r
#3e

  female_driver <- imported[imported$Sex == 2 & imported$FatherOccupation == "Driver",]
  female_driver
```

```
##   X Respondents Sex FatherOccupation PersonAtHome SiblingsAtSchool HouseType
## 2 2           2   2           Driver            7                3         2
## 6 6           6   2           Driver            4                3         3
## 7 7           7   2           Driver            4                1         1
```

```r
#3f

  greaterFive <- imported[imported$SiblingsAtSchool >= 5,]
  greaterFive
```

```
##   X Respondents Sex FatherOccupation PersonAtHome SiblingsAtSchool HouseType
## 4 4           4   1           Others            8                5         1
## 9 9           9   2           Farmer           11                6         3
```

```r
#4
# On July 14, there were more negative sentiments compared to the other sentiments.This could indicate

# On July 15, all sentiments increased, with the negative sentiment as the highest.This could imply tha

# On July 17 and July 18, the negative sentiments stayed high and the neutral and positive sentiments r

# On July 20, all sentiments got to their lowest with but there were still more negative sentiments tha

# On July 21, experienced an increase in all sentiments, with the negative being the highest. This coul

# From this data, we could assume that public sentiment is responsive to external factors and it also v
```