Strawberries Production in the US

An Exploratory Data Analysis

Nicole Kingdon

October 18, 2023

Table of contents

ntroduction	3
Strawberries & Positive Health	3
Strawberries & Pesticides	3
Organic Strawberries	4
Analyzing Strawberry Production	4
Oata Acquisition & Assessment	6
USDA-NASS Data	6
Census Data	6
Survey Data	7
States	8
Years	8
Assumptions & Motivations	8
Census Data	8
Survey Data	9
Oata Cleaning & Organizing	10
R Packages	10
Initial Data Cleaning & Organization	10
Separating Data Frames	10
Census	11
Survey	12
Exploratory Data Analysis	16
Review of Data Frames	16
	16
	18

References 19

Introduction

Strawberries & Positive Health

Strawberries are a fruit that holds several minerals, vitamins, and nutrients (CRAIG 1997), which have positive implications on human health (Afrin et al. 2016). Specifically, strawberries have been found to help reduce likelihood of cancer, diabetes, obesity, neurodegeneration, cardiovascular disease, and metabolic syndrome (see Figure 1) (Afrin et al. 2016). Although strawberries as a healthy food is the norm, pesticides appear to be harming the beneficial factors of this fruit.




Figure 1: Health benefits of strawberries (Afrin et al., 2016)

Strawberries & Pesticides

Pesticides are used on fruit and vegetable crops, including strawberries, with hopes to increase the quantity (Fenik, Tankiewicz, and Biziuk 2011). Pesticides are made of chemical compounds to reduce or completely eliminate pests from impacting crops (Afrin et al. 2016). These chemical compounds may increase the yield of the crop, but may have a large risk on human health. Additionally, they may contaminate bodies of water and soil with the chemicals, help pests develop resistance to the chemicals, and impact helpful organisms from persisting in areas where pesticides are used. Overall, there are positive and negative impacts of using pesticides (see Figure 2), but it is important to further examine these impacts, specifically on strawberries, to understand the implications of using such.





Figure 2: Positive and negative implications of pesticides on fruit and vegetable crops (Fenik et al., 2011)

Organic Strawberries

Strawberries that are organic are simply strawberries. They use alternative and safer methods to production, and those fruits and vegetables, in this case strawberries, are considered organic (Verteramo Chiu and Gomez 2023). Along with these alternative methods comes a higher cost of production and care, which increases the price to the consumer.



Analyzing Strawberry Production

The literature varies on if organic or non-organic is environmentally better (Afrin et al. 2016). This exploratory data analysis on production of strawberries, which encompasses both pro-

cessed (non-organic) and fresh (organic) market data, will help us better understand production of strawberries in the United States. Additionally, it dives deeper into the pesticide usage in states, which will provide more information as to the level of toxic ingredients that go into the production of non-organic (or conventional) strawberries .

Data Acquisition & Assessment

USDA-NASS Data

The data was acquired from U.S. Department of Agriculture (USDA) and the National Agricultural Statistics Service (NASS). The data was uploaded for data cleaning and organizing and exploratory data analysis by Professor Haviland Wright, who chose the following data: USDA-NASS.

The data frame loaded into the environment is titled strawberry.

```
Rows: 4,314
Columns: 21
             <chr> "CENSUS", "CENSUS", "CENSUS", "CENSUS", "~
$ Program
$ Year
             <dbl> 2021, 2021, 2021, 2021, 2021, 2021, 2021, 2021, 202
             <chr> "YEAR", "YEAR", "YEAR", "YEAR", "YEAR", "YEAR", "YE~
$ Period
$ `Week Ending`
             <chr> "STATE", "STATE", "STATE", "STATE", "STATE", "STATE"
$ `Geo Level`
             <chr> "ALASKA", "ALASKA", "ALASKA", "ALASKA", "ALASKA", "~
$ State
$ `State ANSI`
             <chr> "02", "02", "02", "02", "02", "02", "02", "06", "06~
$ `Ag District`
             $ County
$ `County ANSI`
             $ `Zip Code`
             $ Region
$ watershed_code
             $ Watershed
$ Commodity
             <chr> "STRAWBERRIES", "STRAWBERRIES", "STRAWBERRIES", "ST~
$ `Data Item`
             <chr> "STRAWBERRIES, ORGANIC - OPERATIONS WITH SALES", "S~
             <chr> "ORGANIC STATUS", "ORGANIC STATUS", "ORGANIC STATUS~
$ Domain
             <chr> "ORGANIC STATUS: (NOP USDA CERTIFIED)", "ORGANIC ST~
$ `Domain Category`
             <chr> "2", "(D)", "(D)", "(D)", "2", "(D)", "(D)", "142",~
$ Value
$ `CV (%)`
             <chr> "(H)", "(D)", "(D)", "(H)", "(D)", "(D)", "1~
```

Census Data

The data offers census data based on state that represents fresh market (organic) and process market (non-organic) sales.

Rows: 864

```
Columns: 21
             <chr> "CENSUS", "CENSUS", "CENSUS", "CENSUS", "~
$ Program
$ Year
             <dbl> 2021, 2021, 2021, 2021, 2021, 2021, 2021, 2021, 202
$ Period
             <chr> "YEAR", "YEAR", "YEAR", "YEAR", "YEAR", "YEAR", "YE~
             $ `Week Ending`
$ `Geo Level`
             <chr> "STATE", "STATE", "STATE", "STATE", "STATE", "STATE"
$ State
             <chr> "ALASKA", "ALASKA", "ALASKA", "ALASKA", "ALASKA", "~
             <chr> "02", "02", "02", "02", "02", "02", "02", "06", "06~
$ `State ANSI`
$ `Ag District`
             $ County
$ `County ANSI`
             $ `Zip Code`
$ Region
             $ watershed_code
$ Watershed
             $ Commodity
             <chr> "STRAWBERRIES", "STRAWBERRIES", "STRAWBERRIES", "ST~
$ `Data Item`
             <chr> "STRAWBERRIES, ORGANIC - OPERATIONS WITH SALES", "S~
             <chr> "ORGANIC STATUS", "ORGANIC STATUS", "ORGANIC STATUS~
$ Domain
             <chr> "ORGANIC STATUS: (NOP USDA CERTIFIED)", "ORGANIC ST~
$ `Domain Category`
             <chr> "2", "(D)", "(D)", "(D)", "2", "(D)", "(D)", "142",~
$ Value
$ `CV (%)`
             <chr> "(H)", "(D)", "(D)", "(D)", "(H)", "(D)", "(D)", "1~
```

Survey Data

Additionally, it holds survey information for each state, specifically indicating pesticides and bacterium used to preserve strawberry crop yield. In addition, it offers fresh and process market data.

```
Rows: 3,450
Columns: 21
             <chr> "SURVEY", "SURVEY", "SURVEY", "SURVEY", "SURVEY", "~
$ Program
$ Year
             <dbl> 2022, 2022, 2022, 2022, 2022, 2022, 2022, 2022, 202
             <chr> "MARKETING YEAR", "MARKETING YEAR", "MARKETING YEAR~
$ Period
             $ `Week Ending`
$ `Geo Level`
             <chr> "STATE", "STATE", "STATE", "STATE", "STATE", "STATE"
             <chr> "CALIFORNIA", "CALIFORNIA", "CALIFORNIA", "FLORIDA"~
$ State
$ `State ANSI`
             <chr> "06", "06", "06", "12", "12", "12", NA, NA, NA, "06~
$ `Ag District`
             $ County
             $ `County ANSI`
```

```
$ `Zip Code`
            $ Region
            $ watershed_code
            $ Watershed
            <chr> "STRAWBERRIES", "STRAWBERRIES", "STRAWBERRIES", "ST~
$ Commodity
            <chr> "STRAWBERRIES - PRICE RECEIVED, MEASURED IN $ / CWT~
$ `Data Item`
$ Domain
            <chr> "TOTAL", "TOTAL", "TOTAL", "TOTAL", "TOTAL", "TOTAL",
$ `Domain Category`
            <chr> "NOT SPECIFIED", "NOT SPECIFIED", "NOT SPECIFIED", ~
            <chr> "108", "(D)", "(D)", "169", "(D)", "(D)", "0", "135~
$ Value
            $ `CV (%)`
```

States

There were 47 states (c("ALASKA", "CALIFORNIA", "CONNECTICUT", "FLORIDA", "GEORGIA", "IDAHO", "ILLINOIS", "INDIANA", "IOWA", "KENTUCKY", "LOUISIANA", "MAINE", "MARYLAND", "MASSACHUSETTS", "MICHIGAN", "MINNESOTA", "MONTANA", "NEBRASKA", "NEW HAMPSHIRE", "NEW JERSEY", "NEW YORK", "NORTH CAROLINA", "OHIO", "OKLAHOMA", "OREGON", "PENNSYLVANIA", "RHODE ISLAND", "SOUTH CAROLINA", "SOUTH DAKOTA", "TENNESSEE", "VERMONT", "WASHINGTON", "WEST VIRGINIA", "WISCONSIN", "ALABAMA", "ARIZONA", "COLORADO", "KANSAS", "MISSOURI", "NEVADA", "NEW MEXICO", "VIRGINIA", "ARKANSAS", "NORTH DAKOTA", "TEXAS", "UTAH", "OTHER STATES")) with two states considered as "other states".

Years

The data was from the years c(2021, 2019, 2016, 2022, 2020, 2018, 2017).

Assumptions & Motivations

Census Data

The census data was a nation-wide collection of data about the fresh and process markets related to strawberries. This data has values that are indicated as (D), which are data that was withheld upon request by the strawberry market in that particular state. This could leave out important information in the data.

Survey Data

The survey data was collected via a survey sent out to each state in the United States. There were only 11 out of 47 states who returned the survey (c("CALIFORNIA", "FLORIDA", "OTHER STATES", "NEW YORK", "NORTH CAROLINA", "OREGON", "WASHINGTON", "MICHIGAN", "OHIO", "PENNSYLVANIA", "WISCONSIN")), which includes the "other states". (The "other states" did not have any data relating to pesticides and bacterium.) This is only a 23% response rate, which is not comprehensive of all the states and the entire United States process market. The states that did return the survey will still be able to show a report of pesticide and bacterium usage on their processed strawberry crops.

Data Cleaning & Organizing

R Packages

The following R packages were used to clean and organize the data:

```
library(knitr)
library(kableExtra)
library(tidyverse)
library(stringr)
library(dplyr)
```

The data was organized into two main categories: census and survey. The census data was cleaned and organized to show fresh and process market sales, and the survey data consisted of pesticide and fertilizer usage.

Initial Data Cleaning & Organization

The following initial data cleaning derived from Professor Wright.

Removed columns with a single value in all columns using a function. This function produced all columns with one value from the **strawberry** data frame. From there, all but the single value columns were added back to the data frame.

Next, Professor Wright checked if every row in the state column held a value, which it did.

After that, Professor Wright checked to see what state had the most rows.

The state with the most rows is CALIFORNIA.

Separating Data Frames

The following separation of data frames derived from Professor Wright.

By using dplyr, the strawberry data frame was split by Program. The rows that contained CENSUS were moved to strwb_census, and the rows containing SURVEY were moved to strwb_survey.

Census

After splitting CENSUS and SURVEY rows into two data frames, Professor Wright has first organized the CENSUS data.

First, he separated composite columns and cleaned the Value column.

The composite columns in the strwb_census are as follows: Data Item and Domain category.

The column separators in CENSUS are ",", "-", ":".

Separated Data Item into the new columns Fruit, temp1, temp2, and temp3 by ",".

Separated temp1 into the new columns crop_type and prop_acct by "-".

To finish this first section of cleaning and organizing the strwb_census data frame, Professor Wright string trimmed both sides of the following three columns: crop_type, temp2, and temp3.

Now, Professor Haviland created a "Fresh Market" column. To do this, he duplicated the temp2 column with the new column name as Fresh Market. Next, he removed all the cells in the new column that began with MEASURED. Same for the cells that begin with PROCESSING. He substituted NA values for empty strings. Finally, the FRESH MARKET parts of the strings were removed, as they were unneccesary anymore with the new column creation of Fresh Market.

Now, to manipulate the temp2 column, Professor Wright removed all the cells that began with FRESH, which would now set up the cleaning and organizing to create the Process Market column.

Professor Wright created a "Process Market" column.

To do this, he followed the same method as Fresh Market, where he first duplicated temp2, then removed cells beginning with MEASURED, and removed PROCESSING from the beginning of the strings. and remove

Finally, removed the cells starting with PROCESS MARKET from temp2.

Next, Professor Wright removed NA's from prop_acct, temp2, and temp3 by substituting a space.

From here, he combined temp2 with temp3 to create a Metric column. He also removed parts of string that did not matter, beginning with MEASURED IN. To finish, he moved the Metric column to the end.

The Value column was transformed. To do this, Professor Wright first pulled the Value column from strwb_census and put them into vals. From there, he string replaced all vals to remove

the commas in the strings. From there, he converted the strings to numeric values. During this, NA values were automatically implemented.

After this, he aimed to find the location and value of the footnotes in the Value column. He implemented a string detection on this column, which helped discover that all the cells with NA values were also where the footnotes were located. The footnotes indicated that the Value was instructed to be left out during the data collection.

I finished cleaning and organizing the strwb_census data frame, which is detailed below.

First, I selected particular columns that had necessary data.

Next, I removed the "," from the Value column and transformed them into numeric values. This introduced rows with NA values.

After that, I cleaned up the CV (%) column by changing the values to numbers, instead of strings. This also introduced rows with NA values.

Furthermore, I am going to omit all rows in the Value and CV (%) columns with NA values. These are being omitted because they hold no meaning. Only the Value column had to be adjusted, as the CV (%) column did not have any values if the Value column also did not.

Finally, to complete the strwb_census cleaning and organizing, I am going to arrange the State column to be in ascending order.

Survey

Professor Wright had organized the SURVEY data frame splitting the marketing, and production data from the chemical application data. In the strawberry data frame, The CENSUS rows contain marketing, sales, and production data. The SURVEY rows contain rows which may be redundant with the CENSUS rows and chemical application rows. These rows contain fresh and process market sales data, which have been removed.

Began cleaning and organizing strwb_survey by discovering what columns in this data frame that need to be split.

First, Professor Wright separated the Data Item column to temp1, temp2, temp3, and temp4 by ",". Additionally, he separated temp1 into temp1a and temp1b.

Next, he separated the Domain column into temp22 and temp23 by ",".

Also, he separated Domain Category into temp42 and temp43 by ",".

To finish the strwb_survey cleaning and organizing, this data frame was split into two new data frames, where strwb_survey_chem holds pesticide data and strwb_survey_mkt contains all the surveyed market and fertilizer usage data.

Survey: Market

Now, I further cleaned up both strwb_survey_mkt and strwb_survey_chem. First, I worked with the strwb_survey_mkt.

Quickly, to begin, Professor Wright Dropped one-value columns in strwb_survey_mkt.

To begin, I made the Value column into numeric values, which introduced NA values. Then, I changed the format of the numeric values, so they did not appear in scientific notation.

To reduce the duplicity of strwb_survey_mkt with strwb_census, I separated the fresh and process market data in strwb_survey_mkt from the data on the fertilizer usage.

Next, I separated the temp42 column to create the columns type and fertilizer_type by the delimiter ":".

I cleaned up the fertilizer_type strings by removing the parentheses on both sides.

I also removed the empty spaces in these strings.

Finally, I selected the rows that are relevant and renamed temp3 to measurement and temp4 to avg.

Survey: Chemical

Finally, I finished cleaning and organizing strwb_survey_chem.

First, Professor Haviland dropped one-value columns in strwb_survey_chem.

Then, I selected the relevant columns.

From here, I split up temp43 into chemical and PC# by "=".

Furthermore, I cleaned up both of the new chemical and the PC# columns by removing the unnecessary parentheses.

I also want to separate the chemical column by chemical type and chemical name.

Trimmed off the spaces on both sides of the chemical column.

Trimmed the same to chem_type.

Finally, trimmed the same with PC#.

Now to remove the rows with NA values in the PC# column.

Addition to Strwb_Survey_Chem: WHO Chemical Toxicity

Now, to better understand the strwb_survey_chem data, I added two columns of data. We are first going to begin with "chemical toxicity" (toxicity).

To gather this information, I used the World Health Organization (WHO)'s classification of pesticides by hazard [WHO, 2019]. Table 1 shows the WHO's toxicity classification for pesticides.

Class	Description	LD ₅₀ for the Rat (mg/kg Body Weight)			
		Oral		Dermal	
		Solids	Liquids	Solids	Liquids
la	Extremely hazardous	≤5	≤20	≤10	≤40
lb	Highly hazardous	5–50	20–200	10–100	40–400
II	Moderately hazardous	50–500	200–2,000	100–1,000	400–4,000
III	Slightly hazardous	>500	>2,000	>1,000	>4,000

doi:10.1371/journal.pmed.1000357.t001

Figure 3: Table 1. World Health Organization Pesticide Toxicity Classification

Additionally, some chemicals are presented as fatal or toxic if inhaled, as they are gaseous or volatile fumigants [WHO, 2019]. Others are classified as "unlikely to present acute hazard" by WHO [WHO, 2019], which means that they will not present any hazard if used properly. Furthermore, they are also classified as "no significant acute toxicity" when they are not in the WHO classification and are found to be non-toxic, mostly discovered through the Environmental Protection Agency (EPA) [EPA].

I classified each chemical toxicity by the string Highly hazardous, Moderately hazardous, Unlikely to present acute hazard, Slightly hazardous, Fatal if inhaled, Toxic if inhaled, No significant acute toxicity and Not specified, based on the WHO chemical toxicity rating system [WHO, 2019].

I searched through each classification table to find each pesticide. Not all pesticides were on the table. To find the missing chemical toxicities, I used the large language model, Chat GPT [Chat GPT], found information through EPA Pesticide Fact Sheets [EPA-Acibenzolar], [EPA-Ammonium.], [EPA-Aureobasidium.], [EPA-Canola], [EPA-Caprylic], [EPA-Capasaicin], [EPA-Clethodim], [EPA-Cyfluefenamid], [EPA-Cytokinin],

[EPA-Indole.], [EPA-Iron-Phos.], [EPA-Mefenoxam], [EPA-Metam-sodium], [EPA-Polyoxin], [EPA-Potassium], [EPA-Potassium-salts], [EPA-Potassium-silicate], [EPA-Spiromedifen], [EPA-STREPTOMYCES-LYDICUS], [EPA-SULFENTRAZONE], [EPA-SULFUR], and in other sources (Kilani-Morakchi, Morakchi-Goudjil, and Sifi 2021), [PubChem-Capric], [Carfentrazone-ethyl], [Pub-Chem-Copper.], [Copper-Octanoate], [ACS-Copper-Oxide], [Cyprodinil], [DECYLDIMETHYLOCTYL], [Didecyl.], [Dodine], [Sodium-Ferric-Ethyl.], [Garlic-Oil], [Glyphosate], [Hydrogen-Peroxide], [Isofetamind], [Methoxyfenozide], [Mineral-oil], [Mono-Potassium-Salt], [Mustard-oil], [Peroxyacetic-acid], [Quinoline], [CDC-PYRACLOSTROBIN], [SOYBEAN-OIL].

Furthermore, through this search, I discovered through Chat GPT [Chat GPT], that many of the "chemicals" were actually bacterium. I found the toxicity of the bacterium via web search from various sources [BACILLUS AMYLOLIQUEFAC F727.], [BACILLUS AMYLOLIQUEFACIENS MBI 600], [BACILLUS AMYLOLIQUEFACIENS STRAIN D747], [BACILLUS PUMILUS], [BACILLUS SUBTILIS], [ALKYL. DIM. BENZ. AM], [AUREOBASIDIUM PULLULANS DSM 14940], [BACILLUS SUBT. GB03], [BEAUVERIA BASSIANA], [BLAD], [BT KURSTAK ABTS-1857], [BT], [BURKHOLDERIA A396 CELLS & MEDIA], [CAPSICUM OLEORESIN EXTRACT], [CHROMOBAC SUBTSUGAE PRAA4-1 CELLS AND SPENT MEDIA], [GLIOCLADIUM VIRENS], [HARPIN A B PROTEIN], [HELICOVERPA ZEA NPV], [NEEM OIL], [PAECILOMYCES FUMOSOR], [PETROLEUM DISTILLATE], [PSEUDOMONAS CHLORORAPHIS STRAIN AFS009], [REYNOUTRIA SACHALINE], [TRICHODERMA HARZ.], [TRICHODERMA VIRENS STRAIN G-41].

There were three Unknown values for COPPER ETHANOLAMINE, KANTOR, and HALOSULFURON-METHYL, which did not have any clear information on toxicity.

Addition to Strwb_Survey_Chem: CAS Registry Number

Now, the second added column represents each pesticide's Chemical Abstract Service (CAS) Registry Number (CAS#). A CAS Registry Number allows each chemical compound, including molecular formulas, chemical structures, generic, systematic, common, and trade names, to have a clear identification number [CAS].

To find the CAS#s of each pesticide, I used [WHO, 2019], [CHAT GPT], and, mainly, the United States Environmental Protection Agency's Pesticide Chemical Search [EPA-search]. After discovering that some of the rows in chemical are actually bacterium, all values that do not have CAS#s are indicated as Bacteria. All the other chemicals were matched with their appropriate CAS#.

Quickly, I reorganized strwb_survey_chem to have the columns in a different order.

I also changed the column name from chemical to strwb_treatment, as bacterium are not chemicals.

Finally, to finish the data cleaning and organization of strwb_survey_chem, I changed the name of the data frame to strwb_survey_treat (treat = treatment) to avoid confusion, since it is known that all strawberry pesticide treatments are not only chemicals but also bacterium.

Exploratory Data Analysis

Now, to begin the Exploratory Data Analysis of the USDA-NASS data set. First, I have reviewed each data frame associated with the data set. Next, I searched for and examined any missing values in the data frames. After that, I categorized the values in the data set to prepare for visualization. Finally, I analyzed the relationships in the data frames through visualization and located any outliers [EDA-Advice].

Review of Data Frames

strwb_census

Columns & Values

The column names in strwb_census are as follows: Program, Year, State, Fruit, crop_type, Totals, Fresh Market, Process Market, Metric, Value, and CV (%).

Program

Program has a single string of CENSUS in every column. CENSUS represents that the data derived from a census of states based on their strawberry production.

Year

Year represents the years in which the data was collected. The years are as follows: c(2019, 2021, 2016).

State

State holds all the US states that held any census data on their strawberry production (c("ALABAMA", "CALIFORNIA", "COLORADO", "CONNECTICUT", "FLORIDA", "GEORGIA", "IDAHO", "ILLINOIS", "IOWA", "KANSAS", "KENTUCKY", "LOUISIANA", "MAINE", "MARYLAND", "MASSACHUSETTS", "MICHIGAN", "MONTANA", "NEW HAMPSHIRE", "NEW JERSEY", "NEW MEXICO", "NEW YORK", "NORTH CAROLINA", "OHIO", "OKLAHOMA", "OREGON", "PENNSYLVANIA", "SOUTH CAROLINA", "VERMONT", "WISCONSIN")).

Fruit

Fruit has a single string of STRAWBERRY, which shows that all the data is based on this fruit.

crop_type

crop_type also has a single string, ORGANIC, which shows that the entirety of the production data is based on organic strawberries.

Totals

Totals holds three values: c("OPERATIONS WITH SALES", "SALES", "","PRODUCTION"). The SALES string reprents the sales of strawberries, OPERATIONS WITH SALES represents how many operating strawberry production with intent to sell, and PRODUCTION represents the amount of strawberries produced.

Fresh Market

Fresh Market holds two values: OPERATIONS WITH SALES and SALES. The definition of both of these values are the same as the Totals column.

Process Market

Process Market has the same strings as Fresh Market.

Metric

Metric has the following strings: c("","\$","CWT").

The empty string holds no values because that row did not need a metric to help describe the data. \$ represents that the metric for the following Value column is in US Dollars. CWT is weight in 100's

[CWT].

Value

The Value column holds numeric values that correspond with the strings in Totals, Fresh Market, and Process Market, as well as the Metric column.

CV (%)

Finally, the CV (%) column holds the co-efficients of variance of the numeric values in the Value column.

Missing Values

All rows with missing values in the Value column were omitted. The missing values represented values that were omitted by the state upon census data collection. These values have no purpose in the data analysis.

Categorizing Values

All columns, but Value and CV (%) hold nominal variables. Both the Value column and CV (%) hold interval variables.

Data Visualization

Questions


After analyzing the columns and values of strwb_census , I developed the following questions:

- 1. What state held the most operational strawberry sellers?
- 2. What states made the most money producing strawberries?
- 3. What states produced the most strawberries?

These questions will be individually answered below via data visualization.

What state held the most operational strawberry sellers?





References

Afrin, Sadia, Massimiliano Gasparrini, Tamara Y. Forbes-Hernandez, Patricia Reboredo-Rodriguez, Bruno Mezzetti, Alfonso Varela-López, Francesca Giampieri, and Maurizio Battino. 2016. "Promising Health Benefits of the Strawberry: A Focus on Clinical Studies." *Journal of Agricultural and Food Chemistry* 64 (22): 4435–49. https://doi.org/10.1021/acs.jafc.6b00857.

CRAIG, WINSTON J. 1997. "Phytochemicals." Journal of the American Dietetic Association 97 (10): S199–204. https://doi.org/10.1016/s0002-8223(97)00765-7.

Fenik, Jolanta, Maciej Tankiewicz, and Marek Biziuk. 2011. "Properties and Determination of Pesticides in Fruits and Vegetables." *TrAC Trends in Analytical Chemistry* 30 (6): 814–26. https://doi.org/10.1016/j.trac.2011.02.008.

Kilani-Morakchi, Samira, Houda Morakchi-Goudjil, and Karima Sifi. 2021. "Azadirachtin-Based Insecticide: Overview, Risk Assessments, and Future Directions." Frontiers in Agronomy 3 (July). https://doi.org/10.3389/fagro.2021.676208.

Verteramo Chiu, Leslie J., and Miguel I. Gomez. 2023. "A Tale of Two Strawberries: Conventional and Organic Open-Field Production in California." *Sustainability* 15 (19): 14363. https://doi.org/10.3390/su151914363.