

# Napkin exercises — solutions

Jakub Kądziołka

January 4, 2021

## Contents

```
theory Chapter1
  imports HOL-Algebra.Algebra
begin
```

It seems that one cannot use  $\heartsuit$  as variable. My disappointment is immeasurable and my day is ruined.

```
theorem problem1A:
  assumes group: group G
  assumes subgrp: subgroup H G
  assumes proper:  $H \subset \text{carrier } G$ 
  assumes iso:  $G \cong G(\text{carrier} := H)$ 
  shows infinite (carrier G)
proof
  from iso obtain h where bij: bij-betw h (carrier G) H
    unfolding is-iso-def iso-def by auto

  assume finite: finite (carrier G)
  with proper have  $\text{card } H < \text{card } (\text{carrier } G)$ 
    by (simp add: psubset-card-mono)
  moreover from finite and bij have  $\text{card } H = \text{card } (\text{carrier } G)$ 
    using bij-betw-same-card by fastforce
  ultimately show False by auto
qed
```

Problem 1B asks for a special case of Lagrange's theorem, thus we avoid using the general variant.

```
theorem (in comm-group) problem1B:
  assumes finite: finite (carrier G)
  assumes closed:  $g \in \text{carrier } G$ 
  shows  $g \text{ [} \wedge \text{] order } G = 1$ 
proof –
  let  $?f = \lambda x. g \otimes x$ 
  have [simp]:  $?f \text{ ' carrier } G = \text{carrier } G$ 
    by (simp add: closed group.surj-const-mult)
  have inj-on  $?f$  (carrier G)
```

```

    by (simp add: closed group.inj-on-cmult)
  hence  $(\bigotimes x \in \text{carrier } G. x) = (\bigotimes x \in \text{carrier } G. g \otimes x)$ 
  using finprod-reindex[where  $h=?f$  and  $A=\text{carrier } G$  and  $f=\lambda x. x, \text{ symmetric}$ ]
  by simp
  also have  $\dots = (\bigotimes x \in \text{carrier } G. g) \otimes (\bigotimes x \in \text{carrier } G. x)$ 
  using closed by (intro finprod-multf) auto
  finally have  $(\bigotimes x \in \text{carrier } G. g) = 1$ 
  using closed by (intro r-cancel-one'[THEN iffD1]) auto
  thus ?thesis
  using closed unfolding order-def by (simp add: finprod-const)
qed

end

```