```
theory Warmup-Problem-C
 imports
   Complex-Main
  HOL-Library.\ Quadratic-Discriminant
begin
```

## Warmup problem C 0.1

end

This one is a straight-forward equation:

```
theorem warmup3:
 |x-1|*|x+2|*|x-3|*|x+4| = |x+1|*|x-2|*|x+3|*|x-4|
   \longleftrightarrow x \in \{0, sqrt \ 7, -sqrt \ 7,
              sqrt ((13 + sqrt 73) / 2),
             -sqrt((13 + sqrt 73) / 2),
              sqrt ((13 - sqrt 73) / 2),
             -sqrt ((13 - sqrt 73) / 2)
 (is ?egn \leftrightarrow ?sols)
proof -
 have ?eqn \longleftrightarrow |(x-1)*(x+2)*(x-3)*(x+4)| = |(x+1)*(x-2)*(x+3)*(x-4)| (is -\longleftrightarrow |?lhs|
= |?rhs|
   by (simp add: abs-mult)
 also have ... \longleftrightarrow ?lhs - ?rhs = 0 \lor ?lhs + ?rhs = 0 by (auto simp add: abs-eq-iff)
 also have ... \longleftrightarrow x*(x^2 - 7) = 0 \lor x^4 - 13*x^2 + 24 = 0 by algebra
 also have x*(x^2 - 7) = 0 \longleftrightarrow x \in \{0, sqrt 7, -sqrt 7\} using plus-or-minus-sqrt by auto
 also have x^4 - 13 * x^2 + 24 = 0 \longleftrightarrow x^2 \in \{(13 + sqrt 73) / 2, (13 - sqrt 73) / 2\}
   using discriminant-nonneg [where x=x^2, of 1 -13 24]
   by (auto simp add: algebra-simps discrim-def)
 also have ... \longleftrightarrow x \in \{sqrt ((13 + sqrt 73) / 2),
                      -sqrt ((13 + sqrt 73) / 2),
                      sqrt ((13 - sqrt 73) / 2),
                      -sqrt ((13 - sqrt 73) / 2)
 proof -
   have 0 \le (13 - sqrt 73) / 2 by (auto simp add: real-le-lsqrt)
   hence x^2 = (13 - sqrt 73) / 2
         \longleftrightarrow x \in \{sqrt \ ((13 - sqrt \ 73) \ / \ 2),\
                 -sqrt ((13 - sqrt 73) / 2)
     using plus-or-minus-sqrt
     by blast
   moreover have x^2 = (13 + sqrt 73) / 2
     \longleftrightarrow x \in \{sqrt ((13 + sqrt 73) / 2),
            -sqrt ((13 + sqrt 73) / 2)
      by (smt insert-iff power2-minus power-divide real-sqrt-abs real-sqrt-divide real-sqrt-pow2
singletonD)
   ultimately show ?thesis by blast
 qed
 ultimately show ?thesis by blast
qed
```