

 $ENGR142 \quad {\it 2018, 2nd Trimester}$

Lecturers: B. Ruck, F. Natali, and C. Hollitt

Assignment 2 Due date: 11:59 PM, Thursday 2nd August, 2018

Problem 1: Projectile motion

(6 Marks)


Famous American football player Cam Newton throws a football at a speed of 27 ms⁻¹ and at an angle of 10° above the horizontal. He releases it at a height of 2 m above the ground. A defender 10 m away jumps to intercept the ball. If the defender can reach to 3.5 m above the ground are they able to intercept the pass? Assume the only force on the ball is gravity.

Problem 2: Air hockey table

(7 Marks)

Two pucks, with masses $m_1 = 0.1$ kg and $m_2 = 0.1$ kg, slide without friction on an air hockey table. They approach each other and collide, as shown below. Before the collision the velocity vector of m_1 is $\vec{v}_{1i} = (5\hat{i} + 0\hat{j})$ ms⁻¹ and the velocity vector of m_2 is $\vec{v}_{2i} = (-2\hat{i} + 2\hat{j})$ ms⁻¹. After the collision the velocity vector of m_1 is $\vec{v}_{2f} = (2\hat{i} + 3\hat{j})$ ms⁻¹.

- (a) Find the final velocity vector for m_2 .
- (b) Was the collision elastic or inelastic?



Problem 3: Terminal velocity

(7 Marks)

A hailstone of mass $m=5\times 10^{-4}$ kg falls under the effect of gravity. It also experiences a drag force of magnitude $F_D=\eta v^2$, where v is the hailstone's velocity and the coefficient η for the particular hailstone under consideration is equal to $\eta=2\times 10^{-5}$ kg/m.

- (a) Calculate the terminal velocity of the hailstone.
- (b) If the hailstone falls from a height h = 500 m, starting at rest, calculate the velocity it would have when it reaches the ground in the absence of drag ($F_D = 0$).
- (c) Assuming that in the presence of drag the hailstone reaches the ground with terminal velocity, calculate the amount of energy that has been lost due to drag.