

Suggested Solution, DFT

(1)

1) Find $X[k]$ given $x(t)$

$$X[k] = \frac{1}{T} \int_{-1}^1 e^{-t} \cdot e^{-jk\Omega_T t} dt$$

$$= \frac{1}{2} \int_{-1}^1 e^{-t} \cdot e^{-jk\pi t} dt$$

$$= \frac{1}{2} \int_{-1}^1 e^{-(1+jk\pi)t} dt$$

$$= \frac{-1}{2(1+jk\pi)} \cdot \left[e^{-(1+jk\pi)t} \right]_{-1}^1$$

$$= \frac{e^{(1+jk\pi)} - e^{-(1+jk\pi)}}{2(1+jk\pi)} \quad k \in]-\infty; \infty[$$

It is possible though to "polish" this expression a little bit;

$$\begin{aligned}
 X[k] &= \frac{e^{(1+jk\pi)} - e^{-(1+jk\pi)}}{2(1+jk\pi)} & (2) \\
 &= \frac{e \cdot e^{jk\pi} - e^{-1} \cdot e^{-jk\pi}}{2(1+jk\pi)} \\
 &= \frac{e(\cos k\pi + j\sin k\pi) - \frac{1}{e}(\cos k\pi - j\sin k\pi)}{2(1+jk\pi)} \\
 &= \frac{(e - \frac{1}{e})\cos(k\pi)}{2(1+jk\pi)} \\
 &= \frac{(1 - jk\pi)(e - \frac{1}{e})\cos(k\pi)}{2(1 + k^2\pi^2)}
 \end{aligned}$$

Plot $|X[k]|$ for $k \in [-30; 30]$


$$|X[k]| = \left| \frac{(e - \frac{1}{e})\cos(k\pi)}{2(1 + k^2\pi^2)} \right| \cdot \sqrt{1 + (k\pi)^2}$$

$k \in [-30; 30]$

The plot is easily made in
term of a small Matlab program.

(3)

Amplitude of the complex FS coefficients.



If you look at $x(t)$ over one period from $t=-1$ to $t=1$ you will realize that the signal actually is very slowly varying. This is consistent with the plot above which shows a significant DC component and very little "high" frequencies.

Prob. 4 p. 713.

(4)

$$x[n] = a^n u[n] \quad \alpha < 1$$

A periodic sequence is constructed from $x[n]$;

$$\tilde{x}[n] = \sum_{r=-\infty}^{\infty} x[n+rN]$$

a) Fourier Transform $X(e^{j\omega})$ of $x[n]$

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] \cdot e^{-j\omega n}$$

$$= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} = \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n$$

$$= \frac{1}{1 - \alpha e^{-j\omega}} \quad |\alpha| < 1$$

Geometric Series.

(5)

b) The DFS of $\tilde{x}[n]$

$$\tilde{X}[k] = \sum_{n=0}^{N-1} \tilde{x}[n] \cdot W_N^{kn}$$

periodic sequence
which is now inserted.

Twiddle factor.

$$\tilde{X}[k] = \sum_{n=0}^{N-1} \sum_{r=-\infty}^{\infty} x[n+rN] W_N^{kn}$$

$$= \sum_{n=0}^{N-1} \sum_{r=-\infty}^{\infty} \alpha^{n+rN} u[n+rN] W_N^{kn}$$

$$= \sum_{n=0}^{N-1} \sum_{r=0}^{\infty} \alpha^{n+rN} W_N^{kn}$$

Rearranging the summations;

$$\tilde{X}[k] = \sum_{r=0}^{\infty} \alpha^{rN} \sum_{n=0}^{N-1} \alpha^n W_N^{kn}$$

↓

6

Again, let's utilize that there are geometric series;

$$W_N^{kn} = e^{-j \frac{2\pi}{N} kn}$$

$$\Downarrow \quad \alpha^n W_N^{kn} = \alpha^n e^{-j \frac{2\pi}{N} kn} = \left(\alpha \cdot e^{-j \frac{2\pi}{N} k} \right)^n$$

$$\sum_{n=0}^{N-1} \beta^n = \frac{1 - \beta^{(N-1)+1}}{1 - \beta} = \beta^N$$

$$= \frac{1 - \alpha^N e^{-j 2\pi k}}{1 - \alpha \cdot e^{-j \frac{2\pi}{N} k}}$$

\Downarrow

$$\tilde{X}[k] = \sum_{r=0}^{\infty} \underbrace{\alpha^{r \cdot N}}_{(\alpha^N)^r} \cdot \frac{1 - \alpha^N e^{-j 2\pi k}}{1 - \alpha e^{-j \frac{2\pi}{N} k}} \quad |\alpha| < 1$$

\Downarrow

As related to r , this can now be considered a constant.

$$(\alpha^N)^r$$

$$\tilde{X}[k] = \frac{1}{1-\alpha^N} \cdot \frac{1-\alpha^N \cdot \underbrace{e^{-j2\pi k}}_{=1}}{1-\alpha \cdot e^{-j\frac{2\pi}{N} \cdot k}} \quad |\alpha| < 1 \quad (7)$$

$$\Downarrow \quad \tilde{X}[k] = \frac{1}{1-\alpha \cdot e^{-j\frac{2\pi}{N} \cdot k}} \quad |\alpha| < 1$$

c) Compare a) and b)

$$\begin{cases} X(e^{j\omega}) = \frac{1}{1-\alpha e^{-j\omega}} & |\alpha| < 1 \\ \tilde{X}[k] = \frac{1}{1-\alpha e^{-j\frac{2\pi}{N} \cdot k}} & |\alpha| < 1 \end{cases}$$

$$\Downarrow \quad \tilde{X}[k] = X(e^{j\omega}) \Big|_{\omega = \frac{2\pi k}{N}}$$

So, the Fourier Series coefficients are the Fourier transform (freq. response) sampled in frequencies $\omega_k = \frac{2\pi}{N} \cdot k$

⑧

Prob. 5 p. 713.

Compute DFT of finite length sequences considered to be of length N (even).

a) $x[n] = \delta[n]$

$$\begin{aligned}
 X[k] &= \sum_{n=0}^{N-1} \delta[n] \cdot W_N^{kn} & 0 \leq k \leq N-1 \\
 &= \sum_{n=0}^{N-1} \delta[n] e^{-j \frac{2\pi}{N} kn} = \underline{\underline{1}}
 \end{aligned}$$


b) $x[n] = \delta[n - n_0] \quad 0 \leq n_0 \leq N-1$

$$\begin{aligned}
 X[k] &= \sum_{n=0}^{N-1} \delta[n - n_0] \cdot W_N^{kn} & 0 \leq k \leq (N-1) \\
 &= \sum_{n=0}^{N-1} \delta[n - n_0] e^{-j \frac{2\pi}{N} kn} \\
 &= \underline{\underline{W_N^{kn_0}}}
 \end{aligned}$$

Prob. 7 p. 714.

9

We have a six-point uniform sequence $x[n]$ (figure P7 p. 714) which is non-zero for $0 \leq n \leq 5$



$X(z)$ is the Z-transform of $x[n]$

If we sample $X(z)$ @ $z = e^{j\frac{2\pi}{4}k}$ $k=0,1,2,3$
then we obtain $X_1[k] = X(z) \big|_{z=e^{j\frac{2\pi}{4}k}}$ $k=0,1,2,3$.

Sketch the inverse DFT of $X_1[k]$.

$$\left\{ \begin{array}{l} \text{Now; } X(z) = \sum_{n=-\infty}^{\infty} x[n] \cdot z^{-n} = \sum_{n=0}^5 z^{-n} \end{array} \right.$$

Next we substitute $z = e^{j\frac{2\pi}{4}k}$



(10)

$$X[k] = \sum_{n=0}^5 e^{-j \frac{2\pi k}{4} \cdot n} = \sum_{n=0}^5 e^{-j \frac{2\pi}{4} kn}$$

$$= \sum_{n=0}^5 W_4^{kn}$$

$$0 \leq k \leq 4$$

The period $N=4$.

This is a 4-point DFT.
 The original sequence was of length 6

Going back from $X[k]$ to the time domain
 we may therefore expect some aliasing

Let's now do inverse DFT

$$X[k] = W_4^{0k} + W_4^{1k} + W_4^{2k} + W_4^{3k} + W_4^{4k} + W_4^{5k} \quad 0 \leq k \leq 4.$$

Based on the result we found in prob. 5a,
 we can now do inverse DFT by inspection.

Six impulses (one for each value of n)

However;

$$W_4^{4k} = e^{-j \frac{2\pi}{4} \cdot 4k} = e^{-j \frac{2\pi}{4} \cdot 0 \cdot k} = W_4^{0k}$$

and

$$W_4^{5k} = e^{-j \frac{2\pi}{4} \cdot 5k} = e^{-j \frac{2\pi}{4} \cdot k} = W_4^k$$

Therefore two points are aliased;

