

Dossier zur Nutzenbewertung gemäß § 35a SGB V

Zolbetuximab (VYLOY™)

Astellas Pharma GmbH

Modul 4 A, Anhang 4-G4

*Erstlinienbehandlung von erwachsenen Patienten mit
lokal fortgeschrittenem inoperablem oder metastasiertem
HER2-negativem Adenokarzinom des Magens oder des GEJ,
deren Tumore Claudin18.2 positiv sind*

Studie SPOTLIGHT

Primärer Datenschnitt vom 09.09.2022

Finaler Datenschnitt vom 08.09.2023

Stand: 30.10.2024

Inhaltsverzeichnis

Anhang 4-G4 Post hoc Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022 ..5	
Anhang 4-G4 Baseline Charakteristika	5
Anhang 4-G4 Gesamtüberleben und Progressionsfreies Überleben	43
Anhang 4-G4 Gesamtüberleben.....	43
1. Time-to-Event-Analyse	43
2. Kaplan-Meier-Plots	48
Anhang 4-G4 Progressionsfreies Überleben (IRC und INV)	50
1. Time-to-Event-Analysen	50
2. Kaplan-Meier-Plots	59
Anhang 4-G4 Patientenberichtete Endpunkte.....	70
Anhang 4-G4 Symptomatik und Gesundheitsbezogene Lebensqualität anhand des EORTC	
QLQ-C30	70
1. Rücklaufquoten	70
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	119
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	198
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	229
5. Kaplan-Meier-Plots	251
Anhang 4-G4 Symptomatik anhand des EORTC QLQ-OG25.....	273
1. Rücklaufquoten	273
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	325
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	431
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	464
5. Kaplan-Meier-Plots	495
Anhang 4-G4 Schmerzintensität anhand der NRS	526
1. Rücklaufquoten	526
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	530
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	535
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	538
5. Kaplan-Meier-Plots	540
Anhang 4-G4 EQ-5D VAS	542
1. Rücklaufquoten	542
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	546
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	551
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	554
5. Kaplan-Meier-Plots	556
Anhang 4-G4 Sicherheit.....	558
Anhang 4-G4 Unerwünschte Ereignisse.....	558
1. Time-to Event-Analysen	558
2. Kaplan-Meier-Plots	561
Anhang 4-G4 Schwere unerwünschte Ereignisse.....	563
1. Time-to-Event-Analysen	563
2. Kaplan-Meier-Plots	566
Anhang 4-G4 Schwerwiegende unerwünschte Ereignisse.....	568

1. Time-to-Event-Analysen	568
2. Kaplan-Meier-Plots	571
Anhang 4-G4 Abbruch der Studienmedikation aufgrund unerwünschter Ereignisse	573
1. Time-to-Event-Analysen	573
2. Kaplan-Meier-Plots	576
Anhang 4-G4 Unerwünschte Ereignisse - SOC und PT	578
1. Time-to-Event-Analysen	578
2. Kaplan-Meier-Plots	733
Anhang 4-G4 Schwere unerwünschte Ereignisse - SOC und PT	876
1. Time-to-Event-Analysen	876
2. Kaplan-Meier-Plots	921
Anhang 4-G4 Schwerwiegende unerwünschte Ereignisse - SOC und PT	962
1. Time-to-Event-Analysen	962
2. Kaplan-Meier-Plots	986
Anhang 4-G4 Abbruchgründe – SOC und PT	1010
Anhang 4-G4 Unerwünschte Ereignisse von besonderem Interesse	1017
1. Time-to-Event-Analysen	1017
2. Kaplan-Meier-Plots	1102
Anhang 4-G4 Weitere Analysen	1153
Anhang 4-G4 Beobachtungsdauern	1153
Anhang 4-G4 Begleitmedikationen	1163
Anhang 4-G4 Studienmedikation - Exposition	1170
Anhang 4-G4 Patientenfluss	1173

Anhang 4-G4 <i>Post hoc</i> Analysen für die Studie SPOTLIGHT, finaler Datenschnitt 08.09.2023.....	1176
Anhang 4-G4 Baseline Charakteristika	1176
Anhang 4-G4 Gesamtüberleben und Progressionsfreies Überleben	1214
Anhang 4-G4 Gesamtüberleben.....	1214
1. Time-to-Event-Analyse	1214
2. Kaplan-Meier-Plots	1219
Anhang 4-G4 Progressionsfreies Überleben (IRC und INV)	1223
1. Time-to-Event-Analysen	1223
2. Kaplan-Meier-Plots	1232
Anhang 4-G4 Patientenberichtete Endpunkte.....	1237
Anhang 4-G4 Symptomatik und Gesundheitsbezogene Lebensqualität anhand des EORTC	
QLQ-C30	1237
1. Rücklaufquoten	1237
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	1302
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	1384
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	1415
5. Kaplan-Meier-Plots	1437
Anhang 4-G4 Symptomatik anhand des EORTC QLQ-OG25.....	1459
1. Rücklaufquoten	1459
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	1528
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	1641
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	1674
5. Kaplan-Meier-Plots	1705
Anhang 4-G4 Schmerzintensität anhand der NRS	1736
1. Rücklaufquoten	1736
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	1741
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	1746
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	1749
5. Kaplan-Meier-Plots	1751
Anhang 4-G4 EQ-5D VAS	1753
1. Rücklaufquoten	1753
2. Verlauf über die Zeit (Observed Means and Change from Baseline)	1758
3. Time-to-Event-Analysen - Zeit bis zur ersten Verschlechterung.....	1763
4. Graphische Darstellung des Verlaufs (Mean Change from Baseline)	1766
5. Kaplan-Meier-Plots	1768
Anhang 4-G4 Sicherheit.....	1770
Anhang 4-G4 Unerwünschte Ereignisse.....	1770
1. Time-to Event-Analysen	1770
2. Kaplan-Meier-Plots	1773
Anhang 4-G4 Schwere unerwünschte Ereignisse	1775
1. Time-to-Event-Analysen	1775
2. Kaplan-Meier-Plots	1778
Anhang 4-G4 Schwerwiegende unerwünschte Ereignisse	1780
1. Time-to-Event-Analysen	1780
2. Kaplan-Meier-Plots	1783

Anhang 4-G4 Abbruch der Studienmedikation aufgrund unerwünschter Ereignisse	1785
1. Time-to-Event-Analysen	1785
2. Kaplan-Meier-Plots	1788
Anhang 4-G4 Unerwünschte Ereignisse - SOC und PT	1790
1. Time-to-Event-Analysen	1790
2. Kaplan-Meier-Plots	1957
Anhang 4-G4 Schwere unerwünschte Ereignisse - SOC und PT	2110
1. Time-to-Event-Analysen	2110
2. Kaplan-Meier-Plots	2157
Anhang 4-G4 Schwerwiegende unerwünschte Ereignisse - SOC und PT	2199
1. Time-to-Event-Analysen	2199
2. Kaplan-Meier-Plots	2223
Anhang 4-G4 Abbruchgründe – SOC und PT	2247
Anhang 4-G4 Unerwünschte Ereignisse von besonderem Interesse.....	2254
1. Time-to-Event-Analysen	2254
2. Kaplan-Meier-Plots	2339
Anhang 4-G4 Weitere Analysen.....	2396
Anhang 4-G4 Beobachtungsdauern.....	2396
Anhang 4-G4 Begleitmedikationen.....	2406
Anhang 4-G4 Studienmedikation - Exposition	2413
Anhang 4-G4 Patientenfluss.....	2416
Anhang 4-G4 Folgetherapien	2419

Anhang 4-G4 *Post hoc* Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022

Anhang 4-G4 Baseline Charakteristika

Table 301.1.1001.1: Summary of Baseline Demographics - Full Analysis Set

Parameter and Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Sex		
Male	176 (62.2%)	175 (62.1%)
Female	107 (37.8%)	107 (37.9%)
Unknown	0	0
Age (years)		
n	283	282
Mean (SD)	59.74 (11.75)	58.77 (12.96)
Median	62.0	60.0
Range	27.0 - 83.0	20.0 - 86.0
Age (years)		
>=18 to <=64	171 (60.4%)	174 (61.7%)
>=65 to <85	112 (39.6%)	106 (37.6%)
>=85	0	2 (0.7%)
Age Group 1 (years)		
<=65	181 (64.0%)	181 (64.2%)
>65	102 (36.0%)	101 (35.8%)
Age Group 2 (years)		
<=75	267 (94.3%)	260 (92.2%)
>75	16 (5.7%)	22 (7.8%)
Race		
White	140 (53.6%)	134 (53.0%)
Black or African American	5 (1.9%)	2 (0.8%)

Abbreviations: BMI=body mass index; BSA=body surface area; CPS=combined positive score; ECOG=eastern cooperative oncology group; eCRF=electronic case report form; IRT=interactive response technology; N=number of patients; n=number of patients with non-missing values; SD=standard deviation.

ASTELLAS Data Cutoff Date: 09SEP22

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Table 301.1.1001.1: Summary of Baseline Demographics - Full Analysis Set

Parameter and Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Asian	96 (36.8%)	97 (38.3%)
American Indian or Alaska Native	9 (3.4%)	8 (3.2%)
Native Hawaiian or Other Pacific Islander	0	0
Other	11 (4.2%)	12 (4.7%)
Missing	22	29
Ethnicity		
Hispanic or Latino	36 (13.8%)	37 (14.8%)
Not Hispanic or Latino	225 (86.2%)	213 (85.2%)
Missing	22	32
Country 1		
Japan	32 (11.3%)	33 (11.7%)
Non-Japan	251 (88.7%)	249 (88.3%)
Country 2		
China	19 (6.7%)	17 (6.0%)
Non-China	264 (93.3%)	265 (94.0%)
Height (cm)		
n	279	277
Mean (SD)	167.17 (9.25)	166.87 (10.33)
Median	168.0	167.5
Range	145.0 - 188.0	143.0 - 196.0
Weight (kg)		
n	279	278

Abbreviations: BMI=body mass index; BSA=body surface area; CPS=combined positive score; ECOG=eastern cooperative oncology group; eCRF=electronic case report form; IRT=interactive response technology; N=number of patients; n=number of patients with non-missing values; SD=standard deviation.
ASTELLAS Data Cutoff Date: 09SEP22

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Table 301.1.1001.1: Summary of Baseline Demographics - Full Analysis Set

Parameter and Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Mean (SD)	64.66 (14.47)	65.43 (16.39)
Median	63.0	64.8
Range	38.0 - 110.6	28.5 - 128.3
BMI (kg/m^2)		
<18.5	34 (12.2%)	27 (9.7%)
>=18.5 to <25	168 (60.2%)	169 (61.0%)
>=25 to <30	61 (21.9%)	61 (22.0%)
>=30	16 (5.7%)	20 (7.2%)
Missing	4	5
BSA (m^2)		
n	279	277
Mean (SD)	1.73 (0.23)	1.74 (0.25)
Median	1.7	1.7
Range	1.2 - 2.4	1.1 - 2.5
BSA (m^2)		
<1.7	128 (45.9%)	127 (45.8%)
>=1.7	151 (54.1%)	150 (54.2%)
Missing	4	5
Tobacco History		
Never	142 (50.5%)	137 (48.9%)
Current	26 (9.3%)	25 (8.9%)
Former	113 (40.2%)	118 (42.1%)

Abbreviations: BMI=body mass index; BSA=body surface area; CPS=combined positive score; ECOG=eastern cooperative oncology group; eCRF=electronic case report form; IRT=interactive response technology; N=number of patients; n=number of patients with non-missing values; SD=standard deviation.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.1: Summary of Baseline Demographics - Full Analysis Set

Parameter and Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Missing	2	2
Baseline ECOG Status		
0	125 (44.8%)	115 (41.4%)
>=1	154 (55.2%)	163 (58.6%)
Missing	4	4
Region (IRT)		
Asia	88 (31.1%)	89 (31.6%)
Non-Asia	195 (68.9%)	193 (68.4%)
Region (eCRF)		
Asia	88 (31.1%)	89 (31.6%)
Non-Asia	195 (68.9%)	193 (68.4%)
Number of Organs with Metastatic Sites (IRT)		
0-2	219 (77.4%)	219 (77.7%)
>=3	64 (22.6%)	63 (22.3%)
Number of Organs with Metastatic Sites (eCRF)		
0-2	220 (77.7%)	222 (78.7%)
>=3	63 (22.3%)	60 (21.3%)
Prior Gastrectomy (IRT)		
Yes	84 (29.7%)	82 (29.1%)
No	199 (70.3%)	200 (70.9%)
Prior Gastrectomy (eCRF)		
Yes	82 (29.0%)	83 (29.4%)

Abbreviations: BMI=body mass index; BSA=body surface area; CPS=combined positive score; ECOG=eastern cooperative oncology group; eCRF=electronic case report form; IRT=interactive response technology; N=number of patients; n=number of patients with non-missing values; SD=standard deviation.
 ASTELLAS Data Cutoff Date: 09SEP22

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Table 301.1.1001.1: Summary of Baseline Demographics - Full Analysis Set

Parameter and Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
No	201 (71.0%)	199 (70.6%)
CPS Status		
<5	170 (60.1%)	173 (61.3%)
>=5	27 (9.5%)	24 (8.5%)
Unknown	86 (30.4%)	85 (30.1%)

Abbreviations: BMI=body mass index; BSA=body surface area; CPS=combined positive score; ECOG=eastern cooperative oncology group; eCRF=electronic case report form; IRT=interactive response technology; N=number of patients; n=number of patients with non-missing values; SD=standard deviation.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.3: Summary of Primary Diagnosis - Full Analysis Set

Parameter	Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Medical Condition	Gastric Adenocarcinoma	219 (77.4%)	210 (74.5%)
	Gastro-Esophageal Junction Adenocarcinoma	64 (22.6%)	72 (25.5%)
Duration Since Initial Diagnosis (days)	n	273	275
	Mean (SD)	270.36 (478.45)	297.63 (675.21)
	Median	56.0	56.0
	Range	2 - 3010	7 - 5366
Tumor Location			
All	Proximal	103 (36.8%)	85 (30.2%)
	Distal	110 (39.3%)	118 (42.0%)
	Unknown	67 (23.9%)	78 (27.8%)
	Missing	3	1
Gastric	n	219	210
	Proximal	73 (33.6%)	59 (28.1%)
	Distal	91 (41.9%)	87 (41.4%)
	Unknown	53 (24.4%)	64 (30.5%)
GEJ	Missing	2	0
	n	64	72
	Proximal	30 (47.6%)	26 (36.6%)
	Distal	19 (30.2%)	31 (43.7%)
Tumor Type	Unknown	14 (22.2%)	14 (19.7%)
	Missing	1	1
	Diffuse	82 (29.1%)	117 (42.1%)
	Intestinal	70 (24.8%)	66 (23.7%)

Abbreviations: GEJ=gastro-esophageal junction; N=number of patients n=number of patients with non-missing values; SD=standard deviation.

[1] Distant metastases (M1) and Tumor Metastatic (Yes) may differ depending on date of initial diagnosis.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.3: Summary of Primary Diagnosis - Full Analysis Set

Parameter	Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Primary Tumor	Mixed	31 (11.0%)	13 (4.7%)
	Other	50 (17.7%)	42 (15.1%)
	Unknown	49 (17.4%)	40 (14.4%)
	Missing	1	4
	TX	62 (22.1%)	46 (16.4%)
	T0	0	0
	Tis	1 (0.4%)	2 (0.7%)
	T1	2 (0.7%)	4 (1.4%)
	T1a	2 (0.7%)	1 (0.4%)
	T1b	2 (0.7%)	6 (2.1%)
	T2	15 (5.3%)	16 (5.7%)
	T3	86 (30.6%)	98 (34.9%)
	T4	32 (11.4%)	35 (12.5%)
	T4a	56 (19.9%)	56 (19.9%)
	T4b	23 (8.2%)	17 (6.0%)
	Missing	2	1
Regional Lymph Nodes	NX	66 (23.7%)	60 (21.4%)
	N0	40 (14.3%)	38 (13.6%)
	N1	56 (20.1%)	66 (23.6%)
	N2	44 (15.8%)	51 (18.2%)
	N3	42 (15.1%)	32 (11.4%)
	N3a	17 (6.1%)	19 (6.8%)
	N3b	14 (5.0%)	14 (5.0%)
	Missing	4	2
Distant Metastasis [1]	M0	85 (30.4%)	70 (24.8%)
	M1	195 (69.6%)	212 (75.2%)

Abbreviations: GEJ=gastro-esophageal junction; N=number of patients n=number of patients with non-missing values; SD=standard deviation.

[1] Distant metastases (M1) and Tumor Metastatic (Yes) may differ depending on date of initial diagnosis.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.3: Summary of Primary Diagnosis - Full Analysis Set

Parameter	Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
	Missing	3	0
Tumor Metastatic [1]	Yes	239 (84.5%)	238 (84.4%)
	No	44 (15.5%)	44 (15.6%)
Metastasis Location	Abdominal Cavity	19 (6.7%)	17 (6.0%)
	Adrenal Gland	7 (2.5%)	6 (2.1%)
	Bladder	1 (0.4%)	0
	Bone	28 (9.9%)	23 (8.2%)
	Brain	0	1 (0.4%)
	Breast	1 (0.4%)	0
	Chest	4 (1.4%)	1 (0.4%)
	Colon	2 (0.7%)	3 (1.1%)
	Esophagus	3 (1.1%)	4 (1.4%)
	Gallbladder	0	1 (0.4%)
	Heart	2 (0.7%)	1 (0.4%)
	Kidney	2 (0.7%)	0
	Liver	62 (21.9%)	75 (26.6%)
	Lung	36 (12.7%)	33 (11.7%)
	Lymph Node	101 (35.7%)	109 (38.7%)
	Mediastinum	5 (1.8%)	2 (0.7%)
	Neck	0	1 (0.4%)
	Omentum	10 (3.5%)	12 (4.3%)
	Other	23 (8.1%)	17 (6.0%)
	Ovary	16 (5.7%)	19 (6.7%)
	Pancreas	2 (0.7%)	4 (1.4%)
	Pelvis	2 (0.7%)	3 (1.1%)
	Pericardium	1 (0.4%)	0

Abbreviations: GEJ=gastro-esophageal junction; N=number of patients n=number of patients with non-missing values; SD=standard deviation.

[1] Distant metastases (M1) and Tumor Metastatic (Yes) may differ depending on date of initial diagnosis.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.3: Summary of Primary Diagnosis - Full Analysis Set

Parameter	Category/Statistic	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
	Peritoneum	94 (33.2%)	76 (27.0%)
	Pleura	4 (1.4%)	5 (1.8%)
	Rectum	3 (1.1%)	0
	Retroperitoneum	7 (2.5%)	9 (3.2%)
	Skin	1 (0.4%)	0
	Spleen	2 (0.7%)	3 (1.1%)
	Stomach	5 (1.8%)	5 (1.8%)
History of Helicobacter pylori Infection	Yes	31 (11.0%)	45 (16.0%)
	No	139 (49.1%)	136 (48.2%)
	Unknown	113 (39.9%)	101 (35.8%)
Barrett's Esophagus Diagnosed	Yes	7 (2.5%)	11 (3.9%)
	No	166 (58.7%)	173 (61.3%)
	Unknown	110 (38.9%)	98 (34.8%)
CLDN18.2 Testing Result	<75%	0	0
	>=75%	283 (100.0%)	282 (100.0%)
	Not Applicable	0	0
HER2 Status	Positive	0	0
	Negative	283 (100.0%)	282 (100.0%)
	Not Applicable	0	0
Measurable Disease based on Central	Yes	211 (74.6%)	211 (74.8%)
	No	72 (25.4%)	71 (25.2%)
Measurable Disease based on Local	Yes	235 (83.0%)	227 (80.5%)
	No	48 (17.0%)	55 (19.5%)

Abbreviations: GEJ=gastro-esophageal junction; N=number of patients n=number of patients with non-missing values; SD=standard deviation.

[1] Distant metastases (M1) and Tumor Metastatic (Yes) may differ depending on date of initial diagnosis.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Overall	259 (91.5%)	251 (89.0%)	510 (90.3%)
Blood And Lymphatic System Disorders	50 (17.7%)	49 (17.4%)	99 (17.5%)
Anaemia	44 (15.5%)	44 (15.6%)	88 (15.6%)
Iron Deficiency Anaemia	4 (1.4%)	5 (1.8%)	9 (1.6%)
Blood Loss Anaemia	1 (0.4%)	0	1 (0.2%)
Hypochromic Anaemia	1 (0.4%)	0	1 (0.2%)
Lymphadenopathy Mediastinal	1 (0.4%)	0	1 (0.2%)
Microcytic Anaemia	0	1 (0.4%)	1 (0.2%)
Pernicious Anaemia	1 (0.4%)	0	1 (0.2%)
Splenic Vein Thrombosis	1 (0.4%)	0	1 (0.2%)
Splenomegaly	1 (0.4%)	0	1 (0.2%)
Cardiac Disorders	32 (11.3%)	30 (10.6%)	62 (11.0%)
Atrial Fibrillation	6 (2.1%)	7 (2.5%)	13 (2.3%)
Angina Pectoris	1 (0.4%)	6 (2.1%)	7 (1.2%)
Coronary Artery Disease	1 (0.4%)	6 (2.1%)	7 (1.2%)
Bradycardia	3 (1.1%)	1 (0.4%)	4 (0.7%)
Myocardial Ischaemia	3 (1.1%)	1 (0.4%)	4 (0.7%)
Palpitations	2 (0.7%)	2 (0.7%)	4 (0.7%)
Arrhythmia	2 (0.7%)	1 (0.4%)	3 (0.5%)
Myocardial Infarction	2 (0.7%)	1 (0.4%)	3 (0.5%)
Sinus Bradycardia	3 (1.1%)	0	3 (0.5%)
Sinus Tachycardia	1 (0.4%)	2 (0.7%)	3 (0.5%)
Tachycardia	0	3 (1.1%)	3 (0.5%)
Atrial Flutter	2 (0.7%)	0	2 (0.4%)
Cardiovascular Disorder	2 (0.7%)	0	2 (0.4%)
Ischaemic Cardiomyopathy	2 (0.7%)	0	2 (0.4%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Mitral Valve Incompetence	1 (0.4%)	1 (0.4%)	2 (0.4%)
Pericardial Effusion	1 (0.4%)	1 (0.4%)	2 (0.4%)
Sinus Arrhythmia	1 (0.4%)	1 (0.4%)	2 (0.4%)
Acute Myocardial Infarction	1 (0.4%)	0	1 (0.2%)
Aortic Valve Incompetence	0	1 (0.4%)	1 (0.2%)
Arteriosclerosis Coronary Artery	0	1 (0.4%)	1 (0.2%)
Cardiac Arrest	0	1 (0.4%)	1 (0.2%)
Cardiac Flutter	0	1 (0.4%)	1 (0.2%)
Cardiac Tamponade	1 (0.4%)	0	1 (0.2%)
Cardiomegaly	1 (0.4%)	0	1 (0.2%)
Coronary Artery Stenosis	1 (0.4%)	0	1 (0.2%)
Diastolic Dysfunction	1 (0.4%)	0	1 (0.2%)
Hypertensive Heart Disease	1 (0.4%)	0	1 (0.2%)
Pericarditis	1 (0.4%)	0	1 (0.2%)
Supraventricular Extrasystoles	1 (0.4%)	0	1 (0.2%)
Ventricular Extrasystoles	0	1 (0.4%)	1 (0.2%)
Congenital, Familial And Genetic Disorders	3 (1.1%)	10 (3.5%)	13 (2.3%)
Gilbert's Syndrome	1 (0.4%)	2 (0.7%)	3 (0.5%)
Brugada Syndrome	0	1 (0.4%)	1 (0.2%)
Congenital Musculoskeletal Disorder Of Limbs	1 (0.4%)	0	1 (0.2%)
Deafness Congenital	0	1 (0.4%)	1 (0.2%)
Dermoid Cyst	0	1 (0.4%)	1 (0.2%)
Gene Mutation	0	1 (0.4%)	1 (0.2%)
Porokeratosis	0	1 (0.4%)	1 (0.2%)
Sturge-Weber Syndrome	1 (0.4%)	0	1 (0.2%)
Thalassaemia Minor	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Type V Hyperlipidaemia	0	1 (0.4%)	1 (0.2%)
Ventricular Septal Defect	0	1 (0.4%)	1 (0.2%)
Ear And Labyrinth Disorders	5 (1.8%)	10 (3.5%)	15 (2.7%)
Deafness	0	4 (1.4%)	4 (0.7%)
Hypoacusis	2 (0.7%)	1 (0.4%)	3 (0.5%)
Vertigo	1 (0.4%)	2 (0.7%)	3 (0.5%)
Meniere's Disease	2 (0.7%)	0	2 (0.4%)
Tinnitus	0	2 (0.7%)	2 (0.4%)
Deafness Neurosensory	0	1 (0.4%)	1 (0.2%)
Mastoid Disorder	0	1 (0.4%)	1 (0.2%)
Presbycusis	0	1 (0.4%)	1 (0.2%)
Endocrine Disorders	24 (8.5%)	19 (6.7%)	43 (7.6%)
Hypothyroidism	18 (6.4%)	16 (5.7%)	34 (6.0%)
Hyperthyroidism	2 (0.7%)	0	2 (0.4%)
Thyroid Disorder	1 (0.4%)	1 (0.4%)	2 (0.4%)
Autoimmune Thyroiditis	1 (0.4%)	0	1 (0.2%)
Goitre	1 (0.4%)	0	1 (0.2%)
Hyperparathyroidism Primary	0	1 (0.4%)	1 (0.2%)
Hyperplasia Adrenal	0	1 (0.4%)	1 (0.2%)
Thyroiditis	1 (0.4%)	0	1 (0.2%)
Eye Disorders	11 (3.9%)	17 (6.0%)	28 (5.0%)
Glaucoma	5 (1.8%)	3 (1.1%)	8 (1.4%)
Cataract	4 (1.4%)	3 (1.1%)	7 (1.2%)
Dry Eye	0	3 (1.1%)	3 (0.5%)
Diabetic Retinopathy	0	2 (0.7%)	2 (0.4%)
Macular Degeneration	0	2 (0.7%)	2 (0.4%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Retinal Degeneration	0	2 (0.7%)	2 (0.4%)
Retinal Detachment	0	2 (0.7%)	2 (0.4%)
Vision Blurred	0	2 (0.7%)	2 (0.4%)
Visual Impairment	1 (0.4%)	1 (0.4%)	2 (0.4%)
Blepharitis	0	1 (0.4%)	1 (0.2%)
Chalazion	1 (0.4%)	0	1 (0.2%)
Chorioretinal Scar	1 (0.4%)	0	1 (0.2%)
Epiretinal Membrane	0	1 (0.4%)	1 (0.2%)
Lacrimation Decreased	0	1 (0.4%)	1 (0.2%)
Macular Oedema	0	1 (0.4%)	1 (0.2%)
Optic Neuropathy	0	1 (0.4%)	1 (0.2%)
Visual Field Defect	0	1 (0.4%)	1 (0.2%)
Vitreous Detachment	1 (0.4%)	0	1 (0.2%)
Gastrointestinal Disorders	146 (51.6%)	157 (55.7%)	303 (53.6%)
Abdominal Pain	33 (11.7%)	46 (16.3%)	79 (14.0%)
Gastrooesophageal Reflux Disease	35 (12.4%)	44 (15.6%)	79 (14.0%)
Constipation	31 (11.0%)	39 (13.8%)	70 (12.4%)
Dysphagia	34 (12.0%)	31 (11.0%)	65 (11.5%)
Nausea	24 (8.5%)	39 (13.8%)	63 (11.2%)
Abdominal Pain Upper	21 (7.4%)	21 (7.4%)	42 (7.4%)
Vomiting	10 (3.5%)	16 (5.7%)	26 (4.6%)
Dyspepsia	10 (3.5%)	12 (4.3%)	22 (3.9%)
Diarrhoea	7 (2.5%)	12 (4.3%)	19 (3.4%)
Abdominal Distension	6 (2.1%)	11 (3.9%)	17 (3.0%)
Ascites	7 (2.5%)	9 (3.2%)	16 (2.8%)
Gastric Ulcer	4 (1.4%)	12 (4.3%)	16 (2.8%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Gastritis	7 (2.5%)	9 (3.2%)	16 (2.8%)
Abdominal Discomfort	7 (2.5%)	6 (2.1%)	13 (2.3%)
Haemorrhoids	6 (2.1%)	6 (2.1%)	12 (2.1%)
Inguinal Hernia	5 (1.8%)	4 (1.4%)	9 (1.6%)
Chronic Gastritis	4 (1.4%)	4 (1.4%)	8 (1.4%)
Hiatus Hernia	4 (1.4%)	3 (1.1%)	7 (1.2%)
Umbilical Hernia	4 (1.4%)	3 (1.1%)	7 (1.2%)
Diverticulum	1 (0.4%)	5 (1.8%)	6 (1.1%)
Flatulence	1 (0.4%)	4 (1.4%)	5 (0.9%)
Abdominal Pain Lower	2 (0.7%)	2 (0.7%)	4 (0.7%)
Eructation	0	4 (1.4%)	4 (0.7%)
Intestinal Obstruction	1 (0.4%)	3 (1.1%)	4 (0.7%)
Melaena	1 (0.4%)	3 (1.1%)	4 (0.7%)
Peptic Ulcer	2 (0.7%)	2 (0.7%)	4 (0.7%)
Diverticulum Intestinal	1 (0.4%)	2 (0.7%)	3 (0.5%)
Dumping Syndrome	0	3 (1.1%)	3 (0.5%)
Irritable Bowel Syndrome	0	3 (1.1%)	3 (0.5%)
Large Intestine Polyp	3 (1.1%)	0	3 (0.5%)
Barrett's Oesophagus	2 (0.7%)	0	2 (0.4%)
Colitis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Duodenal Ulcer	2 (0.7%)	0	2 (0.4%)
Haematochezia	0	2 (0.7%)	2 (0.4%)
Odynophagia	0	2 (0.7%)	2 (0.4%)
Oesophagitis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Upper Gastrointestinal Haemorrhage	1 (0.4%)	1 (0.4%)	2 (0.4%)
Abdominal Hernia	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Anal Fistula	1 (0.4%)	0	1 (0.2%)
Anal Haemorrhage	0	1 (0.4%)	1 (0.2%)
Aphthous Ulcer	0	1 (0.4%)	1 (0.2%)
Diaphragmatic Hernia	0	1 (0.4%)	1 (0.2%)
Discoloured Vomit	0	1 (0.4%)	1 (0.2%)
Dry Mouth	0	1 (0.4%)	1 (0.2%)
Duodenitis	1 (0.4%)	0	1 (0.2%)
Epigastric Discomfort	1 (0.4%)	0	1 (0.2%)
Gastritis Erosive	0	1 (0.4%)	1 (0.2%)
Gastritis Hypertrophic	1 (0.4%)	0	1 (0.2%)
Gastrointestinal Haemorrhage	0	1 (0.4%)	1 (0.2%)
Hyperchlorhydria	1 (0.4%)	0	1 (0.2%)
Impaired Gastric Emptying	0	1 (0.4%)	1 (0.2%)
Intestinal Cyst	1 (0.4%)	0	1 (0.2%)
Obstruction Gastric	0	1 (0.4%)	1 (0.2%)
Pancreatic Failure	1 (0.4%)	0	1 (0.2%)
Pancreatitis Acute	1 (0.4%)	0	1 (0.2%)
Proctalgia	1 (0.4%)	0	1 (0.2%)
Proctitis Ulcerative	1 (0.4%)	0	1 (0.2%)
Pylorospasm	0	1 (0.4%)	1 (0.2%)
Rectal Polyp	1 (0.4%)	0	1 (0.2%)
Regurgitation	0	1 (0.4%)	1 (0.2%)
Retching	0	1 (0.4%)	1 (0.2%)
Subileus	1 (0.4%)	0	1 (0.2%)
Tongue Coated	1 (0.4%)	0	1 (0.2%)
Toothache	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
General Disorders And Administration Site Conditions	46 (16.3%)	55 (19.5%)	101 (17.9%)
Fatigue	23 (8.1%)	32 (11.3%)	55 (9.7%)
Asthenia	6 (2.1%)	8 (2.8%)	14 (2.5%)
Early Satiety	4 (1.4%)	5 (1.8%)	9 (1.6%)
Pain	5 (1.8%)	3 (1.1%)	8 (1.4%)
Chest Pain	1 (0.4%)	5 (1.8%)	6 (1.1%)
Non-Cardiac Chest Pain	3 (1.1%)	3 (1.1%)	6 (1.1%)
Oedema Peripheral	4 (1.4%)	2 (0.7%)	6 (1.1%)
Pyrexia	3 (1.1%)	2 (0.7%)	5 (0.9%)
Hernia	1 (0.4%)	2 (0.7%)	3 (0.5%)
Malaise	1 (0.4%)	2 (0.7%)	3 (0.5%)
Catheter Site Pain	0	2 (0.7%)	2 (0.4%)
Medical Device Pain	2 (0.7%)	0	2 (0.4%)
Chills	0	1 (0.4%)	1 (0.2%)
Complication Associated With Device	1 (0.4%)	0	1 (0.2%)
Granuloma	1 (0.4%)	0	1 (0.2%)
Impaired Healing	0	1 (0.4%)	1 (0.2%)
Localised Oedema	1 (0.4%)	0	1 (0.2%)
Peripheral Swelling	0	1 (0.4%)	1 (0.2%)
Swelling	1 (0.4%)	0	1 (0.2%)
Temperature Intolerance	0	1 (0.4%)	1 (0.2%)
Hepatobiliary Disorders	17 (6.0%)	9 (3.2%)	26 (4.6%)
Cholelithiasis	4 (1.4%)	2 (0.7%)	6 (1.1%)
Cholecystitis	2 (0.7%)	1 (0.4%)	3 (0.5%)
Hepatic Function Abnormal	2 (0.7%)	1 (0.4%)	3 (0.5%)
Hepatic Steatosis	2 (0.7%)	1 (0.4%)	3 (0.5%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Hepatic Cyst	1 (0.4%)	1 (0.4%)	2 (0.4%)
Bile Duct Stone	1 (0.4%)	0	1 (0.2%)
Biliary Dilatation	1 (0.4%)	0	1 (0.2%)
Biliary Obstruction	1 (0.4%)	0	1 (0.2%)
Gallbladder Polyp	1 (0.4%)	0	1 (0.2%)
Hepatic Cirrhosis	0	1 (0.4%)	1 (0.2%)
Hepatitis	1 (0.4%)	0	1 (0.2%)
Hepatotoxicity	1 (0.4%)	0	1 (0.2%)
Hyperbilirubinaemia	0	1 (0.4%)	1 (0.2%)
Jaundice Cholestatic	1 (0.4%)	0	1 (0.2%)
Non-Alcoholic Fatty Liver	0	1 (0.4%)	1 (0.2%)
Immune System Disorders	12 (4.2%)	18 (6.4%)	30 (5.3%)
Drug Hypersensitivity	5 (1.8%)	7 (2.5%)	12 (2.1%)
Seasonal Allergy	5 (1.8%)	7 (2.5%)	12 (2.1%)
Hypersensitivity	1 (0.4%)	1 (0.4%)	2 (0.4%)
Allergy To Animal	0	1 (0.4%)	1 (0.2%)
Allergy To Arthropod Sting	0	1 (0.4%)	1 (0.2%)
Food Allergy	0	1 (0.4%)	1 (0.2%)
Hypogammaglobulinaemia	1 (0.4%)	0	1 (0.2%)
Immunodeficiency Common Variable	1 (0.4%)	0	1 (0.2%)
Mite Allergy	0	1 (0.4%)	1 (0.2%)
Rubber Sensitivity	0	1 (0.4%)	1 (0.2%)
Infections And Infestations	38 (13.4%)	31 (11.0%)	69 (12.2%)
Helicobacter Infection	4 (1.4%)	4 (1.4%)	8 (1.4%)
Upper Respiratory Tract Infection	1 (0.4%)	4 (1.4%)	5 (0.9%)
Urinary Tract Infection	3 (1.1%)	2 (0.7%)	5 (0.9%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Helicobacter Gastritis	4 (1.4%)	0	4 (0.7%)
Pneumonia	2 (0.7%)	2 (0.7%)	4 (0.7%)
Bronchitis	2 (0.7%)	1 (0.4%)	3 (0.5%)
Hepatitis A	2 (0.7%)	1 (0.4%)	3 (0.5%)
Hepatitis B	3 (1.1%)	0	3 (0.5%)
Pulmonary Tuberculosis	3 (1.1%)	0	3 (0.5%)
Chronic Sinusitis	0	2 (0.7%)	2 (0.4%)
Covid-19	1 (0.4%)	1 (0.4%)	2 (0.4%)
Meningitis Viral	1 (0.4%)	1 (0.4%)	2 (0.4%)
Rhinitis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Tuberculosis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Abdominal Abscess	1 (0.4%)	0	1 (0.2%)
Appendicitis	1 (0.4%)	0	1 (0.2%)
Appendicitis Perforated	0	1 (0.4%)	1 (0.2%)
Cellulitis	0	1 (0.4%)	1 (0.2%)
Citrobacter Sepsis	1 (0.4%)	0	1 (0.2%)
Conjunctivitis	1 (0.4%)	0	1 (0.2%)
Covid-19 Pneumonia	0	1 (0.4%)	1 (0.2%)
Cryptococcosis	0	1 (0.4%)	1 (0.2%)
Device Related Infection	0	1 (0.4%)	1 (0.2%)
Encephalitis Viral	0	1 (0.4%)	1 (0.2%)
Endocarditis	0	1 (0.4%)	1 (0.2%)
Enterobacter Infection	1 (0.4%)	0	1 (0.2%)
Epiglottitis	0	1 (0.4%)	1 (0.2%)
Furuncle	0	1 (0.4%)	1 (0.2%)
Gastrointestinal Candidiasis	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Herpes Simplex	0	1 (0.4%)	1 (0.2%)
Infected Dermal Cyst	0	1 (0.4%)	1 (0.2%)
Infective Aneurysm	0	1 (0.4%)	1 (0.2%)
Labyrinthitis	1 (0.4%)	0	1 (0.2%)
Nasopharyngitis	0	1 (0.4%)	1 (0.2%)
Onychomycosis	0	1 (0.4%)	1 (0.2%)
Oral Candidiasis	0	1 (0.4%)	1 (0.2%)
Oral Herpes	0	1 (0.4%)	1 (0.2%)
Papilloma Viral Infection	1 (0.4%)	0	1 (0.2%)
Parotitis	1 (0.4%)	0	1 (0.2%)
Periodontitis	1 (0.4%)	0	1 (0.2%)
Peritonitis	0	1 (0.4%)	1 (0.2%)
Peritonsillar Abscess	0	1 (0.4%)	1 (0.2%)
Pilonidal Disease	1 (0.4%)	0	1 (0.2%)
Pyelonephritis Acute	1 (0.4%)	0	1 (0.2%)
Sinusitis	0	1 (0.4%)	1 (0.2%)
Tonsillitis	0	1 (0.4%)	1 (0.2%)
Viral Hepatitis Carrier	0	1 (0.4%)	1 (0.2%)
Viral Myocarditis	0	1 (0.4%)	1 (0.2%)
Injury, Poisoning And Procedural Complications	13 (4.6%)	18 (6.4%)	31 (5.5%)
Procedural Pain	3 (1.1%)	2 (0.7%)	5 (0.9%)
Gun Shot Wound	0	3 (1.1%)	3 (0.5%)
Stoma Site Pain	1 (0.4%)	2 (0.7%)	3 (0.5%)
Upper Limb Fracture	2 (0.7%)	1 (0.4%)	3 (0.5%)
Ankle Fracture	1 (0.4%)	1 (0.4%)	2 (0.4%)
Fibula Fracture	2 (0.7%)	0	2 (0.4%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Scar	2 (0.7%)	0	2 (0.4%)
Anastomotic Complication	0	1 (0.4%)	1 (0.2%)
Aortic Injury	0	1 (0.4%)	1 (0.2%)
Bone Contusion	1 (0.4%)	0	1 (0.2%)
Contusion	0	1 (0.4%)	1 (0.2%)
Exposure To Chemical Pollution	0	1 (0.4%)	1 (0.2%)
Exposure To Radiation	0	1 (0.4%)	1 (0.2%)
Femur Fracture	1 (0.4%)	0	1 (0.2%)
Foot Fracture	0	1 (0.4%)	1 (0.2%)
Head Injury	0	1 (0.4%)	1 (0.2%)
Infusion Related Reaction	0	1 (0.4%)	1 (0.2%)
Joint Dislocation	0	1 (0.4%)	1 (0.2%)
Meniscus Injury	1 (0.4%)	0	1 (0.2%)
Reactive Gastropathy	1 (0.4%)	0	1 (0.2%)
Rib Fracture	0	1 (0.4%)	1 (0.2%)
Skin Abrasion	0	1 (0.4%)	1 (0.2%)
Spinal Compression Fracture	1 (0.4%)	0	1 (0.2%)
Tibia Fracture	1 (0.4%)	0	1 (0.2%)
Wound Complication	0	1 (0.4%)	1 (0.2%)
Investigations	25 (8.8%)	39 (13.8%)	64 (11.3%)
Weight Decreased	12 (4.2%)	20 (7.1%)	32 (5.7%)
Alanine Aminotransferase Increased	2 (0.7%)	4 (1.4%)	6 (1.1%)
Blood Cholesterol Increased	3 (1.1%)	3 (1.1%)	6 (1.1%)
Aspartate Aminotransferase Increased	2 (0.7%)	3 (1.1%)	5 (0.9%)
Blood Alkaline Phosphatase Increased	2 (0.7%)	3 (1.1%)	5 (0.9%)
Arthroscopy	1 (0.4%)	1 (0.4%)	2 (0.4%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Blood Creatinine Increased	0	2 (0.7%)	2 (0.4%)
Cardiac Murmur	0	2 (0.7%)	2 (0.4%)
Lymphocyte Count Decreased	1 (0.4%)	1 (0.4%)	2 (0.4%)
Oesophagogastrroduodenoscopy	1 (0.4%)	1 (0.4%)	2 (0.4%)
Biopsy Fallopian Tube	0	1 (0.4%)	1 (0.2%)
Blood Bilirubin Increased	1 (0.4%)	0	1 (0.2%)
Blood Creatinine Decreased	0	1 (0.4%)	1 (0.2%)
Blood Testosterone Decreased	0	1 (0.4%)	1 (0.2%)
Creatinine Renal Clearance Decreased	0	1 (0.4%)	1 (0.2%)
Electrocardiogram Abnormal	0	1 (0.4%)	1 (0.2%)
Electrocardiogram Q Wave Abnormal	0	1 (0.4%)	1 (0.2%)
Electrocardiogram Qt Prolonged	1 (0.4%)	0	1 (0.2%)
Endoscopic Retrograde Cholangiopancreatography	1 (0.4%)	0	1 (0.2%)
Gamma-Glutamyltransferase Increased	0	1 (0.4%)	1 (0.2%)
Glomerular Filtration Rate Decreased	1 (0.4%)	0	1 (0.2%)
Haemoglobin Decreased	1 (0.4%)	0	1 (0.2%)
Helicobacter Test Positive	1 (0.4%)	0	1 (0.2%)
Neutrophil Count Increased	0	1 (0.4%)	1 (0.2%)
Occult Blood	0	1 (0.4%)	1 (0.2%)
Occult Blood Positive	1 (0.4%)	0	1 (0.2%)
Oesophagogastrosocopy	0	1 (0.4%)	1 (0.2%)
Platelet Count Decreased	1 (0.4%)	0	1 (0.2%)
Platelet Count Increased	0	1 (0.4%)	1 (0.2%)
Prostatic Specific Antigen Increased	0	1 (0.4%)	1 (0.2%)
Protein Urine Present	0	1 (0.4%)	1 (0.2%)
Stress Echocardiogram Abnormal	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
White Blood Cell Count Decreased	1 (0.4%)	0	1 (0.2%)
White Blood Cell Count Increased	0	1 (0.4%)	1 (0.2%)
Metabolism And Nutrition Disorders	103 (36.4%)	107 (37.9%)	210 (37.2%)
Decreased Appetite	32 (11.3%)	35 (12.4%)	67 (11.9%)
Diabetes Mellitus	22 (7.8%)	23 (8.2%)	45 (8.0%)
Hyperlipidaemia	16 (5.7%)	19 (6.7%)	35 (6.2%)
Dyslipidaemia	13 (4.6%)	16 (5.7%)	29 (5.1%)
Type 2 Diabetes Mellitus	13 (4.6%)	14 (5.0%)	27 (4.8%)
Hypercholesterolaemia	10 (3.5%)	11 (3.9%)	21 (3.7%)
Gout	4 (1.4%)	5 (1.8%)	9 (1.6%)
Hyperuricaemia	4 (1.4%)	5 (1.8%)	9 (1.6%)
Hypoalbuminaemia	7 (2.5%)	2 (0.7%)	9 (1.6%)
Glucose Tolerance Impaired	2 (0.7%)	4 (1.4%)	6 (1.1%)
Vitamin D Deficiency	3 (1.1%)	3 (1.1%)	6 (1.1%)
Hyperglycaemia	2 (0.7%)	2 (0.7%)	4 (0.7%)
Hypokalaemia	3 (1.1%)	1 (0.4%)	4 (0.7%)
Obesity	0	4 (1.4%)	4 (0.7%)
Dehydration	1 (0.4%)	2 (0.7%)	3 (0.5%)
Lactose Intolerance	2 (0.7%)	1 (0.4%)	3 (0.5%)
Vitamin B12 Deficiency	0	3 (1.1%)	3 (0.5%)
Abnormal Loss Of Weight	0	2 (0.7%)	2 (0.4%)
Hypoglycaemia	0	2 (0.7%)	2 (0.4%)
Hypomagnesaemia	1 (0.4%)	1 (0.4%)	2 (0.4%)
Hyponatraemia	1 (0.4%)	1 (0.4%)	2 (0.4%)
Impaired Fasting Glucose	1 (0.4%)	1 (0.4%)	2 (0.4%)
Cachexia	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Hypercalcaemia	0	1 (0.4%)	1 (0.2%)
Hyperkalaemia	1 (0.4%)	0	1 (0.2%)
Hypertriglyceridaemia	1 (0.4%)	0	1 (0.2%)
Hypocalcaemia	1 (0.4%)	0	1 (0.2%)
Hypoproteinaemia	1 (0.4%)	0	1 (0.2%)
Hypotriglyceridaemia	0	1 (0.4%)	1 (0.2%)
Iron Deficiency	0	1 (0.4%)	1 (0.2%)
Malnutrition	0	1 (0.4%)	1 (0.2%)
Type 1 Diabetes Mellitus	1 (0.4%)	0	1 (0.2%)
Musculoskeletal And Connective Tissue Disorders	58 (20.5%)	60 (21.3%)	118 (20.9%)
Back Pain	19 (6.7%)	21 (7.4%)	40 (7.1%)
Arthralgia	9 (3.2%)	5 (1.8%)	14 (2.5%)
Osteoarthritis	5 (1.8%)	6 (2.1%)	11 (1.9%)
Intervertebral Disc Protrusion	5 (1.8%)	3 (1.1%)	8 (1.4%)
Osteoporosis	3 (1.1%)	5 (1.8%)	8 (1.4%)
Spinal Osteoarthritis	3 (1.1%)	4 (1.4%)	7 (1.2%)
Spinal Pain	4 (1.4%)	2 (0.7%)	6 (1.1%)
Arthritis	1 (0.4%)	4 (1.4%)	5 (0.9%)
Neck Pain	1 (0.4%)	4 (1.4%)	5 (0.9%)
Musculoskeletal Pain	2 (0.7%)	2 (0.7%)	4 (0.7%)
Spinal Stenosis	3 (1.1%)	1 (0.4%)	4 (0.7%)
Fibromyalgia	0	3 (1.1%)	3 (0.5%)
Lumbar Spinal Stenosis	1 (0.4%)	2 (0.7%)	3 (0.5%)
Pain In Extremity	2 (0.7%)	1 (0.4%)	3 (0.5%)
Bone Pain	2 (0.7%)	0	2 (0.4%)
Flank Pain	1 (0.4%)	1 (0.4%)	2 (0.4%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Joint Swelling	2 (0.7%)	0	2 (0.4%)
Muscle Spasms	1 (0.4%)	1 (0.4%)	2 (0.4%)
Musculoskeletal Chest Pain	1 (0.4%)	1 (0.4%)	2 (0.4%)
Myalgia	0	2 (0.7%)	2 (0.4%)
Rheumatoid Arthritis	2 (0.7%)	0	2 (0.4%)
Tendonitis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Ankylosing Spondylitis	1 (0.4%)	0	1 (0.2%)
Bursitis	0	1 (0.4%)	1 (0.2%)
Cervical Spinal Stenosis	0	1 (0.4%)	1 (0.2%)
Chondrocalcinosis	0	1 (0.4%)	1 (0.2%)
Diffuse Idiopathic Skeletal Hyperostosis	0	1 (0.4%)	1 (0.2%)
Dupuytren's Contracture	1 (0.4%)	0	1 (0.2%)
Exostosis	0	1 (0.4%)	1 (0.2%)
Gouty Arthritis	0	1 (0.4%)	1 (0.2%)
Greater Trochanteric Pain Syndrome	1 (0.4%)	0	1 (0.2%)
Intervertebral Disc Degeneration	0	1 (0.4%)	1 (0.2%)
Mobility Decreased	1 (0.4%)	0	1 (0.2%)
Muscular Weakness	0	1 (0.4%)	1 (0.2%)
Musculoskeletal Discomfort	0	1 (0.4%)	1 (0.2%)
Periarthritis	0	1 (0.4%)	1 (0.2%)
Polymyalgia Rheumatica	1 (0.4%)	0	1 (0.2%)
Rotator Cuff Syndrome	1 (0.4%)	0	1 (0.2%)
Sjogren's Syndrome	1 (0.4%)	0	1 (0.2%)
Spondylolisthesis	1 (0.4%)	0	1 (0.2%)
Spondylolysis	0	1 (0.4%)	1 (0.2%)
Temporomandibular Joint Syndrome	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Thoracic Spinal Stenosis	0	1 (0.4%)	1 (0.2%)
Trigger Finger	0	1 (0.4%)	1 (0.2%)
Neoplasms Benign, Malignant And Unspecified (Incl Cysts And Polyps)	33 (11.7%)	37 (13.1%)	70 (12.4%)
Tumour Pain	4 (1.4%)	4 (1.4%)	8 (1.4%)
Prostate Cancer	2 (0.7%)	5 (1.8%)	7 (1.2%)
Basal Cell Carcinoma	3 (1.1%)	2 (0.7%)	5 (0.9%)
Breast Cancer	3 (1.1%)	2 (0.7%)	5 (0.9%)
Cancer Pain	2 (0.7%)	2 (0.7%)	4 (0.7%)
Colorectal Adenoma	1 (0.4%)	3 (1.1%)	4 (0.7%)
Colon Cancer	1 (0.4%)	2 (0.7%)	3 (0.5%)
Malignant Ascites	1 (0.4%)	2 (0.7%)	3 (0.5%)
Uterine Leiomyoma	1 (0.4%)	2 (0.7%)	3 (0.5%)
Acrochordon	1 (0.4%)	1 (0.4%)	2 (0.4%)
Benign Ovarian Tumour	0	2 (0.7%)	2 (0.4%)
Bladder Transitional Cell Carcinoma	1 (0.4%)	1 (0.4%)	2 (0.4%)
Gastric Cancer	2 (0.7%)	0	2 (0.4%)
Squamous Cell Carcinoma Of Skin	1 (0.4%)	1 (0.4%)	2 (0.4%)
Adenocarcinoma Gastric	0	1 (0.4%)	1 (0.2%)
Benign Neoplasm Of Bladder	1 (0.4%)	0	1 (0.2%)
Benign Neoplasm Of Optic Nerve	1 (0.4%)	0	1 (0.2%)
Benign Neoplasm Of Thyroid Gland	0	1 (0.4%)	1 (0.2%)
Benign Pancreatic Neoplasm	0	1 (0.4%)	1 (0.2%)
Benign Salivary Gland Neoplasm	0	1 (0.4%)	1 (0.2%)
Bowen's Disease	0	1 (0.4%)	1 (0.2%)
Carcinoid Tumour Pulmonary	0	1 (0.4%)	1 (0.2%)
Cardiac Valve Fibroelastoma	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Cervix Carcinoma	1 (0.4%)	0	1 (0.2%)
Fibroadenoma Of Breast	0	1 (0.4%)	1 (0.2%)
Fibroma	1 (0.4%)	0	1 (0.2%)
Haemangioma	1 (0.4%)	0	1 (0.2%)
Haemangioma Of Liver	1 (0.4%)	0	1 (0.2%)
Haemangioma Of Skin	0	1 (0.4%)	1 (0.2%)
Hodgkin's Disease	1 (0.4%)	0	1 (0.2%)
Invasive Ductal Breast Carcinoma	1 (0.4%)	0	1 (0.2%)
Laryngeal Cancer	0	1 (0.4%)	1 (0.2%)
Leiomyoma	0	1 (0.4%)	1 (0.2%)
Malignant Melanoma	1 (0.4%)	0	1 (0.2%)
Melanocytic Naevus	0	1 (0.4%)	1 (0.2%)
Neuroendocrine Tumour	1 (0.4%)	0	1 (0.2%)
Neuroma	1 (0.4%)	0	1 (0.2%)
Non-Hodgkin's Lymphoma	0	1 (0.4%)	1 (0.2%)
Osteochondroma	0	1 (0.4%)	1 (0.2%)
Papillary Thyroid Cancer	1 (0.4%)	0	1 (0.2%)
Pericardial Effusion Malignant	1 (0.4%)	0	1 (0.2%)
Pituitary Tumour	1 (0.4%)	0	1 (0.2%)
Pyogenic Granuloma	0	1 (0.4%)	1 (0.2%)
Rectal Adenocarcinoma	1 (0.4%)	0	1 (0.2%)
Rectal Cancer	0	1 (0.4%)	1 (0.2%)
Renal Cancer	0	1 (0.4%)	1 (0.2%)
Seborrhoeic Keratosis	0	1 (0.4%)	1 (0.2%)
Squamous Cell Carcinoma	0	1 (0.4%)	1 (0.2%)
Testis Cancer	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Nervous System Disorders	36 (12.7%)	38 (13.5%)	74 (13.1%)
Peripheral Sensory Neuropathy	5 (1.8%)	5 (1.8%)	10 (1.8%)
Epilepsy	5 (1.8%)	3 (1.1%)	8 (1.4%)
Headache	4 (1.4%)	4 (1.4%)	8 (1.4%)
Neuropathy Peripheral	3 (1.1%)	3 (1.1%)	6 (1.1%)
Carpal Tunnel Syndrome	2 (0.7%)	3 (1.1%)	5 (0.9%)
Migraine	3 (1.1%)	2 (0.7%)	5 (0.9%)
Transient Ischaemic Attack	2 (0.7%)	2 (0.7%)	4 (0.7%)
Diabetic Neuropathy	2 (0.7%)	1 (0.4%)	3 (0.5%)
Dizziness	2 (0.7%)	1 (0.4%)	3 (0.5%)
Paraesthesia	1 (0.4%)	2 (0.7%)	3 (0.5%)
Sciatica	1 (0.4%)	2 (0.7%)	3 (0.5%)
Carotid Arteriosclerosis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Cerebral Infarction	1 (0.4%)	1 (0.4%)	2 (0.4%)
Cerebrovascular Accident	1 (0.4%)	1 (0.4%)	2 (0.4%)
Dysgeusia	1 (0.4%)	1 (0.4%)	2 (0.4%)
Memory Impairment	0	2 (0.7%)	2 (0.4%)
Nerve Compression	0	2 (0.7%)	2 (0.4%)
Neuralgia	0	2 (0.7%)	2 (0.4%)
Polyneuropathy	1 (0.4%)	1 (0.4%)	2 (0.4%)
Tremor	1 (0.4%)	1 (0.4%)	2 (0.4%)
Anosmia	0	1 (0.4%)	1 (0.2%)
Ataxia	1 (0.4%)	0	1 (0.2%)
Cerebellar Infarction	0	1 (0.4%)	1 (0.2%)
Cerebral Small Vessel Ischaemic Disease	0	1 (0.4%)	1 (0.2%)
Cerebral Venous Sinus Thrombosis	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Dystonia	0	1 (0.4%)	1 (0.2%)
Hypoaesthesia	0	1 (0.4%)	1 (0.2%)
Ischaemic Stroke	1 (0.4%)	0	1 (0.2%)
Myelopathy	0	1 (0.4%)	1 (0.2%)
Neurotoxicity	1 (0.4%)	0	1 (0.2%)
Occipital Neuralgia	0	1 (0.4%)	1 (0.2%)
Parkinson's Disease	0	1 (0.4%)	1 (0.2%)
Periodic Limb Movement Disorder	0	1 (0.4%)	1 (0.2%)
Peripheral Motor Neuropathy	1 (0.4%)	0	1 (0.2%)
Peripheral Sensorimotor Neuropathy	0	1 (0.4%)	1 (0.2%)
Peroneal Nerve Palsy	0	1 (0.4%)	1 (0.2%)
Sensory Disturbance	1 (0.4%)	0	1 (0.2%)
Vocal Cord Paresis	0	1 (0.4%)	1 (0.2%)
Pregnancy, Puerperium And Perinatal Conditions	2 (0.7%)	2 (0.7%)	4 (0.7%)
Abortion Spontaneous	1 (0.4%)	0	1 (0.2%)
Abortion Spontaneous Incomplete	0	1 (0.4%)	1 (0.2%)
Cervical Incompetence	1 (0.4%)	0	1 (0.2%)
Delivery	1 (0.4%)	0	1 (0.2%)
Ectopic Pregnancy	0	1 (0.4%)	1 (0.2%)
Gestational Hypertension	1 (0.4%)	0	1 (0.2%)
Postpartum Haemorrhage	0	1 (0.4%)	1 (0.2%)
Product Issues	0	1 (0.4%)	1 (0.2%)
Thrombosis In Device	0	1 (0.4%)	1 (0.2%)
Psychiatric Disorders	51 (18.0%)	59 (20.9%)	110 (19.5%)
Anxiety	16 (5.7%)	30 (10.6%)	46 (8.1%)
Insomnia	17 (6.0%)	28 (9.9%)	45 (8.0%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Depression	16 (5.7%)	11 (3.9%)	27 (4.8%)
Alcoholism	2 (0.7%)	1 (0.4%)	3 (0.5%)
Tobacco Abuse	2 (0.7%)	1 (0.4%)	3 (0.5%)
Mixed Anxiety And Depressive Disorder	1 (0.4%)	1 (0.4%)	2 (0.4%)
Nicotine Dependence	2 (0.7%)	0	2 (0.4%)
Adjustment Disorder	1 (0.4%)	0	1 (0.2%)
Affective Disorder	1 (0.4%)	0	1 (0.2%)
Alcohol Abuse	1 (0.4%)	0	1 (0.2%)
Alcohol Withdrawal Syndrome	1 (0.4%)	0	1 (0.2%)
Anticipatory Anxiety	0	1 (0.4%)	1 (0.2%)
Anxiety Disorder	0	1 (0.4%)	1 (0.2%)
Bipolar Disorder	0	1 (0.4%)	1 (0.2%)
Bruxism	0	1 (0.4%)	1 (0.2%)
Depressed Mood	1 (0.4%)	0	1 (0.2%)
Major Depression	0	1 (0.4%)	1 (0.2%)
Panic Attack	0	1 (0.4%)	1 (0.2%)
Post-Traumatic Stress Disorder	1 (0.4%)	0	1 (0.2%)
Stress	1 (0.4%)	0	1 (0.2%)
Renal And Urinary Disorders	27 (9.5%)	28 (9.9%)	55 (9.7%)
Hydronephrosis	3 (1.1%)	7 (2.5%)	10 (1.8%)
Nephrolithiasis	5 (1.8%)	4 (1.4%)	9 (1.6%)
Dysuria	3 (1.1%)	4 (1.4%)	7 (1.2%)
Renal Cyst	3 (1.1%)	4 (1.4%)	7 (1.2%)
Acute Kidney Injury	1 (0.4%)	2 (0.7%)	3 (0.5%)
Chronic Kidney Disease	0	3 (1.1%)	3 (0.5%)
Pollakiuria	1 (0.4%)	2 (0.7%)	3 (0.5%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Proteinuria	2 (0.7%)	1 (0.4%)	3 (0.5%)
Ureterolithiasis	3 (1.1%)	0	3 (0.5%)
Calculus Urinary	1 (0.4%)	1 (0.4%)	2 (0.4%)
Haematuria	1 (0.4%)	1 (0.4%)	2 (0.4%)
Hypertonic Bladder	2 (0.7%)	0	2 (0.4%)
Nocturia	0	2 (0.7%)	2 (0.4%)
Urinary Retention	2 (0.7%)	0	2 (0.4%)
Urinary Tract Obstruction	0	2 (0.7%)	2 (0.4%)
Bladder Obstruction	1 (0.4%)	0	1 (0.2%)
Bladder Spasm	0	1 (0.4%)	1 (0.2%)
Calculus Bladder	0	1 (0.4%)	1 (0.2%)
Cystitis Glandularis	1 (0.4%)	0	1 (0.2%)
Renal Impairment	1 (0.4%)	0	1 (0.2%)
Renal Pain	0	1 (0.4%)	1 (0.2%)
Stress Urinary Incontinence	0	1 (0.4%)	1 (0.2%)
Ureteric Obstruction	1 (0.4%)	0	1 (0.2%)
Urge Incontinence	0	1 (0.4%)	1 (0.2%)
Reproductive System And Breast Disorders	25 (8.8%)	34 (12.1%)	59 (10.4%)
Benign Prostatic Hyperplasia	16 (5.7%)	16 (5.7%)	32 (5.7%)
Erectile Dysfunction	3 (1.1%)	5 (1.8%)	8 (1.4%)
Pelvic Pain	1 (0.4%)	3 (1.1%)	4 (0.7%)
Ovarian Cyst	2 (0.7%)	1 (0.4%)	3 (0.5%)
Prostatomegaly	0	3 (1.1%)	3 (0.5%)
Adenomyosis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Prostatic Calcification	0	2 (0.7%)	2 (0.4%)
Breast Cyst	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Cervical Dysplasia	1 (0.4%)	0	1 (0.2%)
Endometrial Hyperplasia	1 (0.4%)	0	1 (0.2%)
Female Genital Tract Fistula	0	1 (0.4%)	1 (0.2%)
Menstruation Irregular	0	1 (0.4%)	1 (0.2%)
Penile Pain	1 (0.4%)	0	1 (0.2%)
Polycystic Ovaries	1 (0.4%)	0	1 (0.2%)
Sexual Dysfunction	0	1 (0.4%)	1 (0.2%)
Testicular Atrophy	0	1 (0.4%)	1 (0.2%)
Vaginal Haemorrhage	0	1 (0.4%)	1 (0.2%)
Vulvovaginal Dryness	0	1 (0.4%)	1 (0.2%)
Vulvovaginal Pruritus	0	1 (0.4%)	1 (0.2%)
Respiratory, Thoracic And Mediastinal Disorders	47 (16.6%)	49 (17.4%)	96 (17.0%)
Cough	7 (2.5%)	13 (4.6%)	20 (3.5%)
Asthma	9 (3.2%)	8 (2.8%)	17 (3.0%)
Dyspnoea	9 (3.2%)	6 (2.1%)	15 (2.7%)
Chronic Obstructive Pulmonary Disease	6 (2.1%)	7 (2.5%)	13 (2.3%)
Rhinitis Allergic	8 (2.8%)	4 (1.4%)	12 (2.1%)
Pulmonary Embolism	4 (1.4%)	5 (1.8%)	9 (1.6%)
Pleural Effusion	2 (0.7%)	4 (1.4%)	6 (1.1%)
Obstructive Sleep Apnoea Syndrome	2 (0.7%)	2 (0.7%)	4 (0.7%)
Pneumothorax	0	4 (1.4%)	4 (0.7%)
Pulmonary Mass	1 (0.4%)	3 (1.1%)	4 (0.7%)
Sleep Apnoea Syndrome	2 (0.7%)	2 (0.7%)	4 (0.7%)
Bronchiectasis	2 (0.7%)	1 (0.4%)	3 (0.5%)
Emphysema	2 (0.7%)	1 (0.4%)	3 (0.5%)
Hiccups	3 (1.1%)	0	3 (0.5%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Rhinorrhoea	2 (0.7%)	1 (0.4%)	3 (0.5%)
Atelectasis	1 (0.4%)	1 (0.4%)	2 (0.4%)
Dysphonia	1 (0.4%)	1 (0.4%)	2 (0.4%)
Dyspnoea Exertional	0	2 (0.7%)	2 (0.4%)
Nasal Polyps	1 (0.4%)	1 (0.4%)	2 (0.4%)
Sinus Congestion	1 (0.4%)	1 (0.4%)	2 (0.4%)
Upper-Airway Cough Syndrome	2 (0.7%)	0	2 (0.4%)
Allergic Sinusitis	1 (0.4%)	0	1 (0.2%)
Nasal Cavity Mass	1 (0.4%)	0	1 (0.2%)
Nasal Congestion	0	1 (0.4%)	1 (0.2%)
Nasal Septum Deviation	0	1 (0.4%)	1 (0.2%)
Obstructive Airways Disorder	1 (0.4%)	0	1 (0.2%)
Pleuritic Pain	0	1 (0.4%)	1 (0.2%)
Pneumonitis	0	1 (0.4%)	1 (0.2%)
Pulmonary Congestion	0	1 (0.4%)	1 (0.2%)
Pulmonary Hypertension	0	1 (0.4%)	1 (0.2%)
Pulmonary Thrombosis	0	1 (0.4%)	1 (0.2%)
Rhinitis Hypertrophic	0	1 (0.4%)	1 (0.2%)
Wheezing	1 (0.4%)	0	1 (0.2%)
Skin And Subcutaneous Tissue Disorders	14 (4.9%)	24 (8.5%)	38 (6.7%)
Psoriasis	2 (0.7%)	5 (1.8%)	7 (1.2%)
Dry Skin	1 (0.4%)	5 (1.8%)	6 (1.1%)
Eczema	1 (0.4%)	4 (1.4%)	5 (0.9%)
Pruritus	1 (0.4%)	3 (1.1%)	4 (0.7%)
Actinic Keratosis	0	2 (0.7%)	2 (0.4%)
Alopecia	0	2 (0.7%)	2 (0.4%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Dermatitis Atopic	2 (0.7%)	0	2 (0.4%)
Night Sweats	1 (0.4%)	1 (0.4%)	2 (0.4%)
Palmar-Plantar Erythrodysesthesia Syndrome	2 (0.7%)	0	2 (0.4%)
Skin Lesion	0	2 (0.7%)	2 (0.4%)
Alopecia Areata	1 (0.4%)	0	1 (0.2%)
Dermatitis	0	1 (0.4%)	1 (0.2%)
Hand Dermatitis	1 (0.4%)	0	1 (0.2%)
Hyperhidrosis	0	1 (0.4%)	1 (0.2%)
Itching Scar	0	1 (0.4%)	1 (0.2%)
Lipohypertrophy	0	1 (0.4%)	1 (0.2%)
Rash	1 (0.4%)	0	1 (0.2%)
Seborrhoea	0	1 (0.4%)	1 (0.2%)
Skin Mass	1 (0.4%)	0	1 (0.2%)
Skin Ulcer	1 (0.4%)	0	1 (0.2%)
Vitiligo	1 (0.4%)	0	1 (0.2%)
Social Circumstances	8 (2.8%)	8 (2.8%)	16 (2.8%)
Ex-Tobacco User	4 (1.4%)	1 (0.4%)	5 (0.9%)
Tobacco User	1 (0.4%)	3 (1.1%)	4 (0.7%)
Alcohol Use	1 (0.4%)	2 (0.7%)	3 (0.5%)
Menopause	1 (0.4%)	2 (0.7%)	3 (0.5%)
Social Alcohol Drinker	0	3 (1.1%)	3 (0.5%)
Postmenopause	2 (0.7%)	0	2 (0.4%)
Surgical And Medical Procedures	40 (14.1%)	50 (17.7%)	90 (15.9%)
Appendectomy	4 (1.4%)	11 (3.9%)	15 (2.7%)
Cholecystectomy	3 (1.1%)	6 (2.1%)	9 (1.6%)
Cataract Operation	1 (0.4%)	5 (1.8%)	6 (1.1%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Gastrectomy	4 (1.4%)	2 (0.7%)	6 (1.1%)
Hysterectomy	2 (0.7%)	3 (1.1%)	5 (0.9%)
Inguinal Hernia Repair	3 (1.1%)	1 (0.4%)	4 (0.7%)
Polypectomy	1 (0.4%)	3 (1.1%)	4 (0.7%)
Caesarean Section	1 (0.4%)	2 (0.7%)	3 (0.5%)
Haemorrhoid Operation	2 (0.7%)	1 (0.4%)	3 (0.5%)
Hysterosalpingo-Oophorectomy	1 (0.4%)	2 (0.7%)	3 (0.5%)
Intervertebral Disc Operation	1 (0.4%)	2 (0.7%)	3 (0.5%)
Knee Operation	1 (0.4%)	2 (0.7%)	3 (0.5%)
Prostatectomy	2 (0.7%)	1 (0.4%)	3 (0.5%)
Salpingo-Oophorectomy Bilateral	2 (0.7%)	1 (0.4%)	3 (0.5%)
Splenectomy	3 (1.1%)	0	3 (0.5%)
Tonsillectomy	1 (0.4%)	2 (0.7%)	3 (0.5%)
Vasectomy	1 (0.4%)	2 (0.7%)	3 (0.5%)
Cardiac Pacemaker Insertion	1 (0.4%)	1 (0.4%)	2 (0.4%)
Coronary Arterial Stent Insertion	1 (0.4%)	1 (0.4%)	2 (0.4%)
Hip Arthroplasty	2 (0.7%)	0	2 (0.4%)
Jejunostomy	0	2 (0.7%)	2 (0.4%)
Meniscus Operation	1 (0.4%)	1 (0.4%)	2 (0.4%)
Nephrostomy	2 (0.7%)	0	2 (0.4%)
Oesophageal Prosthesis Insertion	2 (0.7%)	0	2 (0.4%)
Oophorectomy	2 (0.7%)	0	2 (0.4%)
Salpingo-Oophorectomy Unilateral	1 (0.4%)	1 (0.4%)	2 (0.4%)
Shoulder Operation	1 (0.4%)	1 (0.4%)	2 (0.4%)
Thyroidectomy	0	2 (0.7%)	2 (0.4%)
Abdominoplasty	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Adenoidectomy	0	1 (0.4%)	1 (0.2%)
Amblyopia Therapy	0	1 (0.4%)	1 (0.2%)
Ankle Operation	0	1 (0.4%)	1 (0.2%)
Apicectomy	1 (0.4%)	0	1 (0.2%)
Bone Graft	0	1 (0.4%)	1 (0.2%)
Brachytherapy To Prostate	0	1 (0.4%)	1 (0.2%)
Brain Operation	0	1 (0.4%)	1 (0.2%)
Breast Conserving Surgery	1 (0.4%)	0	1 (0.2%)
Breast Cyst Excision	0	1 (0.4%)	1 (0.2%)
Cardiac Ablation	1 (0.4%)	0	1 (0.2%)
Central Venous Catheterisation	1 (0.4%)	0	1 (0.2%)
Colectomy	0	1 (0.4%)	1 (0.2%)
Colostomy	0	1 (0.4%)	1 (0.2%)
Coronary Artery Bypass	0	1 (0.4%)	1 (0.2%)
Female Sterilisation	0	1 (0.4%)	1 (0.2%)
Finger Amputation	0	1 (0.4%)	1 (0.2%)
Foot Operation	1 (0.4%)	0	1 (0.2%)
Gallbladder Operation	0	1 (0.4%)	1 (0.2%)
Gastric Bypass	1 (0.4%)	0	1 (0.2%)
Gastric Operation	0	1 (0.4%)	1 (0.2%)
Gastrointestinal Tube Insertion	0	1 (0.4%)	1 (0.2%)
Haematopoietic Stem Cell Mobilisation	1 (0.4%)	0	1 (0.2%)
Hernia Repair	1 (0.4%)	0	1 (0.2%)
Implantable Defibrillator Insertion	0	1 (0.4%)	1 (0.2%)
Internal Fixation Of Fracture	1 (0.4%)	0	1 (0.2%)
Intraocular Lens Implant	0	1 (0.4%)	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Knee Arthroplasty	0	1 (0.4%)	1 (0.2%)
Large Intestinal Polypectomy	0	1 (0.4%)	1 (0.2%)
Lens Extraction	0	1 (0.4%)	1 (0.2%)
Ligament Operation	0	1 (0.4%)	1 (0.2%)
Mammoplasty	1 (0.4%)	0	1 (0.2%)
Mass Excision	0	1 (0.4%)	1 (0.2%)
Mastectomy	0	1 (0.4%)	1 (0.2%)
Mitral Valve Replacement	0	1 (0.4%)	1 (0.2%)
Oesophagoenterostomy	0	1 (0.4%)	1 (0.2%)
Oophorectomy Bilateral	0	1 (0.4%)	1 (0.2%)
Open Reduction Of Fracture	0	1 (0.4%)	1 (0.2%)
Optic Nerve Operation	1 (0.4%)	0	1 (0.2%)
Pituitary Tumour Removal	1 (0.4%)	0	1 (0.2%)
Pleural Decortication	0	1 (0.4%)	1 (0.2%)
Proctectomy	1 (0.4%)	0	1 (0.2%)
Roux Loop Conversion	1 (0.4%)	0	1 (0.2%)
Salpingectomy	0	1 (0.4%)	1 (0.2%)
Skin Neoplasm Excision	1 (0.4%)	0	1 (0.2%)
Spinal Operation	1 (0.4%)	0	1 (0.2%)
Thrombosis Prophylaxis	1 (0.4%)	0	1 (0.2%)
Tooth Extraction	1 (0.4%)	0	1 (0.2%)
Transurethral Bladder Resection	1 (0.4%)	0	1 (0.2%)
Transurethral Prostatectomy	0	1 (0.4%)	1 (0.2%)
Tumour Excision	1 (0.4%)	0	1 (0.2%)
Umbilical Hernia Repair	0	1 (0.4%)	1 (0.2%)
Urethral Stent Insertion	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1001.4: Summary of Medical History by SOC/PT - Full Analysis Set

MedDRA SOC and PT	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Total (N=565)
Wound Closure	0	1 (0.4%)	1 (0.2%)
Vascular Disorders	86 (30.4%)	97 (34.4%)	183 (32.4%)
Hypertension	80 (28.3%)	90 (31.9%)	170 (30.1%)
Deep Vein Thrombosis	2 (0.7%)	2 (0.7%)	4 (0.7%)
Hypotension	1 (0.4%)	3 (1.1%)	4 (0.7%)
Aortic Aneurysm	0	2 (0.7%)	2 (0.4%)
Peripheral Vascular Disorder	0	2 (0.7%)	2 (0.4%)
Peripheral Venous Disease	0	2 (0.7%)	2 (0.4%)
Phlebitis	0	2 (0.7%)	2 (0.4%)
Aortic Stenosis	0	1 (0.4%)	1 (0.2%)
Arteriosclerosis	1 (0.4%)	0	1 (0.2%)
Atheroembolism	0	1 (0.4%)	1 (0.2%)
Embolism Arterial	0	1 (0.4%)	1 (0.2%)
Essential Hypertension	0	1 (0.4%)	1 (0.2%)
Hot Flush	0	1 (0.4%)	1 (0.2%)
Pallor	1 (0.4%)	0	1 (0.2%)
Pelvic Venous Thrombosis	1 (0.4%)	0	1 (0.2%)
Raynaud's Phenomenon	1 (0.4%)	0	1 (0.2%)
Subclavian Artery Stenosis	0	1 (0.4%)	1 (0.2%)
Subclavian Vein Thrombosis	0	1 (0.4%)	1 (0.2%)
Varicose Vein	1 (0.4%)	0	1 (0.2%)
Vena Cava Thrombosis	1 (0.4%)	0	1 (0.2%)
Venous Thrombosis	1 (0.4%)	0	1 (0.2%)

Abbreviations: N=number of patients; PT=preferred term; SOC=system organ class.

Sorting order: alphabetical order by SOC and descending order by the number of subjects of overall group by PT. In case of ties, alphabetical order by PT is applied.

ASTELLAS Data Cutoff Date: 09SEP22

Anhang 4-G4 *Post hoc* Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022

Anhang 4-G4 Gesamtüberleben und Progressionsfreies Überleben

Anhang 4-G4 Gesamtüberleben

1. Time-to-Event-Analyse

Table 301.1.1002.1.1: Summary of Overall Survival - Full Analysis Set

	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6
Number of patients at risk	283 (100.0%)	282 (100.0%)	
Number of patients with events	149 (52.7%)	177 (62.8%)	
Number of patients censored	134 (47.3%)	105 (37.2%)	
Kaplan-Meier estimates of time to event (months)			
Quartiles, 95% CI [a]			
50%	18.2 [16.4, 22.9]	15.5 [13.5, 16.5]	
Cox proportional hazards model			
Stratified HR, 95% CI			0.750 [0.601, 0.936]
Log-rank test			
Two-sided stratified log-rank p-value			0.0107

Abbreviations: CI=confidence interval; HR=hazard ratio; N=number of patients.
Note: The stratification factors are region (Asia vs. Non-Asia), the number of organs with metastatic sites (0 to 2 vs. ≥ 3) and prior gastrectomy (Yes or No).
[a] Based on the Brookmeyer-Crowley Method.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1002.1.2: Type of Events and Censoring of Overall Survival - Full Analysis Set

	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Number of patients at risk	283 (100.0%)	282 (100.0%)
Number of patients with events	149 (52.7%)	177 (62.8%)
Death Before Analysis Cutoff Date	149 (52.7%)	177 (62.8%)
Number of patients censored	134 (47.3%)	105 (37.2%)
Last Known Alive Date Is Before Cutoff Date	115 (40.6%)	85 (30.1%)
Censored At Cutoff Date	19 (6.7%)	20 (7.1%)

Abbreviations: N=number of patients.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1002.1.3: Summary of Overall Survival by Subgroups - Full Analysis Set

Subgroup	Zolbetuximab + mFOLFOX6			Placebo + mFOLFOX6			Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6		
	N	n (%)	KME [95% CI][a]	N	n (%)	KME [95% CI][a]	HR [95% CI][b]	Log-rank p-value[c]	Test for Interaction p-value[d]
Age Group 1									
≤65 years	181	89 (49.2)	20.2 [17.0, 23.8]	181	112 (61.9)	15.6 [13.5, 17.7]	0.741 [0.561, 0.980]	0.0348	0.9663
>65 years	102	60 (58.8)	16.2 [13.2, 19.8]	101	65 (64.4)	15.3 [10.4, 17.1]	0.761 [0.533, 1.086]	0.1308	
Age Group 2									
≤75 years	267	138 (51.7)	18.9 [16.7, 23.1]	260	165 (63.5)	15.5 [13.5, 16.5]	0.713 [0.568, 0.895]	0.0034	0.1660
>75 years	16	11 (68.8)	9.8 [8.5, 25.3]	22	12 (54.5)	15.6 [9.0, 35.1]	1.315 [0.578, 2.996]	0.5197	
Sex									
Male	176	98 (55.7)	17.4 [15.7, 19.8]	175	113 (64.6)	15.6 [12.2, 17.1]	0.760 [0.579, 0.999]	0.0492	0.7817
Female	107	51 (47.7)	23.8 [17.0, 25.3]	107	64 (59.8)	15.3 [11.6, 19.3]	0.726 [0.502, 1.049]	0.0854	
Race									
White	140	81 (57.9)	16.1 [12.5, 18.6]	134	81 (60.4)	15.3 [13.6, 17.1]	0.948 [0.696, 1.291]	0.7349	0.0550
Asian	96	49 (51.0)	23.3 [19.0, 28.6]	97	65 (67.0)	16.5 [13.1, 19.3]	0.572 [0.393, 0.832]	0.0031	
Tobacco History									
Current	26	15 (57.7)	19.4 [14.3, 23.8]	25	15 (60.0)	15.6 [10.4, 27.5]	0.819 [0.398, 1.686]	0.5866	0.6629
Former	113	64 (56.6)	17.8 [15.1, 24.8]	118	74 (62.7)	15.8 [13.7, 17.8]	0.807 [0.576, 1.131]	0.2123	
Never	142	70 (49.3)	19.0 [15.9, 24.1]	137	87 (63.5)	13.5 [10.7, 17.7]	0.678 [0.494, 0.930]	0.0151	
Region									
Asia	88	47 (53.4)	21.5 [17.7, 28.6]	89	59 (66.3)	17.7 [13.5, 19.7]	0.643 [0.437, 0.947]	0.0244	0.5099
Non-Asia	195	102 (52.3)	17.0 [13.5, 21.5]	193	118 (61.1)	13.7 [11.6, 15.8]	0.796 [0.610, 1.039]	0.0930	
Number of Organs with Metastatic Sites									
0-2	219	110 (50.2)	19.7 [17.0, 23.6]	219	129 (58.9)	15.8 [13.7, 19.0]	0.767 [0.594, 0.990]	0.0410	0.5739
≥3	64	39 (60.9)	16.4 [9.8, 19.4]	63	48 (76.2)	12.0 [9.5, 15.6]	0.670 [0.436, 1.030]	0.0664	
Prior Gastrectomy (total or partial)									
Yes	84	40 (47.6)	24.8 [19.0, 30.0]	82	52 (63.4)	15.7 [13.1, 19.7]	0.575 [0.380, 0.869]	0.0080	0.1607
No	199	109 (54.8)	17.0 [13.9, 18.9]	200	125 (62.5)	14.3 [11.9, 16.5]	0.839 [0.648, 1.086]	0.1807	

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Subgroup	Zolbetuximab + mFOLFOX6			Placebo + mFOLFOX6			Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6		Test for Interaction p-value[d]
	N	n (%)	KME [95% CI][a]	N	n (%)	KME [95% CI][a]	HR [95% CI][b]	Log-rank p-value[c]	
Histology (Tumor Type)									
Diffuse	82	46 (56.1)	20.6 [17.5, 24.8]	117	75 (64.1)	15.8 [11.7, 19.0]	0.766 [0.530, 1.108]	0.1565	0.2257
Intestinal	70	38 (54.3)	22.9 [15.5, 28.9]	66	48 (72.7)	13.8 [10.3, 17.7]	0.552 [0.358, 0.851]	0.0063	
Mixed/Other	81	48 (59.3)	16.1 [10.4, 19.8]	55	34 (61.8)	14.7 [13.1, 19.5]	0.992 [0.638, 1.543]	0.9714	
Tumor Location 1									
Gastric	219	111 (50.7)	20.2 [17.0, 24.1]	210	135 (64.3)	13.8 [11.9, 16.5]	0.666 [0.517, 0.858]	0.0015	0.0766
GEJ	64	38 (59.4)	15.8 [11.7, 17.7]	72	42 (58.3)	16.4 [13.7, 19.9]	1.072 [0.690, 1.666]	0.7596	
Tumor Location 2									
Gastric Proximal	73	40 (54.8)	19.7 [15.9, 26.6]	59	37 (62.7)	17.7 [13.6, 19.7]	0.736 [0.467, 1.158]	0.1827	0.7157
Gastric Distal	91	43 (47.3)	21.5 [16.7, 25.3]	87	53 (60.9)	13.7 [10.8, 15.7]	0.660 [0.441, 0.989]	0.0429	
Tumor Location 3									
GEJ Proximal	30	16 (53.3)	17.5 [15.5, 33.7]	26	18 (69.2)	16.5 [7.4, 26.3]	0.629 [0.316, 1.255]	0.1846	0.0841
GEJ Distal	19	12 (63.2)	15.7 [8.7, 19.4]	31	16 (51.6)	17.1 [11.6, NC]	1.505 [0.710, 3.191]	0.2834	

Abbreviations: CI=confidence interval; GEJ=gastro-esophageal junction; HR=hazard ratio; KME=Kaplan-Meier Estimate of time to event (months); N=number of patients; n=number of patients with event; NC=not calculated.
[a] Based on the Brookmeyer-Crowley Method, [b] Calculated from Cox proportional hazards model, [c] Based on 2-sided log-rank test, unadjusted for multiplicity, and [d] Two-sided Type 3 Wald test p-value from Cox proportional hazards model including treatment, subgroup, and treatment x subgroup interaction.
ASTELLAS Data Cutoff Date: 09SEP22


Anhang 4-G4 *Post hoc* Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022

Anhang 4-G4 Gesamtüberleben und Progressionsfreies Überleben

Anhang 4-G4 Gesamtüberleben

2. Kaplan-Meier-Plots

Figure 301.1.1002.1: Kaplan-Meier Plot of Overall Survival - Full Analysis Set



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; N=number of patients.
ASTELLAS Data Cutoff Date: 09SEP22

Anhang 4-G4 *Post hoc* Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022

Anhang 4-G4 Gesamtüberleben und Progressionsfreies Überleben

Anhang 4-G4 Progressionsfreies Überleben (IRC und INV)

1. Time-to-Event-Analysen

Table 301.1.1002.2.1: Summary of Progression-Free Survival (IRC) - Full Analysis Set

	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6
Number of patients at risk	283 (100.0%)	282 (100.0%)	
Number of patients with events	146 (51.6%)	167 (59.2%)	
Number of patients censored	137 (48.4%)	115 (40.8%)	
Kaplan-Meier estimates of time to event (months)			
Quartiles, 95% CI [a]			
50%	10.6 [8.9, 12.5]	8.7 [8.2, 10.3]	
Cox proportional hazards model			
Stratified HR, 95% CI			0.751 [0.598, 0.942]
Log-rank test			
Two-sided stratified log-rank p-value			0.0132

Abbreviations: CI=confidence interval; HR=hazard ratio; IRC=independent review committee; N=number of patients.
Note: The stratification factors are region (Asia vs. Non-Asia), the number of organs with metastatic sites (0 to 2 vs. ≥ 3) and prior gastrectomy (Yes or No).
[a] Based on the Brookmeyer-Crowley Method.
ASTELLAS Data Cutoff Date: 09SEP22

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Table 301.1.1002.2.2: Type of Events and Censoring of Progression-Free Survival (IRC) - Full Analysis Set

	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Number of patients at risk	283 (100.0%)	282 (100.0%)
Number of patients with events	146 (51.6%)	167 (59.2%)
rPD	87 (30.7%)	98 (34.8%)
No rPD, but death recorded on eCRF	59 (20.8%)	69 (24.5%)
Number of patients censored	137 (48.4%)	115 (40.8%)

Abbreviations: eCRF=electronic case report form; IRC=independent review committee; N=number of patients; rPD=radiological progressive disease.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1002.2.3: Summary of Progression-Free Survival (IRC) by Subgroups - Full Analysis Set

Subgroup	Zolbetuximab + mFOLFOX6			Placebo + mFOLFOX6			Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6			Test for Interaction p-value[d]
	N	n (%)	KME [95% CI][a]	N	n (%)	KME [95% CI][a]	HR [95% CI][b]	Log-rank p-value[c]		
Age Group 1										
<=65 years	181	94 (51.9)	11.0 [8.8, 12.6]	181	106 (58.6)	8.9 [7.9, 10.5]	0.771 [0.583, 1.019]	0.0668	0.6914	
>65 years	102	52 (51.0)	10.6 [8.3, 17.4]	101	61 (60.4)	8.6 [7.4, 10.4]	0.714 [0.492, 1.038]	0.0751		
Age Group 2										
<=75 years	267	138 (51.7)	10.6 [8.9, 12.5]	260	154 (59.2)	8.4 [8.0, 10.3]	0.739 [0.586, 0.931]	0.0098	0.7321	
>75 years	16	8 (50.0)	8.3 [4.4, NC]	22	13 (59.1)	10.4 [7.1, 15.8]	0.955 [0.390, 2.342]	0.9067		
Sex										
Male	176	95 (54.0)	10.6 [8.6, 12.5]	175	106 (60.6)	8.9 [8.2, 10.4]	0.776 [0.587, 1.024]	0.0721	0.6583	
Female	107	51 (47.7)	11.0 [8.5, 17.9]	107	61 (57.0)	8.4 [7.2, 10.5]	0.711 [0.488, 1.034]	0.0724		
Race										
White	140	77 (55.0)	8.9 [8.3, 12.4]	134	82 (61.2)	10.2 [8.4, 10.8]	0.930 [0.681, 1.271]	0.6455	0.0351	
Asian	96	47 (49.0)	14.0 [12.3, 18.1]	97	51 (52.6)	8.2 [6.5, 8.7]	0.527 [0.352, 0.788]	0.0016		
Tobacco History										
Current	26	15 (57.7)	10.3 [7.4, 13.2]	25	14 (56.0)	8.7 [4.4, 17.6]	1.001 [0.480, 2.086]	0.9894	0.7503	
Former	113	56 (49.6)	12.3 [8.8, 15.3]	118	70 (59.3)	9.8 [8.0, 10.8]	0.714 [0.501, 1.018]	0.0603		
Never	142	75 (52.8)	10.3 [8.3, 16.7]	137	82 (59.9)	8.2 [7.1, 10.3]	0.737 [0.538, 1.011]	0.0574		
Region										
Asia	88	45 (51.1)	12.6 [10.4, 17.8]	89	47 (52.8)	8.2 [6.5, 9.1]	0.563 [0.372, 0.852]	0.0060	0.1064	
Non-Asia	195	101 (51.8)	9.7 [8.4, 12.2]	193	120 (62.2)	9.4 [8.2, 10.7]	0.848 [0.650, 1.106]	0.2209		
Number of Organs with Metastatic Sites										
0-2	219	107 (48.9)	12.4 [10.3, 15.8]	219	123 (56.2)	9.2 [8.2, 10.6]	0.726 [0.559, 0.943]	0.0157	0.6418	
>=3	64	39 (60.9)	8.1 [6.2, 9.7]	63	44 (69.8)	8.2 [6.3, 9.8]	0.844 [0.548, 1.301]	0.4412		
Prior Gastrectomy (total or partial)										
Yes	84	40 (47.6)	12.4 [10.4, 18.2]	82	50 (61.0)	9.3 [7.2, 11.3]	0.622 [0.410, 0.943]	0.0242	0.2535	
No	199	106 (53.3)	9.3 [8.4, 12.5]	200	117 (58.5)	8.6 [8.2, 10.3]	0.808 [0.620, 1.053]	0.1130		

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Subgroup	Zolbetuximab + mFOLFOX6					Placebo + mFOLFOX6					Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6			Test for Interaction p-value[d]	
	N	n (%)	KME [95% CI][a]			N	n (%)	KME [95% CI][a]			HR [95% CI][b]	Log-rank p-value[c]			
Histology (Tumor Type)															
Diffuse	82	40 (48.8)	12.5	[9.3,	18.2]	117	64 (54.7)	10.3	[8.4,	13.2]	0.756	[0.506,	1.129]	0.1693	0.1756
Intestinal	70	41 (58.6)	10.3	[8.3,	12.5]	66	46 (69.7)	6.6	[6.0,	9.3]	0.582	[0.379,	0.894]	0.0122	
Mixed/Other	81	49 (60.5)	9.8	[8.1,	14.0]	55	35 (63.6)	8.7	[6.8,	10.5]	0.929	[0.601,	1.434]	0.7403	
Tumor Location 1															
Gastric	219	109 (49.8)	12.2	[9.3,	14.0]	210	126 (60.0)	8.4	[7.8,	10.3]	0.688	[0.531,	0.890]	0.0042	0.1478
GEJ	64	37 (57.8)	8.8	[7.4,	15.8]	72	41 (56.9)	8.9	[8.2,	11.3]	1.015	[0.649,	1.586]	0.9498	
Tumor Location 2															
Gastric Proximal	73	40 (54.8)	10.2	[8.2,	18.2]	59	31 (52.5)	8.3	[7.1,	15.0]	0.876	[0.545,	1.409]	0.5830	0.2419
Gastric Distal	91	39 (42.9)	14.0	[10.4,	18.1]	87	50 (57.5)	9.1	[6.8,	10.5]	0.581	[0.381,	0.885]	0.0105	
Tumor Location 3															
GEJ Proximal	30	14 (46.7)	15.8	[8.3,	20.6]	26	20 (76.9)	7.4	[4.1,	8.7]	0.387	[0.189,	0.794]	0.0076	0.0145
GEJ Distal	19	12 (63.2)	8.8	[6.4,	12.5]	31	16 (51.6)	10.8	[8.2,	18.0]	1.369	[0.646,	2.903]	0.4109	

Abbreviations: CI=confidence interval; GEJ=gastro-esophageal junction; HR=hazard ratio; IRC=independent review committee; KME=Kaplan-Meier Estimate of time to event (months); N=number of patients; n=number of patients with event; NC=not calculated.

[a] Based on the Brookmeyer-Crowley Method, [b] Calculated from Cox proportional hazards model, [c] Based on 2-sided log-rank test, unadjusted for multiplicity, and [d] Two-sided Type 3 Wald test p-value from Cox proportional hazards model including treatment, subgroup, and treatment x subgroup interaction.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1002.3.1: Summary of Progression-Free Survival (INV) - Full Analysis Set

	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)	Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6
Number of patients at risk	283 (100.0%)	282 (100.0%)	
Number of patients with events	171 (60.4%)	198 (70.2%)	
Number of patients censored	112 (39.6%)	84 (29.8%)	
Kaplan-Meier estimates of time to event (months)			
Quartiles, 95% CI [a]			
50%	10.4 [8.4, 11.8]	8.4 [7.8, 9.6]	
Cox proportional hazards model			
Stratified HR, 95% CI			0.728 [0.590, 0.898]
Log-rank test			
Two-sided stratified log-rank p-value			0.0031

Abbreviations: CI=confidence interval; HR=hazard ratio; INV=investigator; N=number of patients.
Note: The stratification factors are region (Asia vs. Non-Asia), the number of organs with metastatic sites (0 to 2 vs. ≥ 3) and prior gastrectomy (Yes or No).
[a] Based on the Brookmeyer-Crowley Method.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1002.3.2: Type of Events and Censoring of Progression-Free Survival (INV) - Full Analysis Set

	Zolbetuximab + mFOLFOX6 (N=283)	Placebo + mFOLFOX6 (N=282)
Number of patients at risk	283 (100.0%)	282 (100.0%)
Number of patients with events	171 (60.4%)	198 (70.2%)
rPD	130 (45.9%)	159 (56.4%)
No rPD, but death recorded on eCRF	41 (14.5%)	39 (13.8%)
Number of patients censored	112 (39.6%)	84 (29.8%)

Abbreviations: eCRF= electronic case report form; INV=investigator; N=number of patients; rPD=radiological progressive disease.
ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.1002.3.3: Summary of Progression-Free Survival (INV) by Subgroups - Full Analysis Set

Subgroup	Zolbetuximab + mFOLFOX6			Placebo + mFOLFOX6			Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6			Test for Interaction p-value[d]
	N	n (%)	KME [95% CI][a]	N	n (%)	KME [95% CI][a]	HR [95% CI][b]	Log-rank p-value[c]		
Age Group 1										
<=65 years	181	113 (62.4)	10.4 [8.3, 11.8]	181	121 (66.9)	8.5 [7.8, 10.3]	0.798 [0.616, 1.033]	0.0862	0.2246	
>65 years	102	58 (56.9)	10.4 [8.3, 12.5]	101	77 (76.2)	8.2 [6.3, 9.3]	0.631 [0.448, 0.890]	0.0080		
Age Group 2										
<=75 years	267	162 (60.7)	10.4 [8.4, 11.8]	260	184 (70.8)	8.3 [7.8, 9.6]	0.726 [0.587, 0.898]	0.0030	0.7667	
>75 years	16	9 (56.3)	9.7 [4.4, 18.1]	22	14 (63.6)	9.3 [5.9, 10.7]	0.843 [0.357, 1.988]	0.6802		
Sex										
Male	176	112 (63.6)	10.4 [8.3, 12.3]	175	128 (73.1)	8.3 [7.6, 9.6]	0.723 [0.559, 0.934]	0.0127	0.9940	
Female	107	59 (55.1)	10.4 [8.3, 12.4]	107	70 (65.4)	8.4 [6.4, 10.5]	0.742 [0.524, 1.051]	0.0907		
Race										
White	140	92 (65.7)	8.3 [6.5, 10.6]	134	94 (70.1)	8.7 [7.6, 10.4]	0.975 [0.731, 1.301]	0.8609	0.0076	
Asian	96	58 (60.4)	12.4 [10.4, 15.2]	97	66 (68.0)	8.3 [5.8, 10.2]	0.521 [0.364, 0.746]	0.0003		
Tobacco History										
Current	26	16 (61.5)	10.4 [7.9, 13.2]	25	19 (76.0)	8.4 [4.3, 11.0]	0.729 [0.374, 1.423]	0.3544	0.9311	
Former	113	73 (64.6)	10.3 [8.3, 12.5]	118	82 (69.5)	8.9 [7.9, 10.2]	0.749 [0.544, 1.031]	0.0747		
Never	142	82 (57.7)	8.4 [7.9, 12.3]	137	96 (70.1)	8.0 [6.3, 10.2]	0.710 [0.528, 0.956]	0.0234		
Region										
Asia	88	54 (61.4)	12.4 [10.4, 15.1]	89	60 (67.4)	8.3 [6.2, 10.2]	0.545 [0.375, 0.792]	0.0013	0.0631	
Non-Asia	195	117 (60.0)	8.8 [7.5, 10.6]	193	138 (71.5)	8.4 [7.6, 10.2]	0.848 [0.663, 1.086]	0.1893		
Number of Organs with Metastatic Sites										
0-2	219	127 (58.0)	11.0 [10.2, 12.5]	219	147 (67.1)	8.9 [8.0, 10.3]	0.709 [0.558, 0.901]	0.0048	0.5801	
>=3	64	44 (68.8)	6.5 [6.1, 9.3]	63	51 (81.0)	6.8 [6.0, 8.4]	0.834 [0.556, 1.252]	0.3754		
Prior Gastrectomy (total or partial)										
Yes	84	49 (58.3)	12.3 [10.4, 15.2]	82	59 (72.0)	9.3 [7.1, 11.3]	0.605 [0.413, 0.887]	0.0091	0.2521	
No	199	122 (61.3)	8.8 [7.7, 11.0]	200	139 (69.5)	8.3 [6.5, 9.6]	0.803 [0.629, 1.026]	0.0787		

Medizinischer Nutzen, medizinischer Zusatznutzen, Patientengruppen mit therap. bedeutsamem Zusatznutzen

Subgroup	Zolbetuximab + mFOLFOX6			Placebo + mFOLFOX6			Zolbetuximab + mFOLFOX6 vs. Placebo + mFOLFOX6			Test for Interaction p-value[d]
	N	n (%)	KME [95% CI][a]	N	n (%)	KME [95% CI][a]	HR [95% CI][b]	Log-rank p-value[c]		
Histology (Tumor Type)										
Diffuse	82	53 (64.6)	10.4 [8.3, 14.9]	117	78 (66.7)	10.2 [8.3, 10.8]	0.828 [0.583, 1.177]	0.2932		0.1594
Intestinal	70	46 (65.7)	10.4 [6.7, 13.2]	66	51 (77.3)	8.0 [5.7, 9.3]	0.548 [0.362, 0.831]	0.0042		
Mixed/Other	81	53 (65.4)	9.7 [7.7, 12.4]	55	42 (76.4)	8.4 [6.3, 10.3]	0.846 [0.564, 1.270]	0.4190		
Tumor Location 1										
Gastric	219	131 (59.8)	10.6 [8.5, 12.3]	210	146 (69.5)	8.3 [7.8, 10.2]	0.710 [0.560, 0.901]	0.0046		0.5492
GEJ	64	40 (62.5)	8.3 [6.2, 11.6]	72	52 (72.2)	8.4 [6.2, 9.8]	0.845 [0.559, 1.279]	0.4242		
Tumor Location 2										
Gastric Proximal	73	48 (65.8)	8.4 [8.0, 12.3]	59	41 (69.5)	8.4 [7.1, 10.7]	0.822 [0.539, 1.253]	0.3611		0.4207
Gastric Distal	91	48 (52.7)	13.2 [10.4, 15.2]	87	57 (65.5)	8.5 [6.3, 10.5]	0.621 [0.422, 0.913]	0.0142		
Tumor Location 3										
GEJ Proximal	30	17 (56.7)	10.4 [6.5, 17.8]	26	23 (88.5)	5.9 [4.1, 8.4]	0.395 [0.206, 0.756]	0.0039		0.0230
GEJ Distal	19	12 (63.2)	8.3 [4.3, 15.1]	31	21 (67.7)	10.2 [6.2, 12.1]	1.189 [0.580, 2.437]	0.6363		

Abbreviations: CI=confidence interval; GEJ=gastro-esophageal junction; HR=hazard ratio; INV=investigator; KME=Kaplan-Meier Estimate of time to event (months); N=number of patients; n=number of patients with event.

[a] Based on the Brookmeyer-Crowley Method, [b] Calculated from Cox proportional hazards model, [c] Based on 2-sided log-rank test, unadjusted for multiplicity, and [d] Two-sided Type 3 Wald test p-value from Cox proportional hazards model including treatment, subgroup, and treatment x subgroup interaction.

ASTELLAS Data Cutoff Date: 09SEP22


Anhang 4-G4 *Post hoc* Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022

Anhang 4-G4 Gesamtüberleben und Progressionsfreies Überleben

Anhang 4-G4 Progressionsfreies Überleben (IRC und INV)

2. Kaplan-Meier-Plots


Figure 301.1.1002.2: Kaplan-Meier Plot of Progression-Free Survival (IRC) - Full Analysis Set



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; IRC=independent review committee; N=number of patients.

ASTELLAS Data Cutoff Date: 09SEP22


Figure 301.1.1002.2.12: Kaplan-Meier Plot of Progression-Free Survival (IRC) by Tumor Location 3 - Full Analysis Set
Tumor Location 3: GEJ Proximal



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; GEJ=gastro-esophageal junction; IRC=independent review committee; N=number of patients.


ASTELLAS Data Cutoff Date: 09SEP22

Figure 301.1.1002.2.12: Kaplan-Meier Plot of Progression-Free Survival (IRC) by Tumor Location 3 - Full Analysis Set
Tumor Location 3: GEJ Distal



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; GEJ=gastro-esophageal junction; IRC=independent review committee; N=number of patients.
ASTELLAS Data Cutoff Date: 09SEP22


Figure 301.1.1002.2.4: Kaplan-Meier Plot of Progression-Free Survival (IRC) by Race - Full Analysis Set
Race: White



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; IRC=independent review committee; N=number of patients.
ASTELLAS Data Cutoff Date: 09SEP22


Figure 301.1.1002.2.4: Kaplan-Meier Plot of Progression-Free Survival (IRC) by Race - Full Analysis Set

Race: Asian



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; IRC=independent review committee; N=number of patients.


ASTELLAS Data Cutoff Date: 09SEP22

Figure 301.1.1002.3: Kaplan-Meier Plot of Progression-Free Survival (INV) - Full Analysis Set

Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; INV=investigator; N=number of patients.

ASTELLAS Data Cutoff Date: 09SEP22


Figure 301.1.1002.3.12: Kaplan-Meier Plot of Progression-Free Survival (INV) by Tumor Location 3 - Full Analysis Set
Tumor Location 3: GEJ Proximal



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; GEJ=gastro-esophageal junction; INV=investigator; N=number of patients.

ASTELLAS Data Cutoff Date: 09SEP22


Figure 301.1.1002.3.12: Kaplan-Meier Plot of Progression-Free Survival (INV) by Tumor Location 3 - Full Analysis Set
Tumor Location 3: GEJ Distal



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; GEJ=gastro-esophageal junction; INV=investigator; N=number of patients.
ASTELLAS Data Cutoff Date: 09SEP22

Figure 301.1.1002.3.4: Kaplan-Meier Plot of Progression-Free Survival (INV) by Race - Full Analysis Set

Race: White




Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; INV=investigator; N=number of patients.

ASTELLAS Data Cutoff Date: 09SEP22

Figure 301.1.1002.3.4: Kaplan-Meier Plot of Progression-Free Survival (INV) by Race - Full Analysis Set

Race: Asian



Abbreviations: # at Risk=number of patients at risk; CI=confidence interval; INV=investigator; N=number of patients.

ASTELLAS Data Cutoff Date: 09SEP22

Anhang 4-G4 *Post hoc* Analysen für die Studie SPOTLIGHT, primärer Datenschnitt 09.09.2022

Anhang 4-G4 Patientenberichtete Endpunkte

Anhang 4-G4 Symptomatik und Gesundheitsbezogene Lebensqualität anhand des EORTC QLQ-C30

1. Rücklaufquoten

Table 301.1.3001.1: EORTC QLQ-C30 - Completion Status - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	257/278 (92.4%)	257/280 (91.8%)	257/282 (91.1%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.1: EORTC QLQ-C30 - Completion Status - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.1: EORTC QLQ-C30 - Completion Status - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.5: EORTC QLQ-C30 - Completion Status of Global Health Status - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.5: EORTC QLQ-C30 - Completion Status of Global Health Status - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.5: EORTC QLQ-C30 - Completion Status of Global Health Status - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.6: EORTC QLQ-C30 - Completion Status of Physical Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.6: EORTC QLQ-C30 - Completion Status of Physical Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.6: EORTC QLQ-C30 - Completion Status of Physical Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.7: EORTC QLQ-C30 - Completion Status of Role Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.7: EORTC QLQ-C30 - Completion Status of Role Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.7: EORTC QLQ-C30 - Completion Status of Role Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.8: EORTC QLQ-C30 - Completion Status of Emotional Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.8: EORTC QLQ-C30 - Completion Status of Emotional Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.8: EORTC QLQ-C30 - Completion Status of Emotional Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.9: EORTC QLQ-C30 - Completion Status of Cognitive Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.9: EORTC QLQ-C30 - Completion Status of Cognitive Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.9: EORTC QLQ-C30 - Completion Status of Cognitive Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.10: EORTC QLQ-C30 - Completion Status of Social Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.10: EORTC QLQ-C30 - Completion Status of Social Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.10: EORTC QLQ-C30 - Completion Status of Social Functioning - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.11: EORTC QLQ-C30 - Completion Status of Fatigue - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.11: EORTC QLQ-C30 - Completion Status of Fatigue - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.11: EORTC QLQ-C30 - Completion Status of Fatigue - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.12: EORTC QLQ-C30 - Completion Status of Nausea and Vomiting - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.12: EORTC QLQ-C30 - Completion Status of Nausea and Vomiting - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.12: EORTC QLQ-C30 - Completion Status of Nausea and Vomiting - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 27 Day 1	4/ 4 (100.0%)	4/227 (1.8%)	4/283 (1.4%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 27 Day 22	3/ 4 (75.0%)	3/227 (1.3%)	3/283 (1.1%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 28 Day 1	3/ 3 (100.0%)	3/227 (1.3%)	3/283 (1.1%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 28 Day 22	2/ 3 (66.7%)	2/227 (0.9%)	2/283 (0.7%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 29 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 29 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 30 Day 1	2/ 2 (100.0%)	2/227 (0.9%)	2/283 (0.7%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 30 Day 22	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 1 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 31 Day 1	1/ 2 (50.0%)	1/227 (0.4%)	1/283 (0.4%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 32 Day 1	0	0/227 (0.0%)	0/283 (0.0%)	1/ 1 (100.0%)	1/235 (0.4%)	1/282 (0.4%)
Study Disc	136/223 (61.0%)	137/270 (50.7%)	137/283 (48.4%)	147/222 (66.2%)	147/264 (55.7%)	147/282 (52.1%)
30 D SFU Z/P	67/203 (33.0%)	68/251 (27.1%)	69/283 (24.4%)	81/199 (40.7%)	81/241 (33.6%)	81/282 (28.7%)
90 D SFU Z/P	83/179 (46.4%)	83/230 (36.1%)	83/283 (29.3%)	70/160 (43.8%)	70/209 (33.5%)	71/282 (25.2%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo; 90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.13: EORTC QLQ-C30 - Completion Status of Pain - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Baseline	257/279 (92.1%)	257/282 (91.1%)	257/283 (90.8%)	258/278 (92.8%)	258/280 (92.1%)	258/282 (91.5%)
Cycle 1 Day 22	186/257 (72.4%)	187/273 (68.5%)	187/283 (66.1%)	212/266 (79.7%)	212/271 (78.2%)	212/282 (75.2%)
Cycle 2 Day 1	214/244 (87.7%)	214/263 (81.4%)	217/283 (76.7%)	230/263 (87.5%)	231/270 (85.6%)	231/282 (81.9%)
Cycle 2 Day 22	156/222 (70.3%)	157/257 (61.1%)	157/283 (55.5%)	184/243 (75.7%)	185/264 (70.1%)	185/282 (65.6%)
Cycle 3 Day 1	191/215 (88.8%)	197/254 (77.6%)	200/283 (70.7%)	204/228 (89.5%)	204/259 (78.8%)	204/282 (72.3%)
Cycle 3 Day 22	160/204 (78.4%)	161/250 (64.4%)	161/283 (56.9%)	156/219 (71.2%)	156/257 (60.7%)	156/282 (55.3%)
Cycle 4 Day 1	169/192 (88.0%)	174/244 (71.3%)	178/283 (62.9%)	170/202 (84.2%)	170/254 (66.9%)	171/282 (60.6%)
Cycle 4 Day 22	128/179 (71.5%)	128/241 (53.1%)	128/283 (45.2%)	132/184 (71.7%)	132/251 (52.6%)	132/282 (46.8%)
Cycle 5 Day 1	149/171 (87.1%)	155/239 (64.9%)	156/283 (55.1%)	147/170 (86.5%)	147/251 (58.6%)	147/282 (52.1%)
Cycle 5 Day 22	113/151 (74.8%)	114/236 (48.3%)	114/283 (40.3%)	119/150 (79.3%)	120/248 (48.4%)	120/282 (42.6%)
Cycle 6 Day 1	121/135 (89.6%)	125/236 (53.0%)	125/283 (44.2%)	120/139 (86.3%)	122/244 (50.0%)	122/282 (43.3%)
Cycle 6 Day 22	100/125 (80.0%)	102/234 (43.6%)	102/283 (36.0%)	93/120 (77.5%)	93/242 (38.4%)	93/282 (33.0%)
Cycle 7 Day 1	107/118 (90.7%)	111/234 (47.4%)	111/283 (39.2%)	91/103 (88.3%)	91/240 (37.9%)	91/282 (32.3%)
Cycle 7 Day 22	80/105 (76.2%)	80/232 (34.5%)	80/283 (28.3%)	66/ 93 (71.0%)	66/239 (27.6%)	66/282 (23.4%)
Cycle 8 Day 1	80/ 96 (83.3%)	81/231 (35.1%)	81/283 (28.6%)	73/ 85 (85.9%)	73/239 (30.5%)	73/282 (25.9%)
Cycle 8 Day 22	70/ 86 (81.4%)	71/231 (30.7%)	71/283 (25.1%)	56/ 75 (74.7%)	56/237 (23.6%)	56/282 (19.9%)
Cycle 9 Day 1	72/ 82 (87.8%)	73/229 (31.9%)	73/283 (25.8%)	53/ 63 (84.1%)	53/237 (22.4%)	53/282 (18.8%)
Cycle 9 Day 22	54/ 69 (78.3%)	54/229 (23.6%)	54/283 (19.1%)	46/ 58 (79.3%)	46/237 (19.4%)	46/282 (16.3%)
Cycle 10 Day 1	55/ 63 (87.3%)	58/229 (25.3%)	58/283 (20.5%)	46/ 49 (93.9%)	47/237 (19.8%)	47/282 (16.7%)
Cycle 10 Day 22	47/ 59 (79.7%)	47/229 (20.5%)	47/283 (16.6%)	35/ 43 (81.4%)	35/237 (14.8%)	35/282 (12.4%)
Cycle 11 Day 1	50/ 56 (89.3%)	50/229 (21.8%)	50/283 (17.7%)	37/ 40 (92.5%)	37/237 (15.6%)	37/282 (13.1%)
Cycle 11 Day 22	34/ 51 (66.7%)	35/229 (15.3%)	35/283 (12.4%)	22/ 35 (62.9%)	22/236 (9.3%)	22/282 (7.8%)
Cycle 12 Day 1	42/ 48 (87.5%)	43/229 (18.8%)	43/283 (15.2%)	32/ 34 (94.1%)	32/236 (13.6%)	32/282 (11.3%)
Cycle 12 Day 22	32/ 45 (71.1%)	32/229 (14.0%)	32/283 (11.3%)	20/ 30 (66.7%)	20/235 (8.5%)	20/282 (7.1%)
Cycle 13 Day 1	37/ 43 (86.0%)	37/229 (16.2%)	37/283 (13.1%)	25/ 27 (92.6%)	25/235 (10.6%)	25/282 (8.9%)
Cycle 13 Day 22	33/ 37 (89.2%)	33/228 (14.5%)	34/283 (12.0%)	15/ 26 (57.7%)	15/235 (6.4%)	15/282 (5.3%)

Abbreviations: EORTC QLQ-C30=european organisation for research and treatment of cancer quality of life questionnaire-core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22

Table 301.1.3001.13: EORTC QLQ-C30 - Completion Status of Pain - Full Analysis Set

Analysis Visit	Zolbetuximab + mFOLFOX6 (N=283)			Placebo + mFOLFOX6 (N=282)		
	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)	Compliance Rate [1] (based on expected assessments)	Compliance Rate [2] (adjusted for deaths)	Completion Rate [3] (unadjusted)
Cycle 14 Day 1	31/ 32 (96.9%)	31/228 (13.6%)	31/283 (11.0%)	23/ 25 (92.0%)	23/235 (9.8%)	23/282 (8.2%)
Cycle 14 Day 22	24/ 30 (80.0%)	24/228 (10.5%)	24/283 (8.5%)	13/ 22 (59.1%)	13/235 (5.5%)	13/282 (4.6%)
Cycle 15 Day 1	26/ 29 (89.7%)	26/228 (11.4%)	26/283 (9.2%)	19/ 20 (95.0%)	19/235 (8.1%)	19/282 (6.7%)
Cycle 15 Day 22	22/ 29 (75.9%)	22/228 (9.6%)	22/283 (7.8%)	8/ 15 (53.3%)	8/235 (3.4%)	8/282 (2.8%)
Cycle 16 Day 1	24/ 26 (92.3%)	25/228 (11.0%)	25/283 (8.8%)	11/ 13 (84.6%)	11/235 (4.7%)	11/282 (3.9%)
Cycle 16 Day 22	18/ 23 (78.3%)	19/228 (8.3%)	19/283 (6.7%)	6/ 11 (54.5%)	6/235 (2.6%)	6/282 (2.1%)
Cycle 17 Day 1	19/ 20 (95.0%)	19/227 (8.4%)	19/283 (6.7%)	10/ 10 (100.0%)	10/235 (4.3%)	10/282 (3.5%)
Cycle 17 Day 22	14/ 18 (77.8%)	14/227 (6.2%)	14/283 (4.9%)	3/ 8 (37.5%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 18 Day 1	16/ 17 (94.1%)	16/227 (7.0%)	16/283 (5.7%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 18 Day 22	11/ 16 (68.8%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 19 Day 1	13/ 15 (86.7%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 19 Day 22	11/ 15 (73.3%)	11/227 (4.8%)	11/283 (3.9%)	4/ 7 (57.1%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 20 Day 1	13/ 14 (92.9%)	13/227 (5.7%)	13/283 (4.6%)	7/ 7 (100.0%)	7/235 (3.0%)	7/282 (2.5%)
Cycle 20 Day 22	9/ 13 (69.2%)	9/227 (4.0%)	9/283 (3.2%)	4/ 6 (66.7%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 21 Day 1	11/ 12 (91.7%)	11/227 (4.8%)	11/283 (3.9%)	5/ 6 (83.3%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 21 Day 22	9/ 11 (81.8%)	9/227 (4.0%)	9/283 (3.2%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 22 Day 1	8/ 10 (80.0%)	8/227 (3.5%)	8/283 (2.8%)	5/ 5 (100.0%)	5/235 (2.1%)	5/282 (1.8%)
Cycle 22 Day 22	5/ 9 (55.6%)	5/227 (2.2%)	5/283 (1.8%)	3/ 5 (60.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 23 Day 1	9/ 9 (100.0%)	9/227 (4.0%)	9/283 (3.2%)	4/ 4 (100.0%)	4/235 (1.7%)	4/282 (1.4%)
Cycle 23 Day 22	4/ 7 (57.1%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 24 Day 1	7/ 7 (100.0%)	7/227 (3.1%)	7/283 (2.5%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 24 Day 22	4/ 6 (66.7%)	4/227 (1.8%)	4/283 (1.4%)	1/ 3 (33.3%)	1/235 (0.4%)	1/282 (0.4%)
Cycle 25 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	3/ 3 (100.0%)	3/235 (1.3%)	3/282 (1.1%)
Cycle 25 Day 22	4/ 5 (80.0%)	4/227 (1.8%)	4/283 (1.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)
Cycle 26 Day 1	5/ 5 (100.0%)	5/227 (2.2%)	5/283 (1.8%)	2/ 2 (100.0%)	2/235 (0.9%)	2/282 (0.7%)
Cycle 26 Day 22	1/ 4 (25.0%)	1/227 (0.4%)	1/283 (0.4%)	0/ 2 (0.0%)	0/235 (0.0%)	0/282 (0.0%)

Abbreviations: EORTC QLQ-C30=European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30; N=number of patients; 30 D SFU Z/P=30 day safety follow up visit - Zolbetuximab/Placebo;

90 D FU Z/P=90 day follow up visit Zolbetuximab/Placebo; Study Disc=study drug discontinuation visit.

[1] Compliance (or adjusted completion) rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to progression, death or other reasons at respective visit. minimum requirements for scoring is defined as at least one scale with non-missing values.

[2] Compliance rate (adjusted for deaths) is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects with PRO from FAS without patients not expected due to death at respective visit.

[3] Unadjusted completion rate is calculated as the number of subjects with minimum requirements for scoring divided by the number of subjects in the FAS.

ASTELLAS Data Cutoff Date: 09SEP22