

Create a webpage which look like the image given below:

[illegible]

```

                                body content
                                &lt;/body&gt;
                                &lt;/html&gt;
                                <CODE>
                                <PRE><hr>
                                <center><b>Created by : Catherine Joshi</b></center>
                                </BODY>
                                </HTML>

```

OUTPUT:



Hypertext Markup Language

Introduction

HTML stands for **Hypertext Markup Language**. It is used to design webpages using markup language. HTML is the combination of *Hypertext* and *Markup* Language. Originally, HTML was developed with the intent of defining the structure of documents like headings, paragraphs, lists, and so forth to facilitate the sharing of scientific information between researchers. Now, HTML is being widely used to format webpages with the help of different tags available in HTML language.

Structure of a simple HTML document

```
<!DOCTYPE html>
<html>
<head>
  <title> Give the title here </title>
</head>
<body>
  body content
</body>
</html>

```

Created by : Catherine Joshi

PROGRAM 2

Create an HTML file to link to different HTML page which contains images and tables.

Source Code:

Home.html

```
<html>
  <head>
    <title>Birds</title>
  </head>
  <body>
    <center><h2>Birds of Kerala</h2>
    <p>Due to the climate and evergreen forests, there are many
    kinds of birds found all over Kerala. Some of them are
    common and some others are found in forest areas. Here are
    some of the birds of Kerala.</p>
    <a href="images.html">Images</a><br>
    <a href="table.html">Tabular data</a></center>
  </body>
</html>
```

Images.html

```
<html>
  <head>
    <title>Images</title>
  </head>
  <body>
    <center><h2>Birds of Kerala</h2>
    <table>
      <tr>
        <td></td>
        <td></td>
        <td></td>
      </tr>
      <tr>
```

```

        <td></td>
        <td></td>
        <td></td>
    </tr>
</table><br><br>
<a href="home.html">Home</a><br>
<a href="table.html">Tabular data</a><center>
</body>
</html>

```

Table.html

```

<html>
    <head>
        <title>Tabular data</title>
    </head>
    <body>
        <center><h2>Birds of Kerala</h2>
        <table border="1">
            <tr>
                <th>Species Name</th>
                <th>Local Name</th>
                <th>Size</th>
            </tr>
            <tr>
                <td>Grey Hornbill</td>
                <td>Vezhambal</td>
                <td>41.4x30.0</td>
            </tr>
            <tr>
                <td>Black Eagle</td>
                <td>Karimparundu </td>
                <td>62.7x49.9 mm</td>
            </tr>
            <tr>
                <td>Whistling Teal</td>
                <td>Yeranda</td>
            </tr>
        </table>
    </body>
</html>

```

```

<td>46.9X36.8 mm</td>
</tr>
</table><br><br>
<a href="home.html">Home</a><br>
<a href="images.html">Images</a></center>
</body>
</html>

```

Output:



Birds of Kerala

Due to the climate and evergreen forests, there are many kinds of birds found all over Kerala. Some of them are common and some others are found in forest areas. Here are some of the birds of Kerala.

[Images](#)
[Tabular data](#)



Birds of Kerala



[Home](#)
[Tabular data](#)



Birds of Kerala

Species Name	Local Name	Size
Grey Hornbill	Vezhambal	41.4x30.0
Black Eagle	Karimparundu	62.7x49.9 mm
Whistling Teal	Yeranda	46.9X36.8 mm

[Home](#)
[Images](#)

PROGRAM 3

Create an HTML file to link to different HTML page which contains images and tables and link within page.

Source Code:

Home.html

```
<html>
  <head>
    <title>Birds</title>
  </head>
  <body>
    <center><h2>Birds of Kerala</h2>
    <p>Due to the climate and evergreen forests, there are many
    kinds of birds found all over Kerala. Some of them are
    common and some others are found in forest areas. Here are
    some of the birds of Kerala.</p>
    <a href="images.html">Images</a><br>
    <a href="table.html">Tabular data</a></center>
    <center><a href=#Little Cormorant>Little Cormorant</a><br>
      <a href=#Indian Darter or Snake –Bird >Indian Darter
      or Snake –Bird </a><br>
      <a href=#Eastern Grey Heron>Eastern Grey
Heron</a><br>
      <a href=#Indian Little Green Bittern>Indian Little
      Green Bittern</a><br>
      <a href=#Openbill Stork>Openbill Stork</a><br>
      <a href=#Cattle Egret>Cattle Egret</a><br>
    </center>
    <a id=Little Cormorant></a><br>
    <p><h2>Little Cormorant</h2>
      Local Name: Kakkattaravu Kakkattaravu<br>Size:
      Somewhat larger than the Jungle Crow, with longer
      neck and tail.<br>
      Field Characters: A glistening black duck-like water
      bird with a longer stiff tail and slender compressed bill
      sharply hooked at the tip.
      A small white patch on throat, and suggestion of a
      nuchal crest. Sexes alike. Singly or in flocks, on or
      near water.
```

Cormorants are almost exclusively fish-eaters. They chase and capture their quarry under water, being expert divers and submarine swimmers.

Flocks sometimes hunt in concert in the manner of pelicans. They also hunt individually. When satiated, the birds may be seen sitting on a rock or sandbank, or on stakes or trees in or near the water, drying themselves with outstretched wings.

Eggs: 4 or 5, long narrow ovals, bluish white in color with a chalky texture.
Average Size:

44.8x29.0mm </p>

<p><h2>Indian Darter or Snake –Bird </h2>

Local Name: Chera-Kozhi
Size: About that of the Kite
Field Characters: A black, cormorant-like water bird with a long and very slender S-shaped neck, narrow head and pointed dagger bill. Back heavily streaked with silver-grey; head and neck velvety brown; chin and throat whitish.

Tail long, stiff and wedge-shaped(almost rounded).

Sexes alike. Singly or in loose parties on tanks.

Resident. Fairly common, but not abundant.

Affects open placid streams, village tanks, and dammed reservoirs. On the last the exposed tops of the submerged dead tress provide favorite lookout and wing-drying posts. This is also found in Ceylon , all India, Burma, Malaysia, Indochinese countries to the Phluppines.

Other races occur in America, Africa and

Australia .
Eggs: 3 or 4, very elongated,pale greenish blue with whitish chalky coating.

Average size: 52.9x33.5mm</p>

<p><h2>Eastern Grey Heron </h2>

Local Name: Charamunti
Size: About the same as the Purple Heron
Field Characters: Like the

Purple Heron but ashy grey above with white

crown and neck; grayish white below. Long black occipital crest. Elongated white feathers on the breast with some black streaks.

A conspicuous black dotted line down the middle of the fore neck. In the female the crest and pectoral plumes are less developed.

Solitary, on reedy marshes. Distributed outside Beralia is in Ceylon, all India, Burma, Malaysia. This race ranges from Iraq eastward to

the Philippines. The species overall is found throughout the old World. Eggs: Almost

indistinguishable from those of the Purple Heron.

Average size: 58.6X43.5 mm.

Indian Little Green Bittern

Local Name: Chinna-kokku. Size: About the same as, or slightly smaller than, the Pond Heron.

Field Characters: Like the Pond Heron in general effect but largely blackish grey, glossy dark green and bronze-green above; ashy grey below. Top of head and long occipital crest black, glossed with dark green. Chin and throat white. Sexes alike. Solitary, in screw pine thickets and brushwood near water. Resident. Not common.

Low country, and to at least 2000 ft. (c. 600 m). Keeps to brushwood in and on the banks of rocky streams.

Distributed outside Kerala

is in Ceylon, India, Burma, Thailand, South China, Malaysia. The species in numerous races is practically cosmopolitan.

Eggs: 3 to 5, very like the Pond Heron's. Average size: 39.5X29.7 mm

Openbill Stork

Local Name: Cherakokkan. Size: Smaller than vulture. Long legged; standing about 2 1/2 ft. (75 cm.) high.

Field Characters: A small stork of grayish white or white plumage, with black in the wings. The curious reddish black bill,

with arching mandibles leaving a narrow open gap between them, is diagnostic. Sexes alike. Twos and threes or flocks, at tanks and marshes. Resident. According to Ferguson (1904, JBNHS 16: 15) 'very common in Travancore where large flocks may be met with on the marshy borders of all the larger tanks and freshwater lakes, sharing the trees for roosting with flocks of Darters'. Since then its status in Kerala seems to have altered considerably. It is now found only sparsely and in small numbers. Distributed outside Kerala is in Ceylon , all India , Assam , Burma , Thailand and Cambodia . It is apparently adapted in some way to deal with the thick hard shells of the large Pila snails found on marshes, the soft body and viscera of which form such an important food item of this species. It also eats frogs, crabs, large insects.
Eggs: 2 to 4 , sullied white.
Average size: 57.9X41.2 mm.</p>

<p><h2>Cattle Egret</h2>

Local Name: Kalimunti
 Size: Same as the Little Egret. About that of the domestic hen; lankier and more leggy.

Field Characters: Snow-white overall, very similar to the Little Egret, with which it may sometimes be confused. But the colour of its bill-yellow (instead of black) – is always diagnostic. In the breeding season the head, neck and back turn a distinctive golden, orange-buff. Sexes alike. Flocks, attending on grazing cattle. Resident (presumably). Apparently much more abundant in Ferguson 's day than at present, but still the commonest of all the egrets. Frequents paddy fields along the backwaters, cultivated and fallow land, chiefly in attendance on grazing cattle. Distributed outside Kerala is in Ceylon, all India, Burma, Thailand, Malaysia to the Philippines

and Eastern China. The nominate race occurs in Africa, S. Europe and W. Asia. Its food consists very largely of bluebottle flies picked off the grass or weeds. The bird poises its bill, sways its neck comically from side to side to get a correct aim, as it were, and jabs at the quarry. Grasshoppers and other insects, as well as frogs and lizards, are also eaten.
 Eggs: 3 to 5, pale skim-milk blue in colour, unmarked.
 Average size: 44.1X33.6 mm. </p>

 Go to Top

</body>

</html>

Images.html

<html>

<head>

<title>Images</title>

</head>

<body>

<center><h2>Birds of Kerala</h2>

<table>

<tr>

<td></td>

<td></td>

<td></td>

</tr>

<tr>

<td></td>

<td></td>

<td></td>

</tr>

</table>


```
        <a href="home.html">Home</a><br>
        <a href="table.html">Tabular data</a><center>
</body>
</html>
```

Table.html

```
<html>
  <head>
    <title>Tabular data</title>
  </head>
  <body>
    <center><h2>Birds of Kerala</h2>
    <table border="1">
      <tr>
        <th>Species Name</th>
        <th>Local Name</th>
        <th>Size</th>
      </tr>
      <tr>
        <td>Grey Hornbill</td>
        <td>Vezhambal</td>
        <td>41.4x30.0</td>
      </tr>
      <tr>
        <td>Black Eagle</td>
        <td>Karimparundu </td>
        <td>62.7x49.9 mm</td>
      </tr>
      <tr>
        <td>Whistling Teal</td>
        <td>Yeranda</td>
        <td>46.9X36.8 mm</td>
      </tr>
    </table><br><br>
    <a href="home.html">Home</a><br>
    <a href="images.html">Images</a></center>
  </body>
</html>
```

Output:

Birds of Kerala

Due to the climate and evergreen forests, there are many kinds of birds found all over Kerala. Some of them are common and some others are found in forest areas. Here are some of the birds of Kerala.

[Images](#)
[Tabular data](#)
[Little Cormorant](#)
[Indian Darter or Snake –Bird](#)
[Eastern Grey Heron](#)
[Indian Little Green Bittern](#)
[Openbill Stork](#)
[Cattle Egret](#)

Little Cormorant

Local Name: Kakkattaravu Kakkattaravu

Size: Somewhat larger than the Jungle Crow, with longer neck and tail.

Field Characters: A glistening black duck-like water bird with a longer stiff tail and slender compressed bill sharply hooked at the tip. A small white patch on throat, and suggestion of a nuchal crest. Sexes alike. Singly or in flocks, on or near water. Cormorants are almost exclusively fish-eaters. They chase and capture their quarry under water, being expert divers and submarine swimmers. Flocks sometimes hunt in concert in the manner of pelicans. They also hunt individually. When satiated, the birds may be seen sitting on a rock or sandbank, or on stakes or trees in or near the water, drying themselves with outstretched wings.

Eggs: 4 or 5, long narrow ovals, bluish white in color with a chalky texture.

Average Size: 44.8x29.0mm

Indian Darter or Snake –Bird

Local Name: Chera-Kozhi

Size: About that of the Kite

Field Characters: A black, cormorant-like water bird with a long and very slender S-shaped neck, narrow head and pointed dagger bill. Back heavily streaked with silver-grey; head and neck velvety brown; chin and throat whitish. Tail long, stiff and wedge-shaped(almost rounded). Sexes alike. Singly or in loose parties on tanks. Resident. Fairly common, but not abundant. Affects open placid streams, village tanks, and dammed reservoirs. On the last the exposed tops of the submerged dead tress provide favorite lookout and wing-drying posts. This is also found in Ceylon , all India, Burma, Malaysia, Indochinese countries to the Philippines. Other races occur in America, Africa and Australia .

Eggs: 3 or 4, very elongated,pale greenish blue with whitish chalky coating.

Average size: 52.9x33.5mm

Eastern Grey Heron

Local Name: Charamunti

Size: About the same as the Purple Heron

Field Characters: Like the Purple Heron but ashy grey above with white crown and neck; grayish white below. Long black occipital crest. Elongated white feathers on the breast with some black streaks. A conspicuous black dotted line down the middle of the fore neck. In the female the crest and pectoral plumes are less developed. Solitary, on reedy marshes. Distributed outside Berala is in Ceylon, all India, Burma, Malaysia . This race ranges from Iraq eastward to the Philippines The species overall is found throughout the old World.

Eggs: Almost indistinguishable from those of the Purple Heron.

Average size: 58.6X43.5 mm.

Openbill Stork

Local Name: Cherakokkan

Size: Smaller than vulture. Long legged; standing about 2 1/2 ft. (75 cm.) high.

Field Characters: A small stork of grayish white or white plumage, with black in the wings. The curious reddish black bill, with arching mandibles leaving a narrow open gap between them, is diagnostic.

Sexes alike. Twos and threes or flocks, at tanks and marshes. Resident. According to Ferguson (1904, JBNHS 16: 15) 'very common in Travancore where large flocks may be met with on the marshy borders of all the larger tanks and freshwater lakes, sharing the trees for roosting with flocks of Darters'. Since then its status in Kerala seems to have altered considerably. It is now found only sparsely and in small numbers. Distributed outside Kerala is in Ceylon , all India , Assam , Burma , Thailand and Cambodia . It is apparently adapted in some way to deal with the thick hard shells of the large Pila snails found on marshes, the soft body and viscera of which form such an important food item of this species. It also eats frogs, crabs, large insects.

Eggs: 2 to 4 , sullied white.

Average size: 57.9X41.2 mm.

Cattle Egret

Local Name: Kalimunti

Size: Same as the Little Egret. About that of the domestic hen; lankier and more leggy.

Field Characters: Snow-white overall, very similar to the Little Egret, with which it may sometimes be confused. But the colour of its bill-yellow (instead of black) – is always diagnostic. In the breeding season the head, neck and back turn a distinctive golden, orange-buff. Sexes alike. Flocks, attending on grazing cattle. Resident (presumably). Apparently much more abundant in Ferguson 's day than at present, but still the commonest of all the egrets. Frequents paddy fields along the backwaters, cultivated and fallow land, chiefly in attendance on grazing cattle. Distributed outside Kerala is in Ceylon, all India, Burma, Thailand, Malaysia to the Philippines and Eastern China. The nominate race occurs in Africa, S Europe and W.Asia. Its food consists very largely of bluebottle flies picked off the grass or weeds.

The bird poises its bill, sways its neck comically from side to side to get a correct aim, as it were, and jabs at the quarry. Grasshoppers and other insects, as well as frogs and lizards, are also eaten.

Eggs: 3 to 5, pale skim-milk blue in colour, unmarked.

Average size: 44.1X33.6 mm.

[Go to Top](#)



Birds of Kerala



[Home](#)
[Tabular data](#)



Birds of Kerala

Species Name	Local Name	Size
Grey Hornbill	Vezhambal	41.4x30.0
Black Eagle	Karimparundu	62.7x49.9 mm
Whistling Teal	Yeranda	46.9X36.8 mm

[Home](#)
[Images](#)

PROGRAM 4

Create a HTML page with different types of frames

Source Code:

Mainframe.html

```
<html>
  <head>
    <title>home</title>
  </head>
  <frameset rows="20%,*" border="1">
    <frame name = "f1" src = "heading.html"/>
    <frame name = "f2" src = "cont.html"/>
  </frameset>
</html>
```

Heading.html

```
<html>
  <head>
    <title>heading</title>
  </head>
  <body bgcolor="DarkTurquoise">
    <br><center><h2>Vidya Academy of Science and
      Technology</h2></center>
  </body>
</html>
```

Cont.html

```
<html>
  <head>
    <title>nav1</title>
  </head>
  <body bgcolor="PaleTurquoise">
    <p>Vidya Academy of Science & Technology (VAST) is a
    state-of-the-art engineering college conforming
    to international standards. This model engineering college is
    approved by AICTE vide approval
```

NO./06/ KER/ENGG/2002/91 dated 12/05/03 and affiliated to APJ Abdul Kalam Technological University.

In few years VAST has evolved and achieved recognition as a notable School of Engineering with its competent and committed faculty, high quality infrastructure and high technology teaching aids, and by providing a serene atmosphere that complements academic life. VAST has a holistic approach to education where academic training goes hand in hand with offerings that develop the body, mind and soul to prepare its graduates to be future leaders.</p>

<h3>Vision</h3>

<p>Progress through education</p>

<h3>Mission</h3>

<p>To seek, strive for and scale greater heights of quality education</p>

<iframe src="frames.html" width="1000" height="500" frameborder="1">

</iframe>

</body>

</html>

Frames.html

<html>

<head>

<title>home</title>

</head>

<frameset cols="20%,80%">

<frame name = "top" src = "nav1.html"/>

<frame name = "main" src = "navframe.html"/>

</frameset>

</html>

Nav1.html

<html>

<head>

<title>nav1</title>

</head>

```

<body bgcolor="Aqua">
    <h2>Photo Gallery</h2>
    <hr>
    <a href = "vyvidh.jpg" target= "main">VYVIDH
    2020</a><br><br><br>
    <a href = "exhibition.jpg" target= "main">ISRO
    EXHIBITION</a><br><br>
    <a href = "graduation.jpg" target=
    "main">GRADUATION      CEREMONY</a>
    <hr>
</body>
</html>

```

Navframe.html

```

<html>
    <head>
        <title>nav1</title>
    </head>
    <body bgcolor="LightSkyBlue">
    </body>
</html>

```

OUTPUT



Photo Gallery

[VYVIDH 2020](#)

[ISRO EXHIBITION](#)

[GRADUATION
CEREMONY](#)



PROGRAM 5

Program to create registration form in HTML

Source Code:

```
<html>
  <head>
    <title>
      Registration form
    </title>
  </head>
  <body>
    <center>
      <h1>Registration form</h1>
      <form name="regform" action="">
        <table>
          <tr>
            <td>
              <label for="fname">First name</label>
            </td>
            <td>
              <input name="fname" id="fname" type="text">
            </td>
          </tr>
          <tr>
            <td>
              <label for="midname">Middle name</label>
            </td>
            <td>
              <input name="midname" id="midname"
type="text">
            </td>
          </tr>
          <tr>
            <td>
              <label for="lname">last name</label>
            </td>
            <td>
              <input name="lname" id="lname" type="text">
            </td>
          </tr>
          <tr>
            <td>
            </td>
          </tr>
        </table>
      </form>
    </center>
  </body>
</html>
```

```

        <td>
            <label for="pwd">Password</label>
        </td>
        <td>
            <input name="pwd" id="pwd" type="password"
                placeholder="*****">
        </td>
    </tr>
    <tr>
        <td>
            <label for="rpwd">Re enter Password</label>
        </td>
        <td>
            <input name="rpwd" id="rpwd" type="password"
                placeholder="*****">
        </td>
    </tr>
    <tr>
        <td>
            <label for="addr">Address</label>
        </td>
        <td>
            <textarea name="addr" id="addr" cols="22"
                rows="5"></textarea>
        </td>
    </tr>
    <tr>
        <td>
            <label for="gen">Gender</label>
        </td>
        <td>
            <input id="male" name="gender" type="radio"
                value="m"><label
for="male">Male</label>
            <input id="female" name="gender" type="radio"
                value="f"><label
for="female">Female</label>
        </td>
    </tr>
    <tr>
        <td>
            <label for="addr">Qualifications</label>
        </td>

```

```

        <td>
            <input id="btech" type="checkbox"
            value="btech"><label for="btech">B
Tech</label>
            <input id="mtech" type="checkbox"
            value="mtech"><label for="mtech">M Tech</label>
            <input id="phd" type="checkbox"
            value="phd"><label for="phd">PhD</label>
        </td>
    </tr>
    <tr>
        <td>
            <label for="dob">Date of birth</label>
        </td>
        <td>
            <input name="dob" id="dob" type="date">
        </td>
    </tr>
    <tr>
        <td>
            <label for="photo">Photo</label>
        </td>
        <td>
            <input id="photo" type="file">
        </td>
    </tr>
    <tr>
        <td>
            <label for="">Post applied for</label>
        </td>
        <td>
            <select>
                <option value="assist">Assistant
professor</option>
                <option value="asso">Associate
professor</option>
                <option value="prof">professor</option>
            </select>
        </td>
    </tr>
    <tr>
        <td>
            <label for="email">Email</label>

```

```

</td>
<td>
    <input id="email" name="email" type="email">
</td>
</tr>
<tr>
<td colspan="2">
    <center><button
type="submit">Submit</button></center>
</td>
</tr>
</table>
</form>
</center>
</body>
</html>

```

Output :

Registration form

First name	
Middle name	
last name	
Password	
Re enter Password	
Address	
Gender	☐ Male ☐ Female
Qualifications	☐ B Tech ☐ M Tech ☐ PhD
Date of birth	dd-mm-yyyy 📅
Photo	Choose File No file chosen
Post applied for	Assistant professor ▼
Email	
Submit	

PROGRAM 6

Create a HTML file by applying the different styles using inline, external & internal style sheets.

INLINE

CODE:

```
<html>
  <head>
    <style>
      .style1
      {
        background-color:PaleGreen
      }
      .style2
      {
        color:SeaGreen
      }
      .style3
      {
        color:DarkGreen
      }
      .style4
      {
        color:Green;
        padding: 20px;
      }
    </style>
  </head>
  <body class="style1">

    <h1 style="color:Teal;text-align:center;">Fish</h1>
    <p class="style2">Fish are gill-bearing aquatic craniate animals that lack
limbs with digits.
    They form a sister group to the tunicates, together forming the olfactores.
    Included in this definition are the living hagfish, lampreys, and cartilaginous
and bony fish as well as various extinct related groups. Around 99% of living
fish
    species are ray-finned fish, belonging to the class Actinopterygii, with over
95%
    belonging to the teleost subgrouping.</p>
    <p class="style3">The earliest organisms that can be classified as fish were
soft-bodied chordates
```

that first appeared during the Cambrian period. Although they lacked a true spine,

they possessed notochords which allowed them to be more agile than their invertebrate

counterparts. Fish would continue to evolve through the Paleozoic era, diversifying into

a wide variety of forms. Many fish of the Paleozoic developed external armor that protected

them from predators. The first fish with jaws appeared in the Silurian period, after which many (such as sharks) became formidable marine predators rather than just the prey of arthropods. Most fish are ectothermic ("cold-blooded"), allowing their body temperatures to vary as ambient temperatures change, though some of the large active swimmers like white shark and tuna can hold a higher core temperature.

Fish can communicate in their underwater environments through the use of acoustic communication. Acoustic communication in fish involves the transmission of acoustic signals from one individual of a species to another. The production of sounds as a means of communication among fish is most often used in the context of feeding, aggression or courtship behaviour. The sounds emitted by fish can vary depending on the species and stimulus involved. They can produce either stridulatory sounds by moving component of the skeletal system, or can produce non-stridulatory sounds by manipulating specialized organs such as the swim bladder. Fish are abundant in most bodies of water. They can be found in nearly all aquatic environments, from high mountain streams (e.g., char and gudgeon) to the abyssal and even hadal depths of the deepest oceans (e.g., cusk-eels and snailfish), although no species has yet been documented in the deepest 25% of the ocean. With 34,300 described species, fish exhibit greater species diversity than any other group of vertebrates.

Fish are an important resource for humans worldwide, especially as food. Commercial and subsistence fishers hunt fish in wild fisheries or farm them in ponds or in cages in the ocean (in aquaculture). They are also caught by recreational fishers, kept as pets, raised by fishkeepers, and exhibited in public aquaria. Fish have had a role in culture through the ages, serving as deities, religious symbols, and as the subject of art, books and movies. Tetrapods emerged within lobe-finned fishes, so cladistically they are fish as well. However, traditionally fish are rendered paraphyletic by excluding the tetrapods (i.e., the amphibians, reptiles, birds and mammals which all descended from within the same ancestry). Because in this manner the term "fish" is defined negatively as a paraphyletic group, it is not considered a formal taxonomic grouping in systematic biology, unless it is used in the cladistic sense, including tetrapods. The traditional term pisces (also ichthyoes) is considered a typological, but not a phylogenetic classification.

<marquee>

</marquee>

</body>

</html>

OUTPUT:


Fish

Fish are gill-bearing aquatic craniate animals that lack limbs with digits. They form a sister group to the tunicates, together forming the outgroup. Included in this definition are the living hagfish, lampreys, and cartilaginous and bony fish as well as various extinct related groups. Around 99% of living fish species are ray-finned fish, belonging to the class Actinopterygii, with over 95% belonging to the teleost subgrouping.

The earliest organisms that can be classified as fish were soft-bodied chordates that first appeared during the Cambrian period. Although they lacked a true spine, they possessed notochords which allowed them to be more agile than their invertebrate counterparts. Fish would continue to evolve through the Paleozoic era, diversifying into a wide variety of forms. Many fish of the Paleozoic developed external armor that protected them from predators. The first fish with jaws appeared in the Silurian period, after which many (such as sharks) became formidable marine predators rather than just the prey of arthropods. Most fish are ectothermic ("cold-blooded"), allowing their body temperatures to vary as ambient temperatures change, though some of the large active swimmers like white shark and tuna can hold a higher core temperature.

Fish can communicate in their underwater environments through the use of acoustic communication. Acoustic communication in fish involves the transmission of acoustic signals from one individual of a species to another. The production of sounds as a means of communication among fish is most often used in the context of feeding, aggression or courtship behaviour. The sounds emitted by fish can vary depending on the species and stimulus involved. They can produce either stridulatory sounds by moving component of the skeletal system, or can produce non-stridulatory sounds by manipulating specialized organs such as the swimbladder. Fish are abundant in most bodies of water. They can be found in nearly all aquatic environments, from high mountain streams (e.g., char and gudgeon) to the abyssal and even hadal depths of the deepest oceans (e.g., cusk-eels and snailfish), although no species has yet been documented in the deepest 25% of the ocean. With 34,300 described species, fish exhibit greater species diversity than any other group of vertebrates.

Fish are an important resource for humans worldwide, especially as food. Commercial and subsistence fisheries hunt fish in wild fisheries or farm them in ponds or in cages in the ocean (in aquaculture). They are also caught by recreational fishers, kept as pets, raised by fishkeepers, and exhibited in public aquaria. Fish have had a role in culture through the ages, serving as deities, religious symbols, and as the subjects of art, books and movies. Tetrapods emerged within lobe-finned fishes, so cladistically they are fish as well. However, traditionally fish are rendered paraphyletic by excluding the tetrapods (i.e., the amphibians, reptiles, birds and mammals which all descended from within the same ancestry). Because in this manner the term "fish" is defined negatively as a paraphyletic group, it is not considered a formal taxonomic grouping in systematic biology, unless it is used in the cladistic sense, including tetrapods. The traditional term pisces (also ichthyes) is considered a typological, but not a phylogenetic classification.



INTERNAL

CODE:

```
<html>
  <head>
    <style>
      body
      {
        background-color: Linen;
      }

      h1
      {
        text-align : center;
        color: maroon;
        margin-left: 40px;
      }
      p
      {
        color : red;
      }
    </style>
  </head>
  <body>

    <h1>Fish</h1>
    <p>Fish are gill-bearing aquatic craniate animals that lack limbs with digits.
    They form a sister group to the tunicates, together forming the outgroup.
```


Included in this definition are the living hagfish, lampreys, and cartilaginous and bony fish as well as various extinct related groups. Around 99% of living fish species are ray-finned fish, belonging to the class Actinopterygii, with over 95% belonging to the teleost subgrouping.

The earliest organisms that can be classified as fish were soft-bodied chordates that first appeared during the Cambrian period. Although they lacked a true spine, they possessed notochords which allowed them to be more agile than their invertebrate counterparts. Fish would continue to evolve through the Paleozoic era, diversifying into a wide variety of forms. Many fish of the Paleozoic developed external armor that protected them from predators. The first fish with jaws appeared in the Silurian period, after which many (such as sharks) became formidable marine predators rather than just the prey of arthropods.

Most fish are ectothermic ("cold-blooded"), allowing their body temperatures to vary as ambient temperatures change, though some of the large active swimmers like white shark and tuna can hold a higher core temperature. Fish can communicate in their underwater environments through the use of acoustic communication. Acoustic communication in fish involves the transmission of acoustic signals from one individual of a species to another. The production of sounds as a means of communication among fish is most often used in the context of feeding, aggression or courtship behaviour. The sounds emitted by fish can vary depending on the species and stimulus involved. They can produce either stridulatory sounds by moving component of the skeletal system, or can produce non-stridulatory sounds by manipulating specialized organs such as the swimbladder.

Fish are abundant in most bodies of water. They can be found in nearly all aquatic environments, from high mountain streams (e.g., char and gudgeon) to the abyssal and even hadal depths of the deepest oceans (e.g., cusk-eels and snailfish), although no species has yet been documented in the deepest 25% of the ocean. With 34,300 described species, fish exhibit greater species diversity than any other group of vertebrates. Fish are an important resource for humans worldwide, especially as food. Commercial and subsistence fishers hunt fish in wild fisheries or farm them in ponds or in cages in the ocean (in aquaculture). They are also caught by recreational fishers, kept as pets, raised by fishkeepers, and exhibited in public aquaria.

Fish have had a role in culture through the ages, serving as deities, religious symbols, and as the subjects of art, books and movies. Tetrapods emerged within lobe-finned fishes, so cladistically they are fish as well.

However, traditionally fish are rendered paraphyletic by excluding the tetrapods (i.e., the amphibians, reptiles, birds and mammals which all descended from within the same ancestry). Because in this manner the term "fish" is

defined negatively as a paraphyletic group, it is not considered a formal taxonomic grouping in systematic biology, unless it is used in the cladistic sense, including tetrapods. The traditional term pisces (also ichthyes) is considered a typological, but not a phylogenetic classification.







</body>

</html>

OUTPUT:

Fish

Fish are gill-bearing aquatic craniate animals that lack limbs with digits. They form a sister group to the tunicates, together forming the outgroup. Included in this definition are the living hagfish, lampreys, and cartilaginous and bony fish as well as various extinct related groups. Around 99% of living fish species are ray-finned fish, belonging to the class Actinopterygii, with over 95% belonging to the teleost subgrouping.

The earliest organisms that can be classified as fish were soft-bodied chordates that first appeared during the Cambrian period. Although they lacked a true spine, they possessed notochords which allowed them to be more agile than their invertebrate counterparts. Fish would continue to evolve through the Paleozoic era, diversifying into a wide variety of forms. Many fish of the Paleozoic developed external armor that protected them from predators. The first fish with jaws appeared in the Silurian period, after which many (such as sharks) became formidable marine predators rather than just the prey of arthropods.

Most fish are ectothermic ("cold-blooded"), allowing their body temperatures to vary as ambient temperatures change, though some of the large active swimmers like white shark and tuna can hold a higher core temperature. Fish can communicate in their underwater environments through the use of acoustic communication. Acoustic communication in fish involves the transmission of acoustic signals from one individual of a species to another. The production of sounds as a means of communication among fish is most often used in the context of feeding, aggression or courtship behaviour. The sounds emitted by fish can vary depending on the species and stimulus involved. They can produce either stridulatory sounds by moving component of the skeletal system, or can produce non-stridulatory sounds by manipulating specialized organs such as the swimbladder. Fish are abundant in most bodies of water. They can be found in nearly all aquatic environments, from high mountain streams (e.g., char and gudgeon) to the abyssal and even hadal depths of the deepest oceans (e.g., cusk-eels and snailfish), although no species has yet been documented in the deepest 25% of the ocean. With 34,300 described species, fish exhibit greater species diversity than any other group of vertebrates. Fish are an important resource for humans worldwide, especially as food. Commercial and subsistence fishers hunt fish in wild fisheries or farm them in ponds or in cages in the ocean (in aquaculture). They are also caught by recreational fishers, kept as pets, raised by fishkeepers, and exhibited in public aquaria. Fish have had a role in culture through the ages, serving as deities, religious symbols, and as the subjects of art, books and movies. Tetrapods emerged within lobe-finned fishes, so cladistically they are fish as well. However, traditionally fish are rendered paraphyletic by excluding the tetrapods (i.e., the amphibians, reptiles, birds and mammals which all descended from within the same ancestry). Because in this manner the term "fish" is defined negatively as a paraphyletic group, it is not considered a formal taxonomic grouping in systematic biology, unless it is used in the cladistic sense, including tetrapods. The traditional term pisces (also ichthyes) is considered a typological, but not a phylogenetic classification.



EXTERNAL

CODE:

style1.css

```
body{
    background-color:Aqua;
}

h1
{
    text-align : center;
    color: Indigo;
    margin-left: 60px;
}
```

```

p
{
    color : Darkblue;
}

```

external.html

```

<html>
  <head>
    <link href="style1.css" rel="stylesheet" type="text/css">
  </head>
  <body>

    <h1>Fish</h1>
    <p>Fish are gill-bearing aquatic craniate animals that lack limbs with digits.
    They form a sister group to the tunicates, together forming the olfactores.
    Included in this definition are the living hagfish, lampreys, and cartilaginous
    and bony fish as well as various extinct related groups. Around 99% of living
fish
95%
species are ray-finned fish, belonging to the class Actinopterygii, with over
belonging to the teleost subgrouping.<br>
The earliest organisms that can be classified as fish were soft-bodied chordates
that first appeared during the Cambrian period. Although they lacked a true
spine,
they possessed notochords which allowed them to be more agile than their
invertebrate counterparts. Fish would continue to evolve through the Paleozoic
era, diversifying into a wide variety of forms. Many fish of the Paleozoic
developed external armor that protected them from predators. The first fish
with jaws appeared in the Silurian period, after which many (such as sharks)
became formidable marine predators rather than just the prey of arthropods.
Most fish are ectothermic ("cold-blooded"), allowing their body temperatures
to vary as ambient temperatures change, though some of the large active
swimmers like white shark and tuna can hold a higher core temperature.<br>
Fish can communicate in their underwater environments through the use of
acoustic communication. Acoustic communication in fish involves the
transmission of acoustic signals from one individual of a species to another.
The production of sounds as a means of communication among fish is most
often used
in the context of feeding, aggression or courtship behaviour. The sounds
emitted by fish can vary depending on the species and stimulus involved. They
can produce either stridulatory sounds by moving component
of the skeletal system, or can produce non-stridulatory sounds by
manipulating specialized organs such as the swimbladder.
Fish are abundant in most bodies of water. They can be found in nearly all
aquatic environments, from high mountain streams

```

(e.g., char and gudgeon) to the abyssal and even hadal depths of the deepest oceans (e.g., cusk-eels and snailfish), although no species has yet been documented in the deepest 25% of the ocean. With 34,300 described species, fish exhibit greater species diversity than any other group of vertebrates. Fish are an important resource for humans worldwide, especially as food. Commercial and subsistence fishers hunt fish in wild fisheries or farm them in ponds or in cages in the ocean (in aquaculture). They are also caught by recreational fishers, kept as pets, raised by fishkeepers, and exhibited in public aquaria. Fish have had a role in culture through the ages, serving as deities, religious symbols, and as the subjects of art, books and movies. Tetrapods emerged within lobe-finned fishes, so cladistically they are fish as well. However, traditionally fish are rendered paraphyletic by excluding the tetrapods (i.e., the amphibians, reptiles, birds and mammals which all descended from within the same ancestry). Because in this manner the term "fish" is defined negatively as a paraphyletic group, it is not considered a formal taxonomic grouping in systematic biology, unless it is used in the cladistic sense, including tetrapods. The traditional term pisces (also ichthyoes) is considered a typological, but not a phylogenetic classification.







OUTPUT:

Fish

Fish are gill-bearing aquatic craniate animals that lack limbs with digits. They form a sister group to the tunicates, together forming the outgroup. Included in this definition are the living hagfish, lampreys, and cartilaginous and bony fish as well as various extinct related groups. Around 99% of living fish species are ray-finned fish, belonging to the class Actinopterygii, with over 95% belonging to the teleost subgrouping.

The earliest organisms that can be classified as fish were soft-bodied chordates that first appeared during the Cambrian period. Although they lacked a true spine, they possessed notochords which allowed them to be more agile than their invertebrate counterparts. Fish would continue to evolve through the Paleozoic era, diversifying into a wide variety of forms. Many fish of the Paleozoic developed external armor that protected them from predators. The first fish with jaws appeared in the Silurian period, after which many (such as sharks) became formidable marine predators rather than just the prey of arthropods. Most fish are ectothermic ("cold-blooded"), allowing their body temperatures to vary as ambient temperatures change, though some of the large active swimmers like white shark and tuna can hold a higher core temperature.

Fish can communicate in their underwater environments through the use of acoustic communication. Acoustic communication in fish involves the transmission of acoustic signals from one individual of a species to another. The production of sounds as a means of communication among fish is most often used in the context of feeding, aggression or courtship behaviour. The sounds emitted by fish can vary depending on the species and stimulus involved. They can produce either stridulatory sounds by moving component of the skeletal system, or can produce non-stridulatory sounds by manipulating specialized organs such as the swimbladder. Fish are abundant in most bodies of water. They can be found in nearly all aquatic environments, from high mountain streams (e.g., char and gudgeon) to the abyssal and even hadal depths of the deepest oceans (e.g., cusk-eels and snailfish), although no species has yet been documented in the deepest 25% of the ocean. With 34,300 described species, fish exhibit greater species diversity than any other group of vertebrates.

Fish are an important resource for humans worldwide, especially as food. Commercial and subsistence fishers hunt fish in wild fisheries or farm them in ponds or in cages in the ocean (in aquaculture). They are also caught by recreational fishers, kept as pets, raised by fishkeepers, and exhibited in public aquaria. Fish have had a role in culture through the ages, serving as deities, religious symbols, and as the subjects of art, books and movies. Tetrapods emerged within lobe-finned fishes, so cladistically they are fish as well. However, traditionally fish are rendered paraphyletic by excluding the tetrapods (i.e., the amphibians, reptiles, birds and mammals which all descended from within the same ancestry). Because in this manner the term "fish" is defined negatively as a paraphyletic group, it is not considered a formal taxonomic grouping in systematic biology, unless it is used in the cladistic sense, including tetrapods. The traditional term pisces (also ichthyoes) is considered a typological, but not a phylogenetic classification.



PROGRAM 7

Create a HTML page to explain the use of various predefined functions in a string and math object in java script.

Code:

```
<html>
  <head>
    <title>Sting and math objects</title>
  </head>
  <body>
    <script>
      document.write("<h2>Predefined Math
Objects</h2>");
      document.write("Value of PI: ",Math.PI);
      document.write("<br>");
      document.write("Round value of 4.6:
",Math.round(4.6));
      document.write("<br>");
      document.write("2th power of 8: ",Math.pow(8,2));
      document.write("<br>");
      document.write("Square root of 25: ",Math.sqrt(25));
      document.write("<br>");
      document.write("Absolute value of -4.7: ",Math.abs(-
4.7));
      document.write("<br>");
      document.write("Ceil value of 4.4: ",Math.ceil(4.4));
      document.write("<br>");
      document.write("Floor value of 5.2: ",Math.floor(5.2));
      document.write("<br>");
      document.write("Value of sin 90: ",Math.sin(90));
      document.write("<br>");
      document.write("Value of cos 180: ",Math.cos(180));
      document.write("<br>");
      document.write("Numbers are 0,150,30,5,-10,-2");
      document.write("<br>");
      document.write("Minimum Value :
",Math.min(0,150,30,5,-10,-2));
      document.write("<br>");
```

```

document.write("Maximum Value:
",Math.max(0,150,30,5,-10,-2));
document.write("<br>");
document.write("A random value between 0 and 1:
",Math.random());
document.write("<br>");
document.write("<h2>Predefined String
Objects</h2>");
document.write("The string : Iam studying in Vidya
Academy of Science and Technology");
var str="Iam studying in Vidya Academy of Science
and Technology"
document.write("<br>");
document.write("Length of the string: ",str.length);
document.write("<br>");
document.write("Position of word Vidya:
",str.indexOf("Vidya"));
document.write("<br>");
document.write("The last index of letter T:
",str.lastIndexOf("T"));
document.write("<br>");
document.write("Searching word Science:
",str.search("Science"));
document.write("<br>");
document.write("Part of the string: ",str.slice(10,20));
document.write("<br>");
document.write("Substring of the string:
",str.substring(16,29));
document.write("<br>");
document.write("Replaceing Iam by We are: ",
str.replace("Iam","We are"));
document.write("<br>");
document.write("Text to lower case:
",str.toLowerCase());
document.write("<br>");
document.write("Text to Upper case:
",str.toUpperCase());
</script>
</body>
</html>

```

Output:

Predefined Math Objects

Value of PI: 3.141592653589793
Round value of 4.6: 5
2th power of 8: 64
Square root of 25: 5
Absolute value of -4.7: 4.7
Ceil value of 4.4: 5
Floor value of 5.2: 5
Value of sin 90: 0.8939966636005579
Value of cos 180: -0.5984600690578581
Numbers are 0,150,30,5,-10,-2
Minimum Value : -10
Maximum Value: 150
A random value between 0 and 1: 0.7963664836211608

Predefined String Objects

The string : Iam studying in Vidya Academy of Science and Technology
Length of the string: 55
Position of word Vidya: 16
The last index of letter T: 45
Searching word Science: 33
Part of the string: ng in Vidy
Substring of the string: Vidya Academy
Replaceing Iam by We are: We are studying in Vidya Academy of Science and Technology
Text to lower case: iam studying in vidya academy of science and technology
Text to Upper case: IAM STUDYING IN VIDYA ACADEMY OF SCIENCE AND TECHNOLOGY

PROGRAM 8

Create a HTML registration form and to validate the form using JavaScript code.

Source Code:

```
<html>
  <head>
    <title>
      Registration form
    </title>
  </head>
  <body>
    <script language="javascript">
      function check()
      {
        var fname=document.form1.fname.value;
        var mname=document.form1.midname.value;
        var lname=document.form1.lname.value;
        var pass=document.form1.pwd.value;
        var conpass=document.form1.rpwd.value;
        var address=document.form1.addr.value;
        var email=document.form1.email.value;
        if(fname==null || fname=="")
          alert("First name cannot be empty.");
        else if(lname==null || lname=="")
          alert("Lastname name cannot be empty.");
        else if(pass==null || pass=="")
          alert("Password cannot be empty.");
        else if(pass.length<8)
          alert("Password should contain atleast 8
characters");
        else if(conpass==null || conpass=="")
          alert("Password should be confirmed.");
        else if(pass!=conpass)
          alert("Password doesnot match!!!.");
        else if(address==null || address=="")
          alert("Address cannot be empty.");

        else if(email==null || email=="")
```



```

        alert("Email cannot be empty.");
    else
        alert("Registration Successfull");
    }
</script>
<center>
    <h1>Registration form</h1>
    <form name="form1">
        <table width="30%">
            <tr>
                <td>
                    First name
                </td>
                <td>
                    <input name="fname" id="fname" type="text">
                </td>
            </tr>
            <tr>
                <td>
                    Middle name
                </td>
                <td>
                    <input name="midname" id="midname" type="text">
                </td>
            </tr>
            <tr>
                <td>
                    last name
                </td>
                <td>
                    <input name="lname" id="lname" type="text">
                </td>
            </tr>
            <tr>
                <td>
                    Password
                </td>
                <td>
                    <input name="pwd" id="pwd" type="password"
placeholder="*****">

```

```

        </td>
    </tr>
    <tr>
        <td>
            Re enter Password
        </td>
        <td>
            <input name="rpwd" id="rpwd" type="password"
placeholder="*****">
        </td>
    </tr>
    <tr>
        <td>
            Address
        </td>
        <td>
            <textarea name="addr" id="addr" cols="22"
rows="5"></textarea>
        </td>
    </tr>
    <tr>
        <td>
            Email
        </td>
        <td>
            <input id="email" name="email" type="email">
        </td>
    </tr><br>
    <tr>
        <td colspan="2">

<center><br>&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;&nbsp;<input
type="button" name="Submit" value="Register" onclick="check()">
            <input type="reset" name="clear"
value="Reset"></center>
        </td>
    </tr>
</table>
</form>

```

```
</center>
</body>
</html>
```

Output:

This page says

Password should contain atleast 8 charcters

OK

First name	Catherine
Middle name	
last name	Joshi
Password	*****
Re enter Password	*****
Address	Tharakan House Adat
Email	catherine271099

Register Reset

Password doesnot match!!!.

OK

First name	Catherine
Middle name	
last name	Joshi
Password	*****
Re enter Password	*****
Address	Tharakan House Adat
Email	catherine271099

Register Reset

Lastname name cannot be empty.

OK

First name	Catherine
Middle name	
last name	
Password	*****
Re enter Password	*****
Address	Tharakan House Adat
Email	catherine271099

Register Reset

Registration Successfull

OK

First name Catherine

Middle name

last name Joshi

Password

Re enter Password

Address Tharakan House
Adat

Email catherine271099

Register

Reset

PROGRAM 9

Generate the calendar using JavaScript code by getting the year from the user.

Source Code:

```
<html>
  <head>
    <title>Calender</title>
  <script>
    function genCalender()
    {
      var year=prompt("Enter the year");
      if(year==null)
        year=2021;
      var month=prompt("Enter the month");
      if(month==null)
        month=1;
      var numDays=(new Date(year,month,0)).getDate();
      var wDay=(new Date(year,month-1,1)).getDay();
      document.body.innerHTML="
      document.write("<center>");
      document.write("<h1>Calender("+year+"-
"+month+"))");

      document.write("<table width='400px' border='1'
      align='center'>");
      document.write("<thead>");
      document.write("<tr bgcolor='cyan' align='center'>");
      document.write("<td width='50px'>Sun</td>");
      document.write("<td width='50px'>Mon</td>");
      document.write("<td width='50px'>Tue</td>");
      document.write("<td width='50px'>Wed</td>");
      document.write("<td width='50px'>Thu</td>");
      document.write("<td width='50px'>Fri</td>");
      document.write("<td width='50px'>Sat</td>");
      document.write("</tr>");
      document.write("</thead>");
      var i;
      document.write("<tr align='center'>");
      for(i=0;i<wDay;i++)
```

```

        {
            document.write("<td width='50px'></td>");
        }
        var d;
        for(d=1;d<=numDays;d++)
        {
            document.write("<td
width='50px'>" + d + "</td>");
            if(i==6)
            {
                document.write("</tr>");
                document.write("<tr align='center'>");
                i=0;
            }
            else
                i++;
        }
        for(;i<7;i++)
        {
            document.write("<td width='50px'></td>");
        }
        document.write("</table><br>");
        document.write("<button
onclick='genCalender()'>Read year and
Month</button>");
        document.write("</center>");
    }
</script>
</head>
<body>
    <center>
        <button onclick="genCalender()">Read Year
        and Month</button>
    </center>
</body>
</html>

```

OUTPUT:

Enter the year

OK Cancel

Enter the month

OK Cancel

Calender(2021-2)

Sun	Mon	Tue	Wed	Thu	Fri	Sat
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28						

Read year and Month

PROGRAM 10


Create a HTML page to change the background color when clicking of button of different colors using JavaScript Event Handling.

Source Code:

```
<html>
  <head>
    <title>ChangeColor</title>
    <script language="javascript">
      function changecolor(color)
      {
        document.body.style.backgroundColor=color;
      }

    </script>
  </head>
  <body>
    <center>
      <input type="button" onclick="changecolor('blue')"
value="blue">
      <input type="button" onclick="changecolor('cyan')"
value="cyan">
      <input type="button" onclick="changecolor('black')"
value="black">
      <input type="button" onclick="changecolor('violet')"
value="violet">
      <input type="button" onclick="changecolor('lightgreen')"
value="lightgreen">
      <input type="button" onclick="changecolor('orange')"
value="orange">
      <input type="button" onclick="changecolor('green')"
value="green">
    </center>
  </body>
</html>
```


Output:



PROGRAM 11

Create a HTML page to display a new image and text when the mouse comes over the existing content in the page using JavaScript Event Handling.

Source Code:

```
<html>
  <head>
    <title>changeimage</title>
    <script language="javascript">
      function setfirstimage()
      {
        document.getElementById("image").src="bird1.jpg";
      }
      function setnextimage()
      {
        document.getElementById("image").src="bird2.jpg";
      }
    </script>
  </head>
  <body>
    
  </body>
</html>
```

Output:



PROGRAM 12

Create a HTML page to show online quiz using JavaScript

Code:

```
<html>
  <head>
    <title>Online Quiz</title>
    <style>
      div
      {
        margin: auto;
        width : 50%
      }
    </style>
    <script>
      function getResults()
      {
        var marks=0;
        if(document.getElementById('q1a3').checked)
          marks=marks+1;
        if(document.getElementById('q2a3').checked)
          marks=marks+1;
        if(document.getElementById('q3a2').checked)
          marks=marks+1;
        if(document.getElementById('q4a2').checked)
          marks=marks+1;
        if(document.getElementById('q5a1').checked)
          marks=marks+1;
        alert("RESULT\nYour Score : "+marks);
      }
    </script>
  </head>
  <body>
    <div>
      <h2>Online Quiz on HTML</h2>
      1.HTML stands for:- <br>
      <input type="radio" id="q1a1" name="q1"
      value="q1a1">Hypertext Machine Language<br>
```

☐Hypertext and Links Markup Language
☐Hypertext Markup Language
☐None of these

2.The correct sequence of HTML tags for creating a webpage is:-

☐Head, Title, HTML, Body
☐HTML, Body, Title, Head
☐HTML, Head, Title, Body
☐HTML, Head, Body, Title

3.Which element makes the text bold:-

☐bold tag
☐strong tag
☐t tag
☐None of these

4.How to create an ordered list:-

☐ul tag
☐ol tag
☐li tag
☐i tag

5.Which element makes text italic :-

☐i tag
☐italic tag
☐pre tag

```
<input type="radio" id="q5a4" name="q5" value="q5a4">it  
tag<br><br>  
<center><input type="button" id="submit" value="submit"  
onclick="getResults()"></center>  
</div>  
</body>  
</html>
```

OUTPUT:

Online Quiz on HTML

1.HTML stands for:-

- ☐ Hypertext Machine Language
- ☐ Hypertext and Links Markup Language
- ☐ Hypertext Markup Language
- ☐ None of these

2.The correct sequence of HTML tags for creating a webpage is:-

- ☐ Head, Title, HTML, Body
- ☐ HTML, Body, Title, Head
- ☐ HTML, Head, Title, Body
- ☐ HTML, Head, Body, Title

3.Which element makes the text bold:-

- ☐ bold tag
- ☐ strong tag
- ☐ t tag
- ☐ None of these

4.How to create an ordered list:-

- ☐ ul tag
- ☐ ol tag
- ☐ li tag
- ☐ i tag

5.Which element makes text italic :-

- ☐ i tag
- ☐ italic tag
- ☐ pre tag
- ☐ it tag

submit

Online Quiz

RESULT

Your Score : 4

OK

1. HTML stands for

- ☐ Hypertext Machine
- ☐ Hypertext and Language
- ☒ Hypertext Markup Language
- ☐ None of these

2. The correct sequence of HTML tags for creating a webpage is:-

- ☐ Head, Title, HTML, Body
- ☐ HTML, Body, Title, Head
- ☒ HTML, Head, Title, Body
- ☐ HTML, Head, Body, Title

3. Which element makes the text bold:-

- ☐ bold tag
- ☐ strong tag
- ☒ b tag
- ☐ None of these

4. How to create an ordered list:-

- ☐ ul tag
- ☒ ol tag
- ☐ li tag
- ☐ i tag

5. Which element makes text italic :-

- ☒ i tag
- ☐ italic tag
- ☐ pre tag
- ☐ it tag

submit