

Chapter 1: Part II: Asymptotic Notations

Devesh C Jinwala , IIT Jammu, India

June 9, 2021

Design and Analysis of Algorithms
IIT Jammu, Jammu

1 Asymptotic Notations

The Big-Oh Notation

- def: for a given function $g(n)$, we say that $O(g(n)) = f(n)$ — if there exists positive constants c and n_0 such that, $0 \leq f(n) \leq cg(n)$, for all $n \geq n_0$

$$f(n) = O(g(n)) \Rightarrow$$

$f(n)$ is dominated in the growth by $g(n)$ i.e. $f(n)$ is of the order at the most $g(n)$ i.e. $g(n)$ grows at least as fast as $f(n)$

The Big-Oh Notation

- def: for a given function $g(n)$, we say that $O(g(n)) = f(n)$ — if there exists positive constants c and n_0 such that,
 $0 \leq f(n) \leq cg(n)$, for all $n \geq n_0$
- The Big-oh defines an upper bound for a function within a constant factor i.e. except for a constant factor and a finite number of exceptions, f is bounded above by g .

$$f(n) = O(g(n)) \Rightarrow$$

$f(n)$ is dominated in the growth by $g(n)$ i.e. $f(n)$ is of the order at the most $g(n)$ i.e. $g(n)$ grows at least as fast as $f(n)$


The Big-Oh Notation

- Can $f(n)$ grow faster than $g(n)$?

The Big-Oh Notation

$$f(n) = o(g(n))$$


- Can $f(n)$ grow faster than $g(n)$?
- Can $g(n)$ grow faster than $f(n)$?



The Big-Oh Notation

- Can $f(n)$ grow faster than $g(n)$?
- Can $g(n)$ grow faster than $f(n)$?

$\lim_{n \rightarrow \infty}$



- What does the growth rate imply ?

The Big-Oh Notation Illustrations

Function	notation in O
$f(n) = 5n + 8$	$f(n) = O(?)$
$f(n) = \underline{n^2} + 3n - 8$	$f(n) = O(?)$
$F(n) = 12n^2 - 11$	$f(n) = O(?)$
$F(n) = 5 \cdot 2^n + n^2$	$f(n) = O(?)$
$f(n) = 3n + 8$	$F(n) = O(n^2)?$
$f(n) = 5n + 8$	$f(n) = O(1)?$

 $O(1)$ $O(n)$ $O(n^2)$ $O(n^3)$ 2^n γ

The Big-Oh Notation

$$f(n) = 5n + 8$$

p.t. $f(n) = \underline{\underline{O(n)}}$

- allows us to keep track of the leading term while ignoring smaller terms

$$f(n) = 5n + 8$$

$$\leq 5n + 8n$$

$$= 13n$$

$C, n \geq n_0$

$\therefore c = 13$ and $n_0 = 1$ we have
 $f(n) \leq \cancel{O(n)}^{c \cdot n} \therefore f(n) = O(n)$

The Big-Oh Notation

- allows us to keep track of the leading term while ignoring smaller terms
- allows us to make concise statements that give approximations to the quantities to analyze.

The Big-Oh Notation

- allows us to keep track of the leading term while ignoring smaller terms

The Big-Oh Notation

- allows us to keep track of the leading term while ignoring smaller terms
- allows us to make concise statements that give approximations to the quantities to analyze.

Tutorial Problem No 1


- if $f(n) = O(g(n))$, what is the upper bound ?

Tutorial Problem No 1

- if $f(n) = O(g(n))$, what is the upper bound ?
- Do we specify how tight this upper bound is?

Tutorial Problem No 1

- if $f(n) = O(g(n))$, what is the upper bound ?
- Do we specify how tight this upper bound is?
- Consider that $f(n) = O(n)$ & $g(n) = O(n^2)$. Is $f(n) = O(g(n))$ saying the same as reverse i.e. $g(n) = O(f(n))$?

Tutorial Problem No 1

- if $f(n) = O(g(n))$, what is the upper bound ?
- Do we specify how tight this upper bound is?
- Consider that $f(n) = O(n)$ & $g(n) = O(n^2)$. Is $f(n) = O(g(n))$ saying the same as reverse i.e. $g(n) = O(f(n))$?
- The symbol $=$ is not proper truly it is \in which should be used i.e. $f(n) \in O(g(n))$ \in

The Big-Oh notation...

- When O notation bounds the worst case running time of an algorithm, by implication we also bound the running time of an algorithm on EVERY input.

$$O(n^2) \rightarrow n$$

Abuse

Technically, it is abuse to say that the running time of insertion sort is $O(n^2)$. Why?

Wrong

The Big-Oh notation...

- When O notation bounds the worst case running time of an algorithm, by implication we also bound the running time of an algorithm on EVERY input.
- this is not so when using other notations i.e. the worst case $\theta(n^2)$ or $\theta(n)$ does not apply to every input.

Abuse

Technically, it is abuse to say that the running time of insertion sort is $O(n^2)$. Why?

The Big- Ω notation...

- the big- Ω notation is used to specify this aspect i.e. to define a lower bound for a function within a constant factor..

The Big- Ω notation...

- the big- Ω notation is used to specify this aspect i.e. to define a lower bound for a function within a constant factor..
- definition: the function $f(n) = \Omega(g(n))$ is true iff there exists positive constants c and n_0 such that $f(n) \geq c(g(n))$ for all $n - n \geq n_0$ i.e. $0 \leq c(g(n)) \leq f(n)$.

The Big- Ω notation...

- the big- Ω notation is used to specify this aspect i.e. to define a lower bound for a function within a constant factor..
- definition: the function $f(n) = \Omega(g(n))$ is true iff there exists positive constants c and n_0 such that $f(n) \geq c(g(n))$ for all $n - n \geq n_0$ i.e. $0 \leq c(g(n)) \leq f(n)$.
- i.e. except for a constant factor and a finite number of exceptions, f is bounded below by g .

The Big- Ω notation...

- the big- Ω notation is used to specify this aspect i.e. to define a lower bound for a function within a constant factor..
- definition: the function $f(n) = \Omega(g(n))$ is true iff there exists positive constants c and n_0 such that $f(n) \geq c(g(n))$ for all $n - n \geq n_0$ i.e. $0 \leq c(g(n)) \leq f(n)$.
- i.e. except for a constant factor and a finite number of exceptions, f is bounded below by g .
- $f(n) = \Omega(g(n))$ implies that

The Big- Ω notation...

- the big- Ω notation is used to specify this aspect i.e. to define a lower bound for a function within a constant factor..
- definition: the function $f(n) = \Omega(g(n))$ is true iff there exists positive constants c and n_0 such that $f(n) \geq c(g(n))$ for all $n - n \geq n_0$ i.e. $0 \leq c(g(n)) \leq f(n)$.
- i.e. except for a constant factor and a finite number of exceptions, f is bounded below by g .
- $f(n) = \Omega(g(n))$ implies that

$\Omega(n) \Rightarrow$


$f(n)$ always dominates the growth of $g(n)$ i.e. $f(n)$ is of the order at least $g(n)$ i.e. $g(n)$ grows at the most as fast as $f(n)$.

The Big- Ω notation...

The Big Omega

- Can $f(n)$ grow faster than $g(n)$?
- Can $g(n)$ grow faster than $f(n)$?

} if $f(n) \neq o(g(n))$



The Big-Oh Notation Illustrations

Function	notation in Ω
$f(n) = 3n + 8$	$f(n) = \Omega(?)$ n
$f(n) = n^2 + 3n - 8$	$f(n) = \Omega(?)$ n^2
$F(n) = 12n^2 - 11$	$f(n) = \Omega(?)$ n^2
$F(n) = 6 \cdot 2^n + n^2$	$f(n) = \Omega(?)$ 2^n
$f(n) = 3n + 8$	$f(n) = \Omega(n^2)?$
$f(n) = 5n + 8$	$f(n) = \Omega(1)?$ <i>yes</i>

Tutorial Problem No 2

- Is it correct to say that $3n + 8 = \Omega(1)$?

Tutorial Problem No 2

- Is it correct to say that $3n + 8 = \Omega(1)$?
- There could be several functions $g(n)$ for which $f(n) = \Omega(g(n))$ - the function $g(n)$ is only lower bound on n .

Tutorial Problem No 2

- Is it correct to say that $3n + 8 = \Omega(1)$?
- There could be several functions $g(n)$ for which $f(n) = \Omega(g(n))$ - the function $g(n)$ is only lower bound on n .
- hence, for $f(n) = \Omega(g(n))$ to be meaningful, $g(n)$ should be as large a function of n as possible.

Tutorial Problem No 2

- Is it correct to say that $3n + 8 = \Omega(1)$?
- There could be several functions $g(n)$ for which $f(n) = \Omega(g(n))$ - the function $g(n)$ is only lower bound on n .
- hence, for $f(n) = \Omega(g(n))$ to be meaningful, $g(n)$ should be as large a function of n as possible.
- Given two choices viz. $3n+3 = \Omega(n)$ and $3n+3 = \Omega(1)$, which one shall we choose ?

Tutorial Problem No 2

- Is it correct to say that $3n + 8 = \Omega(1)$?
- There could be several functions $g(n)$ for which $f(n) = \Omega(g(n))$ - the function $g(n)$ is only lower bound on n .
- hence, for $f(n) = \Omega(g(n))$ to be meaningful, $g(n)$ should be as large a function of n as possible.
- Given two choices viz. $3n+3 = \Omega(n)$ and $3n+3 = \Omega(1)$, which one shall we choose ?
- if $f(n) = O(\Omega(n))$, what is the lower bound ?

Tutorial Problem No 2

- Is it correct to say that $3n + 8 = \Omega(1)$?
- There could be several functions $g(n)$ for which $f(n) = \Omega(g(n))$ - the function $g(n)$ is only lower bound on n .
- hence, for $f(n) = \Omega(g(n))$ to be meaningful, $g(n)$ should be as large a function of n as possible.
- Given two choices viz. $3n+3 = \Omega(n)$ and $3n+3 = \Omega(1)$, which one shall we choose ?
- if $f(n) = O(\Omega(n))$, what is the lower bound ?
- Do we specify how tight this lower bound is?

Tutorial Problem No 2...

- When we say $f(n) = \Omega(g(n))$, does it mean that $g(n) = O(f(n))$?

$$\underline{f(n) = \Omega(g(n))} \quad \xrightarrow{\quad} \quad \underline{g(n) = O(f(n))}$$

$f(n) \geq c \cdot g(n)$ $g(n) = O(f(n))$

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?
- Consider $f(n)=\underline{n^2}$ & $g(n)=\underline{2n^2}$

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?
- Consider $f(n)=n^2$ & $g(n)=2n^2$... $f(n) = O(g(n))$

When we use Ω -notation to describe a lower bound,

$$f(n) = \Omega(g(n))$$

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?
- Consider $f(n)=n^2$ & $g(n)=2n^2$

When we use Ω -notation to describe a lower bound,

- we are also implying a lower bound on the running time of the algorithm on arbitrary inputs as well.

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?
- Consider $f(n)=n^2$ & $g(n)=2n^2$

When we use Ω -notation to describe a lower bound,

- we are also implying a lower bound on the running time of the algorithm on arbitrary inputs as well.
- the bounds $O(n^2)$ and $\Omega(n)$ are as tight bounds as possible.

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?
- Consider $f(n)=n^2$ & $g(n)=2n^2$

When we use Ω -notation to describe a lower bound,

- we are also implying a lower bound on the running time of the algorithm on arbitrary inputs as well.
- the bounds $O(n^2)$ and $\Omega(n)$ are as tight bounds as possible.
- Can we say that the running time of insertion sort is $\Omega(n^2)$? //

Tutorial Problem No 2...

- When we say $f(n)=\Omega(g(n))$, does it mean that $g(n)=O(f(n))$?
- Would the growth of $f(n)$ dominate the growth of $(g(n))$?
- Consider $f(n)=n^2$ & $g(n)=2n^2$

When we use Ω -notation to describe a lower bound,

- we are also implying a lower bound on the running time of the algorithm on arbitrary inputs as well.
- the bounds $O(n^2)$ and $\Omega(n)$ are as tight bounds as possible.
- Can we say that the running time of insertion sort is $\Omega(n^2)$?
- Can we say that the running time of insertion sort is $O(n)$?

The Big- Θ notation...

- Neither the big-O notation nor the big- Ω notation describe the asymptotically tight bounds.
- Θ -notation to express tighter bounds - used to specify the exact order of growth of functions.
- def: we say that $f(n) = \Theta(g(n))$ iff there exists positive constants c_1 and c_2 and a number n_0 such that $0 \leq c_1 g(n) \leq f(n) \leq c_2 g(n)$ for all $n \geq n_0$
- $f(n) = \Theta(g(n))$ iff $f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$

The Big- Θ notation...

- Neither the big- O notation nor the big- Ω notation describe the asymptotically tight bounds.
- Θ -notation to express tighter bounds - used to specify the exact order of growth of functions.
- def: we say that $f(n)=\Theta(g(n))$ iff there exists positive constants c_1 and c_2 and a number n_0 such that $0 \leq c_1g(n) \leq f(n) \leq c_2g(n)$ for all $n \geq n_0$
- $f(n)=\Theta(g(n))$ iff $f(n)=O(g(n))$ and $f(n)=\Omega(g(n))$

The Big- Θ Notation...

Given $f(n) = \Theta(g(n))$

$\rightarrow f(n) = O(g(n))$

- Can $f(n)$ grow faster than $g(n)$?

$=$


$f(n) = \Omega(g(n))$

\rightarrow NO

The Big- Θ Notation...

Given $f(n) = \Theta(g(n))$


- Can $f(n)$ grow faster than $g(n)$?
- Can $g(n)$ grow faster than $f(n)$?



The Big- Θ Notation...

Given $f(n) = \Theta(g(n))$

- Can $f(n)$ grow faster than $g(n)$?
- Can $g(n)$ grow faster than $f(n)$?




- What does the growth rate imply ?

The Big- Θ Notation...

Function	notation in θ
$f(n) = 3n + 8$	$f(n) = \theta(?)$
$f(n) = 10n^2 + 3n - 8$	$f(n) = \theta(?)$
$F(n) = 12n^2 - 11$	$f(n) = \theta(?)$
$F(n) = 6 \cdot 2^n + n^2$	$f(n) = \theta(2^n) ?$
$F(n) = 6 \cdot 2^n + n^2$	$f(n) = \theta(n^2) ?$
$f(n) = 3n + 8$	$f(n) = \theta(n^2) ?$
$f(n) = 5n + 8$	$f(n) = \theta(1) ?$

The Asymptotic Classes



Complexity of Data Structures

Ref: Eric- <http://bigocheatsheet.com>

Data Structure	Time Complexity								Space Complexity
	Average				Worst				Worst
	Access	Search	Insertion	Deletion	Access	Search	Insertion	Deletion	
Array	$O(1)$	$O(n)$	$O(n)$	$O(n)$	$O(1)$	$O(n)$	$O(n)$	$O(n)$	$O(n)$
Stack	$O(n)$	$O(n)$	$O(1)$	$O(1)$	$O(n)$	$O(n)$	$O(1)$	$O(1)$	$O(n)$
Singly-Linked List	$O(n)$	$O(n)$	$O(1)$	$O(1)$	$O(n)$	$O(n)$	$O(1)$	$O(1)$	$O(n)$
Doubly-Linked List	$O(n)$	$O(n)$	$O(1)$	$O(1)$	$O(n)$	$O(n)$	$O(1)$	$O(1)$	$O(n)$
Skip List	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(n)$	$O(n)$	$O(n)$	$O(n)$	$O(n \log(n))$
Hash Table	-	$O(1)$	$O(1)$	$O(1)$	-	$O(n)$	$O(n)$	$O(n)$	$O(n)$
Binary Search Tree	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(n)$	$O(n)$	$O(n)$	$O(n)$	$O(n)$
Cartesian Tree	-	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	-	$O(n)$	$O(n)$	$O(n)$	$O(n)$
B-Tree	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(n)$
Red-Black Tree	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(n)$
Splay Tree	-	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	-	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(n)$
AVL Tree	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(\log(n))$	$O(n)$

Complexities of Sorting Algorithms


linear time sorting + type of sorting algorithms - m.

Algorithm	Time Complexity			Space Complexity
	Best	Average	Worst	Worst
<u>Quicksort</u>	$O(n \log(n))$	$O(n \log(n))$	$O(n^2)$	$O(\log(n))$
<u>Mergesort</u>	$O(n \log(n))$	$O(n \log(n))$	$O(n \log(n))$	$O(n)$
<u>Timsort</u>	$O(n)$	$O(n \log(n))$	$O(n \log(n))$	$O(n)$
<u>Heapsort</u>	$O(n \log(n))$	$O(n \log(n))$	$O(n \log(n))$	$O(1)$
<u>Bubble Sort</u>	$O(n)$	$O(n^2)$	$O(n^2)$	$O(1)$
<u>Insertion Sort</u>	$O(n)$	$O(n^2)$	$O(n^2)$	$O(1)$
<u>Selection Sort</u>	$O(n^2)$	$O(n^2)$	$O(n^2)$	$O(1)$
<u>Shell Sort</u>	$O(n)$	$O((n \log(n))^2)$	$O((n \log(n))^2)$	$O(1)$
<u>Bucket Sort</u>	$O(n+k)$	$O(n+k)$	$O(n^2)$	$O(n)$
<u>Radix Sort</u>	$O(nk)$	$O(nk)$	$O(nk)$	$O(n+k)$

Ref: Eric- <http://bigcheatsheet.com/>

Lower Bound on Sorting

- To sort n elements, one needs information about the sequence of these elements
i.e. given two elements, the relative order could be



Lower Bound on Sorting

- To sort n elements, one needs information about the sequence of these elements
i.e. given two elements, the relative order could be
- Decision tree at a higher level of abstraction

Lower Bound on Sorting

- To sort n elements, one needs information about the sequence of these elements
i.e. given two elements, the relative order could be
- Decision tree at a higher level of abstraction
- For an input sequence of three elements, what would be the total number of 2-element comparisons ?

Lower Bound on Sorting

- To sort n elements, one needs information about the sequence of these elements
i.e. given two elements, the relative order could be
- Decision tree at a higher level of abstraction
- For an input sequence of three elements, what would be the total number of 2-element comparisons ?
- Draw a decision tree showing the number of comparisons of any n -distinct elements - say for $n = 3$.

Lower Bound on Sorting

- To sort n elements, one needs information about the sequence of these elements
i.e. given two elements, the relative order could be
- Decision tree at a higher level of abstraction
- For an input sequence of three elements, what would be the total number of 2-element comparisons ?
- Draw a decision tree showing the number of comparisons of any n -distinct elements - say for $n = 3$.
 - represent each internal node by $a_i:a_j$ in the range $1 \leq i \leq n$

Lower Bound on Sorting

- To sort n elements, one needs information about the sequence of these elements

i.e. given two elements, the relative order could be

- Decision tree at a higher level of abstraction

permutation

- For an input sequence of three elements, what would be the total number of 2-element comparisons ?

$$\frac{n!}{(n-1)!}$$


$$\frac{3!}{(3-2)!}$$

$$= 6$$

- Draw a decision tree showing the number of comparisons of any n -distinct elements - say for $n = 3$.

- represent each internal node by $a_i : a_j$ in the range $1 \leq i \leq n$

- denote each leaf by permutation $(\pi(1), \pi(2), \pi(3), \pi(4) \dots \pi(n))$

Blank $a_1 a_2 a_3$: Decision - tree

Blank

Theorem: Any decision tree that sorts n elements has height $\boxed{\Omega(n \lg n)}$

Proof:

→ Consider a decision tree of height h that sorts n elements.

→ How many leaves does this tree have?

→ If the height of a tree is h , then
 (a) maximally it can have 2^h leaves.
 $n! \leq 2^h$ i.e. $\boxed{2^h}$ ①

→ (b) if a decision tree represent sorting of n elements, then it will have $n!$ leaves ②

Blank

$$n! \leq [2^n]$$

$$\text{i.e. } \log n! \leq \log [2^n]$$

$$h \gg n \log n. \\ = \Omega(n \log n)$$

$$\log n! \leq h \quad \dots (i)$$

\Rightarrow Stirling's approximation applied to our case:

$$\log n \leftarrow n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n (1 + \Theta(1/n))$$

$$\boxed{\log e} \quad n! \geq \left(\frac{n}{e}\right)^n \quad \dots (ii)$$

$$h > \log \left(\frac{n^n}{e^n}\right) \quad \left(h = \underline{n \log n} - \boxed{n \log e} \right)$$

Blank

Corollary: Heapsort and Mergesort are optimal comparison based sorts.

Proof: Optimal? \rightarrow the best we can do.

① Any comparison $\Omega(n \log n)$

\downarrow at least $n \log n$

② complexities of Hs and Ms

$O(n \log n)$ || ③ $\Omega(n^2)$

Blank

Counting sort

→ not comparison based
 $O(n)$

Self-reading