Minimizing exotic matter in rotating Ellis–Bronnikov wormholes

Nikita M. Shirokov[1, *] and Ilya Nachevsky[2, †]

[1] *Moscow Institute of Physics and Technology, Moscow, Russia*
[2] *Higher School of Economics, Moscow, Russia*

(Dated: October 31, 2026)

We ask whether a scalar potential can reduce the exotic matter needed to support a rotating Ellis–Bronnikov wormhole while keeping its throat size and angular momentum fixed. An automated search varies the potential and solves the stationary field equations for each trial. Additional Einstein constraints check the resulting geometry. For a selected slowly rotating drainhole, a calculation on a finer grid retains a 61% reduction in volume-integrated null-energy violation relative to the massless scalar solution. Comparisons of three selected drainholes show that the scalar variation becomes concentrated near the throat: the smaller volume integral accompanies stronger peak violation, stronger peak tides and longer finite-distance light transit. Three-dimensional evolutions initially show decreasing constraint errors as the grid is refined. The throat later contracts, with onset moving later at higher resolution. The potential therefore lowers the stationary matter requirement for this solution, while its subsequent evolution remains sensitive to numerical resolution.

I. INTRODUCTION

A traversable wormhole must violate the null energy condition (NEC) near its throat [3, 4]. The Ellis–Bronnikov solution provides a simple field model: a massless scalar with a reversed kinetic sign supports the throat [1, 2]. This phantom scalar supplies the required exotic matter, but the static wormhole is radially unstable [5, 6, 15, 16].

How much exotic matter is needed depends on both its distribution and the geometry. Visser, Kar, and Dadhich showed that the integrated violation can be made arbitrarily small in idealized thin-shell constructions [7, 8]. Our interest is in smooth scalar-field solutions that can also serve as initial data for nonlinear evolution. Rotation already reduces their exotic-matter integral along the families of Kleihaus and Kunz [11, 12, 16].

Those rotating solutions use a massless scalar without a potential. We instead allow a potential that changes how the scalar varies between the two ends of the wormhole. This can alter both the distribution of exotic matter and the geometry it supports. To isolate that effect, we compare solutions at the same equatorial throat radius and angular momentum.

Ordinary bosonic matter provides another way to modify the throat geometry. Dzhunushaliev et al. [10] constructed boson stars containing phantom-supported wormholes. Their solutions retain a radial unstable mode, although suitable boson self-interactions can bring its eigenvalue close to zero.

Here we vary the phantom potential, solve for the scalar and metric together, and measure the integrated NEC violation. The optimizer uses these measurements to choose further trials (Fig. 1). Independent Einstein constraints test whether each numerical solution is consistent with equations that were not used to construct it. Promising candidates are then checked on a finer grid before a selected solution is evolved in three dimensions.

Section II defines the model and the matter measure. Sections III and IV explain the search and evolution methods. Section V presents the search and evolution results, and Sec. VI examines the matter distribution and physical trade-offs, with background, analytical and evolution checks in Appendices A–C. We use units in which Newton's constant and the speed of light satisfy $G = c = 1$.

II. THE MATTER MODEL

We consider Einstein gravity coupled to the classical Ellis–Bronnikov phantom scalar ψ, augmented by a potential $\mathcal{V}(\psi)$,

$$S = \frac{1}{16\pi}\int d^4x\,\sqrt{-g}\,(R + 2\,g^{\mu\nu}\partial_\mu\psi\,\partial_\nu\psi + 2\,\mathcal{V}(\psi))\,, \quad (1)$$

where g is the determinant of the spacetime metric and R is its Ricci scalar. This extends the massless model of Refs. [11, 12, 16]. The reversed kinetic sign permits NEC violation.

We use metric signature $(-,+,+,+)$. Defining the matter stress tensor by $T_{\mu\nu} = -2(-g)^{-1/2}\delta S_{\rm m}/\delta g^{\mu\nu}$ gives

$$8\pi T_{\mu\nu} = -2\partial_\mu\psi\partial_\nu\psi + g_{\mu\nu}\big[(\partial\psi)^2 + \mathcal{V}\big], \quad (2)$$

where $(\partial\psi)^2 = g^{\mu\nu}\partial_\mu\psi\partial_\nu\psi$. Its trace is $8\pi T = 2(\partial\psi)^2 + 4\mathcal{V}$, and the trace of $G_{\mu\nu} = 8\pi T_{\mu\nu}$ therefore gives

$$R = -2(\partial\psi)^2 - 4\mathcal{V}. \quad (3)$$

Subtracting half the stress-tensor trace from Einstein's equation, and integrating the scalar variation by parts, yields

$$R_{\mu\nu} = -2\,\partial_\mu\psi\,\partial_\nu\psi - \mathcal{V}\,g_{\mu\nu}, \qquad \Box\psi = \tfrac{1}{2}\mathcal{V}'(\psi). \quad (4)$$

* shirokov.nm@phystech.edu
† ommnammashivaya@gmail.com

Here $R_{\mu\nu}$ is the Ricci tensor, $\Box$ is the covariant wave operator, and a prime denotes a derivative with respect to ψ. Without a potential, the scalar has the analytic arctan profile of Ref. [12]. With a potential, both the scalar and the metric must respond. Since $\partial_t\psi = \partial_\varphi\psi = 0$, the mixed Einstein equations imply

$$R^t{}_t + R^\varphi{}_\varphi = -2\mathcal{V}. \tag{5}$$

The massless choice $B = 1$ solves the corresponding homogeneous metric equation; retaining it for nonzero $\mathcal{V}$ would discard this source. We therefore solve for an additional metric function B:

$$\begin{aligned} ds^2 = -e^f dt^2 + B^2 e^{-f}\Big[e^\nu\left(d\eta^2 + h\, d\theta^2\right) \\ + h\sin^2\theta\left(d\varphi - \omega\, dt\right)^2\Big], \end{aligned} \tag{6}$$

with $h = \eta^2 + \eta_0^2$. The coordinates η and θ describe the radial and polar directions; η_0 is the throat parameter. The metric functions f, ω, ν and B, together with ψ, depend on both coordinates. The massless solutions have $B = 1$. We solve for the field value $\psi_- = \psi(-\infty)$ along with these functions.

The two families differ in the relation between their asymptotic ends. We set $f \to 0$ at $\eta \to +\infty$ and $f \to \gamma$ at $\eta \to -\infty$. The symmetric family has $\gamma = 0$, whereas the drainhole family has $\gamma = -\pi$ [16]. For the latter we use $\eta_0 = 2$, so its static massless member has unit mass.

The asymptotic data for the five fields are

$$\begin{aligned} (f, \omega, \nu, B, \psi)_{+\infty} &= (0, 0, 0, 1, 0), \\ (f, \omega, \nu, B, \psi)_{-\infty} &= (\gamma, \omega_-, 0, 1, \psi_-). \end{aligned} \tag{7}$$

The constants ω_- and ψ_- are solved for; ω_- specifies the relative rotation of the asymptotic frames. At the negative end, $t_- = e^{\gamma/2}t$ and $\varphi_- = \varphi - \omega_- t$ give the local nonrotating time and azimuth. Axis regularity requires $\nu = 0$ and $\partial_\theta(f, \omega, B, \psi) = 0$ at $\theta = 0, \pi$, with $\nu = O(\sin^2\theta)$. Equatorial reflection symmetry requires $\partial_\theta(f, \omega, \nu, B, \psi) = 0$ at $\theta = \pi/2$. The even angular basis enforces these reflection conditions.

The potential must vanish at both ends so that they remain asymptotically flat. Its derivative must also vanish at the limiting field values. We choose a double-well form with vacua at $\psi = 0$ and $\psi = \psi_-$:

$$\mathcal{V}(\psi) = \frac{m^2}{\psi_-^2}\,\psi^2(\psi - \psi_-)^2\, Q(u), \qquad u = \frac{\psi}{\psi_-}, \tag{8}$$

where $Q(u) = 1 + q_1 u + q_2 u^2 + q_3 u^3$. The prefactor fixes the two vacua, while the polynomial changes the barrier between them.

The search varies (m, q_1, q_2, q_3). The parameter m controls the curvature of the potential near the first vacuum, $\mathcal{V}''(0) = 2m^2$. The coefficients q_i change the height and asymmetry of the barrier. We require $Q(u) > 0$ between the vacua; individual coefficients may be negative. Setting $m = 0$ recovers the massless model.

The potential also fixes the asymptotic decay. Write $\delta\psi_\pm = \psi - \psi_\pm$, with $\psi_+ = 0$. Linearization in the locally inertial frame at each end gives

$$(\Box - \mu_\pm^2)\delta\psi_\pm = 0, \qquad \mu_+^2 = m^2, \quad \mu_-^2 = m^2 Q(1). \tag{9}$$

Thus the two ends generally have different decay lengths. In the leading flat-space approximation, the decaying stationary branch behaves as $\delta\psi_\pm \propto e^{-\mu_\pm\rho_\pm}/\rho_\pm$, where $\rho_+ \simeq \eta$ and $\rho_- \simeq e^{-\gamma/2}|\eta|$ are asymptotic proper radii. Positive $Q(1)$ also gives positive mass squared for local vacuum perturbations. This asymptotic statement does not determine the stability of the coupled wormhole. Replacing the massless algebraic tails by exponential decay explains how a potential can localize the scalar variation; the spectral grid must still resolve that variation.

To compare the matter content of different solutions, we integrate the radial NEC violation and normalize it by the equatorial throat radius R_e [7, 16]:

$$X = \frac{1}{R_e}\left|\int(\rho + p_\eta)\, dV\right|. \tag{10}$$

Here ρ is the energy density and p_η the radial pressure, measured by a local zero-angular-momentum observer (ZAMO), whose four-velocity and radial unit vector are

$$n = e^{-f/2}(\partial_t + \omega\partial_\varphi), \qquad e_{\hat\eta} = B^{-1}e^{(f-\nu)/2}\partial_\eta. \tag{11}$$

The scalar is stationary in this frame. Contracting Eq. (2) with the locally unit-energy radial null vector $n + e_{\hat\eta}$ gives

$$\begin{aligned} \rho + p_\eta &= -\frac{(\nabla_{\hat\eta}\psi)^2}{4\pi}, \\ dV &= B^3 e^{-3f/2+\nu} h\sin\theta\, d\eta\, d\theta\, d\varphi. \end{aligned} \tag{12}$$

This remains a radial NEC diagnostic when the scalar has angular gradients. We call X the exoticity. The potential cancels directly from $\rho + p_\eta$, but it changes the scalar gradient and the volume element through the coupled field equations. Solving those equations is therefore essential to finding whether a trial potential lowers X.

We hold R_e and the angular momentum J at the values of a chosen massless reference solution. This makes the comparison about the potential's effect, rather than a change in throat size or spin.

III. OPTIMIZATION

The search alternates between constructing a wormhole and choosing a new potential. For each trial, a spectral Newton solver adjusts the scalar and geometry until the stationary field equations are satisfied numerically. The covariance-matrix adaptation evolution strategy (CMA-ES) [18] ranks the trials and uses them to guide the next proposals. Figure 1 shows the checks and measurements linking these steps.




FIG. 1. Optimization of the confining potential at fixed throat size and angular momentum. Each CMA-ES proposal starts Newton iteration from a nearby solved candidate; JAX supplies the Jacobian for the GPU solve. Exoticity and background-constraint penalties feed back into the search; inadmissible or failed solves receive a penalty. Dashed arrows lead to separate resolution and evolution tests. The loop ranks candidates; the subsequent tests assess their reliability.

A. Spectral Newton solver

The spectral representation covers both asymptotic ends on a finite computational domain. We use $x = (2/\pi)\arctan(\eta/\eta_0)$ to map them to $x = \pm 1$, and expand the fields in Chebyshev polynomials radially and even cosine modes angularly. A tensor-product Lobatto grid supplies the points at which the equations are evaluated. Larger vacuum masses shorten the physical decay lengths. Under the algebraic compactification, exponential radial decay need not give exponential decay of Chebyshev coefficients; we monitor the spectral tail directly.

Newton iteration computes a correction to the fields from the residual equations and their Jacobian. Damping reduces the correction when a full step would not improve the solution. We solve for f, ω, $\nu/\sin^2\theta$, B, and ψ; dividing out $\sin^2\theta$ regularizes the metric variable on the symmetry axis. The vacuum value ψ_- and a radial length scale are additional unknowns, with two equations enforcing the chosen R_e and J.

The Jacobian describes how the residual changes when the unknowns change. We compute it using automatic differentiation in `JAX` [19], then solve the resulting dense linear system on a graphics processing unit (GPU). Two independent Einstein constraints check the converged fields. We also inspect the highest spectral coefficients: a slowly decaying tail signals structure that the grid may not resolve. Appendix A gives the background checks.

B. CMA-ES

CMA-ES draws a population of trial potentials from a distribution in parameter space. After each population is evaluated, the better candidates guide the location and spread of subsequent proposals. For each trial, Newton iteration starts from the nearest previously solved potential, which supplies an initial guess for the fields.

The score rewards low exoticity subject to numerical admissibility. Let $C = \max(|E_{19}|, |E_c|)$ be the larger independent background residual, using the dimensionless residuals of Appendix A. The bounded search uses

$$S_{\mathrm{bg}} = \begin{cases} X, & C \le 10^{-8}, \\ X + 0.1, & 10^{-8} < C \le 10^{-6}, \\ 10^3 + \log_{10}(C/10^{-6}), & C > 10^{-6}, \\ 1010, & \text{invalid trial.} \end{cases} \tag{13}$$

An invalid trial violates the parameter bounds, fails Newton convergence, has nonfinite diagnostics or has a nonpositive sampled Q: we evaluate it at 1001 uniformly spaced values of $u \in [0, 1]$. The bounded drainhole and S0 searches use $0.15 \le m \le 1.2$ and $-2 \le q_i \le 8$, in the length units of their reference solutions. CMA-ES starts with parameter scales $(0.1, 0.3, 0.3, 0.3)$. The earlier S27, S50 and S85 searches used the same mass interval without box bounds on q_i; their scoring is specified in Appendix A.

For the resolution-screened searches, a proposed new minimum is also checked on finer grids. A failed check receives score 1001; a passing one is ranked by its finer-grid X. The finalist is passed to the independent background validation. Stationary admissibility defines the reported sample; dynamical conclusions come from the evolutions, not from a radial-stability penalty.

The searches explore different regions of potential space for the two wormhole families (Fig. 2). Labels D and S denote drainhole and symmetric reference solutions; the suffix gives $100v_e$, where v_e is the reference equatorial rotation velocity. Each search keeps its own reference throat size and angular momentum fixed.

The parameter projections show which combinations were sampled, rather than an interpolated surface of solutions. Points can overlap even when the other two potential parameters differ. The progress curves divide each search's best value by its first candidate below the constraint ceiling, so their slopes show improvement within a search.




FIG. 2. CMA-ES exploration of the four potential parameters at fixed R_e and J, for the drainhole (top) and symmetric (bottom) families. Panels (a,b,d,e) project recorded proposals, coloured by evaluation order; diamonds mark each search's lowest X with both background-constraint residuals below 10^{-6}. The q_3 axes are symmetric-logarithmic, linear for $|q_3| \leq 1$. Panels (c,f) show the running minimum relative to the first candidate below that threshold. The searches concentrate in distinct parameter regions and approach plateaus at different rates.

IV. NUMERICAL SETUP

A. Interpolation onto the grid

The stationary solution supplies the initial metric and scalar field for evolution on a Cartesian grid with adaptive mesh refinement (AMR). We interpolate the background table using cubic splines with continuous second derivatives. The isotropic radius r is related to the stationary coordinate by $\eta = r - \eta_0^2/(4r)$, so the second asymptotic end is represented by a puncture at $r = 0$ [16].

The additional metric function B enters the conformal factor through $\chi \to \chi/B^2$. The lapse α, shift β^i, and scalar ψ are read from the stationary data. The extrinsic curvature K_{ij} follows from the stationary shift, and the scalar's initial conjugate momentum vanishes in the corotating frame. The exported data also contain the potential coefficients needed during evolution.

The puncture represents an asymptotic end, rather than a material center. Initially, $B \to 1$, $\nu \to 0$, $\alpha \to e^{\gamma/2}$ and

$$\chi \sim \frac{16 e^{\gamma} r^4}{\eta_0^4}, \qquad \ln|\psi - \psi_-| \sim -\frac{c_-}{r}, \qquad c_- = \frac{\mu_- e^{-\gamma/2}\eta_0^2}{4}. \tag{14}$$

For a massive vacuum, the scalar's spatial derivatives therefore vanish at the puncture in the continuum initial data. The massless case instead has $\psi - \psi_- = O(r)$. We evolve conformal variables on a cell-centered grid, with a positive floor on χ; gauge evolution subsequently changes the lapse and conformal factor. Fixed nested refinement boxes are centered on $r = 0$ and cover the throat as well. They do not refine only a shell at $r = \eta_0/2$, which denotes $\eta = 0$ and need not be the physical throat. The convergence runs use a χ floor of 10^{-8} and exclude the puncture region $r < 0.125$ from the constraint norms.

B. Evolution scheme

We evolve these data with `GRTeclyn` [23–25], following the formulation used in Refs. [15, 16]. The metric evolution uses the conformal and covariant Z4 (CCZ4) equations [20], with 1+log slicing [21] and the Gamma-driver shift [22]. Spatial derivatives and Runge–Kutta time integration are fourth order, with Kreiss–Oliger dissipation.

The scalar evolution includes the potential and its derivative. The Hamiltonian constraint diagnostic includes the potential contribution as well. Mesh refinement concentrates resolution near the throat. The op-

tional outer sponge term that drives fields toward their reference values is disabled. Appendix C specifies the grid and timestep used for the resolution study.

C. Criteria

A low score identifies a promising trial, but the reported stationary candidates must also satisfy the independent background equations. We distinguish the loose constraint ceiling used during exploration from the stricter threshold used for Tables I and II. The former is 10^{-6}; the latter is 10^{-8} for each of the two residuals. Trials between these thresholds receive a constraint penalty.

The tables select the lowest recorded X meeting the stricter threshold in each search. The selected finer-grid backgrounds are listed separately from the search minima. Perturbative stability requires an additional test: a converged mode must satisfy the independent linearized Einstein constraints as well as its eigenvalue equations.

V. RESULTS

A. Stationary exoticity at fixed throat and spin

The search finds lower exoticity at most of the sampled rotations in both families. Tables I and II give the potential parameters and the corresponding matter integrals, using the same model order. Their search-grid rows describe the minima found at that numerical resolution. Figure 3 places these candidates relative to the massless families.

The reduction is not uniform across spin. The high-spin D90 candidate remains above its massless counterpart and lies close to the imposed lower mass bound $m = 0.15$. The search excludes the exactly massless point $m = 0$, even though the analytic potential family includes it. This is a minimum found in the restricted search, not evidence that every nonzero potential raises exoticity at this spin.

For the selected D21* solution, the finer-grid calculation retains a reduction of about 61% in exoticity at the same throat radius and angular momentum. Its background residuals satisfy the independent constraints (Appendix A). This is the solution used for the three-dimensional evolution study; it is distinct from the lowest point in the search history.

B. Three-dimensional evolution and contraction

We evolved D21* on three grids to examine how the dynamics change with resolution. The domain and refinement hierarchy were held fixed while the grid spacing decreased. Early in the evolution, the constraint errors near the throat fall approximately at fourth order. Other




FIG. 3. Exoticity relative to each family's massless static member, against the reference rotation v_e. D and S denote drainhole and symmetric families. Black curves show the massless solutions; blue markers show selected potential-bearing solutions. Filled markers identify the five 210 × 42 backgrounds; open markers show search-grid candidates. Curves are shape-preserving interpolants through the sampled values, used as visual guides within the sampled ranges. Most selected solutions require less exotic matter than their massless counterparts.

refinement levels show more variable behavior, as quantified in Appendix C.

The throat later contracts on all three grids, with contraction starting later as resolution increases. Figure 7 shows this shift in the radius histories alongside the wave and lapse diagnostics. A later onset is consistent with a smaller truncation-error perturbation growing exponentially. Appendix C quantifies this interpretation using the logarithmic dependence expected for such a perturbation. The constraint order and the order of the perturbation amplitude are different measurements.

At fixed numerical resolution, the potential-bearing D0, D21*, D30 and D40 runs reach the same fractional contraction later than their classical counterparts (Fig. 4). We measure the time at which each throat's areal radius first falls by 1% from its own initial value. The difference between these times measures a delay in contraction, separately from the subsequent exponential growth rate. D0, D30 and D40 use their reference throat size, angular momentum and evolution settings. The D21 comparison also changes the background spin slightly and the outer damping: classical D21 uses $J = 1$ and drive strength 1, while D21* uses $J = 1.0138$ and drive strength 0. Holding area fixed, the classical quadratic rate fit predicts only a 0.50% change from this spin shift. This estimates local spin sensitivity; it does not isolate the sponge effect. We therefore interpret the D21 delay as a comparison of these two numerical setups rather than a controlled potential-only effect.

The growth rate answers a separate question: how rapidly does contraction amplify once it develops? We fit $\ln|dR_A/dt|$ over the same 0.2–1% areal-radius decrease

TABLE I. Potential parameters of the lowest-X candidates with both background-constraint residuals at most 10^{-8} on the 150×30 search grid. D and S denote the drainhole and symmetric families. D21* is the separately selected solution re-solved on 210×42, rather than the search-grid minimum.

model	m	q_1	q_2	q_3
Drainhole, 150×30 search grid				
D0	1.199	0.227	0.485	0.274
D21	1.171	0.762	0.351	0.206
D30	0.740	0.607	0.201	−0.664
D80	0.234	0.083	0.504	1.187
D90	0.150	−0.337	0.366	0.445
Symmetric, 150×30 search grid				
S0	1.198	2.433	−0.046	1.197
S27	0.855	3.232	3.776	−0.256
S50	0.962	3.739	0.922	−0.828
S85	0.425	2.438	4.267	0.445
Selected 210×42 solution				
D21*	1.023	0.672	0.048	0.310

TABLE II. Exoticity of the same candidates as Table I, compared with the massless solution at the same R_e and J. Here $\Delta X = 100(X/X_{\mathcal{V}=0} - 1)$. The search-grid rows and the selected finer-grid solution are listed separately.

model	X	$X_{\mathcal{V}=0}$	ΔX
Drainhole, 150×30 search grid			
D0	1.839	4.898	−62.5%
D21	1.785	4.645	−61.6%
D30	1.880	4.376	−57.0%
D80	1.105	1.419	−22.2%
D90	0.787	0.600	+31.3%
Symmetric, 150×30 search grid			
S0	1.627	3.142	−48.2%
S27	1.634	2.994	−45.4%
S50	1.477	2.625	−43.7%
S85	1.098	1.462	−24.9%
Selected 210×42 solution			
D21*	1.829	4.645	−60.6%




FIG. 4. Time to the first 1% decrease in throat areal radius at $N = 128$, for classical (black circles) and optimized (open blue squares) runs. Crossings are interpolated between saved samples; lines join measured points, and grey segments show the delay. Horizontal positions use the classical reference spins. D21* has the spin and damping differences described in the text. Every optimized run reaches this contraction threshold later.




FIG. 5. Measured growth rates at $N = 128$ for the classical models and four potential-bearing solutions (open blue squares). Positions retain $8\pi J/A$; ticks name the classical references. Grey and blue curves fit $a - b(8\pi J/A)^2$ to contracting models; the dashed blue extension is an extrapolation, not a measured stability threshold. Over the common 0.2–1% contraction interval, the potential lowers the rate for D0–D30 but raises it at D40.

(Fig. 5). For D21*, the growth time increases from $\tau = 6.59$ to 9.49. D0 and D30 likewise have lower fitted rates. D40 instead has a 20% higher rate despite its delayed contraction. Thus the potential delays the observed contraction in these runs, but does not uniformly suppress its growth rate. The resolution dependence of the onset distinguishes these finite-resolution delays from a continuum instability measurement.

VI. PHYSICS OF THE OPTIMIZED SOLUTIONS

The potential lowers the volume-integrated exoticity by concentrating the scalar variation into a narrower region near the throat. It does not weaken the NEC violation everywhere. We examine this redistribution in three selected 210×42 drainholes, denoted D0*, D21* and D30*. These are finer-grid selections, distinct from the search minima in Tables I and II. Table III lists their potential parameters and residuals. Selection requires the background constraints at the target size and spin on the finer grids; a smaller search-grid X alone does

not establish that accuracy. D0* is a rescaled alternative candidate, D21* is a separate selection from its original search, and D30* comes from a fresh search with finer-grid screening of each new minimum. Each has its own massless reference at the same R_e, J and asymmetry. In particular, both members of the stationary D21 comparison have $J = 1.0138$.

A. Matter distribution and throat geometry

Figure 6 compares profiles against physical proper distance, rather than the radial coordinates of different numerical solutions. The local radial NEC deficit is the squared radial orthonormal scalar gradient $(\nabla_{\hat{\eta}}\psi)^2$ divided by 4π. Here the radial null vector has unit energy in the local zero-angular-momentum observer's frame. The potential cancels from its direct null contraction, but changes the scalar and metric through the field equations.

In D21*, the scalar excursion between the two ends falls from 3.46 to 1.25. Its remaining variation occurs over a much shorter distance, so the peak radial NEC violation is 3.4 times stronger. The proper width between the 5% and 95% cumulative contributions to X decreases from $27.16R_e$ to $0.65R_e$. This measure includes the long classical tails beyond the plotted interval; it describes the matter distribution, not a change in throat radius. Angular scalar gradients remain small, giving a narrow radial layer rather than an equatorial ring.

The same pattern occurs in the other two solutions (Table IV). Their narrower distributions outweigh the higher peaks, reducing X by 56–62% across the three models. The equatorial throat radius is fixed, and the throat area changes by at most 0.6%. The rotating throats become slightly less oblate. Thus the reduction mainly reflects a changed scalar distribution, while the overall throat dimensions remain similar.

A simple variational bound helps separate localization from a reduced field excursion. In a spherical radial model, let $A(\ell)$ be the area of a proper-distance sphere. Cauchy–Schwarz gives

$$\int A(\ell)\left(\frac{d\psi}{d\ell}\right)^2 d\ell \geq \frac{(\Delta\psi)^2}{\int A(\ell)^{-1}d\ell}. \tag{15}$$

At fixed geometry and excursion, narrowing the scalar transition cannot lower this integral without limit. Here the excursion itself decreases, and the geometry responds. The support also moves toward smaller-area surfaces, removing the large-area weighting of the classical tails. These changes allow a lower volume integral together with a higher peak.

Equality in Eq. (15) requires $A\,d\psi/d\ell$ to be constant. A static massless scalar instead obeys $\partial_\ell(\alpha A\,d\psi/d\ell) = 0$. The bound is therefore saturated by the symmetric ultrastatic Ellis solution, but not generally by a redshifted drainhole. For the rotating solutions it supplies radial intuition; the measurements use the full axisymmetric volume element in Eq. (12).

B. Tides, redshift and light propagation

A smaller volume integral does not imply a gentler passage through the throat. We measure tides by the largest absolute eigenvalue of the electric Riemann tensor in a radially moving observer's frame. At the same local speed $v = 0.5$, its peak over the matched interval increases in all three solutions. This compares local forces along a fixed-speed scan; a freely falling traveller's accumulated tidal exposure also depends on its trajectory. The stronger peak is consistent with the sharper scalar and metric variation visible in Fig. 6.

The asymptotic redshift remains fixed by the common asymmetry γ, while the interior profile changes. For a zero-angular-momentum photon, the local energy is $E_{\rm local}/E_{+\infty} = 1/\alpha$, where $\alpha = e^{f/2}$ is the lapse and $E_{+\infty}$ is the conserved energy normalized at positive infinity. The optimized profile makes the transition between the two ends more abrupt.

We compute a finite light-crossing time along the equatorial null geodesic by integrating $dt/d\ell = 1/\alpha$ between endpoints at equal proper distances from each throat. The same positive-infinity clock normalizes t in both solutions. Transit is longer for the optimized solutions (Table IV); the increase also occurs when the endpoint distances are halved or doubled, although the ratio changes. These are wormhole-to-wormhole comparisons at specified endpoints.

The magnitude of the NEC integral along the complete equatorial null ray also increases. Its affine parameter is normalized by unit Killing energy at positive infinity, unlike the local normalization used for the peak NEC deficit. The ray and volume integrals therefore measure different aspects of the same distribution. Minimizing X reduces the volume-weighted matter requirement while concentrating the violation encountered near the throat.

VII. CONCLUSIONS

Adding a scalar potential can lower the stationary exotic-matter requirement without changing the reference throat size or angular momentum. The search combines numerical solutions of the field equations with independent Einstein constraints, and it retains a reduction of about 61% for the selected D21 solution on a finer grid. The benefit depends on the reference solution; the high-spin D90 search does not improve on its massless counterpart.

The D21 evolutions initially show decreasing constraint errors under refinement and later develop contraction. Its onset moves later on finer grids. At a common numerical resolution, the four evolved potential-bearing models

TABLE III. Selected 210×42 backgrounds used for the stationary physics comparison. Parameters are in the same convention as Table I; residuals test the two unused Einstein equations. All three meet the 10^{-8} background threshold at their matched reference size and spin.

model	m	q_1	q_2	q_3	X	$\lvert E_{19}\rvert$	$\lvert E_c\rvert$
D0*	1.248	0.341	0.291	0.173	1.842	1.1×10^{-9}	2.5×10^{-9}
D21*	1.023	0.672	0.048	0.310	1.829	1.6×10^{-10}	4.3×10^{-10}
D30*	0.585	0.956	0.226	−0.012	1.919	4.2×10^{-9}	3.8×10^{-9}




FIG. 6. Stationary profiles of D21* and its massless reference at the same R_e and J. The signed equatorial proper distance ℓ is zero at the minimum-area throat. Panels show the equatorial scalar gradient, the angularly integrated contribution to X, the area radius of constant-η surfaces, and the energy of a zero-angular-momentum photon measured by a local observer relative to its conserved energy at positive infinity. The potential concentrates the scalar gradient and exoticity near the throat while sharpening the interior redshift profile.

reach a fixed fractional contraction later than their classical counterparts. This delay is distinct from the fitted growth rate: D40 contracts later even though its subsequent rate is higher.

The stationary comparisons explain the exoticity reduction: the scalar variation is confined to a narrower region near the throat, with little change in the throat's overall dimensions. The local violation becomes stronger, and the selected solutions have larger peak tides and longer finite-distance light transit. Reducing the volume-integrated matter requirement therefore carries a trade-off in these measures of passage through the wormhole.

VIII. REPRODUCIBILITY

The solver, optimization and figure-generation scripts are available in the `GRTeclyn` research repository [26]. The archived search traces identify every candidate used in Tables I and II; a checked selection ledger generates both tables. A separate checked ledger supplies Table IV from the saved D0*, D21* and D30* fields. Those fields, their massless references, physical profiles and numerical quadrature checks are archived alongside the final evolution diagnostic streams. All article figures and tables rebuild from those tracked data.

TABLE IV. Stationary quantities for each massless reference ($V = 0$) and its optimized counterpart at the same R_e, J and asymmetry. Lengths are normalized by the common R_e within each pair; the displayed quantities are dimensionless. Optimized columns use the selected finer-grid solutions D0*, D21* and D30*. The width includes the complete radial distribution; the null NEC integral follows the complete equatorial ray with unit Killing energy at positive infinity. Peak tides and light transit use equatorial proper distances $-10R_e$ to $+10R_e$. A smaller volume integral accompanies stronger local violation, stronger peak tides and longer transit.

	D0		D21		D30	
Quantity	$V = 0$	Optimized	$V = 0$	Optimized	$V = 0$	Optimized
Total exoticity X	4.898	1.842	4.645	1.829	4.376	1.919
Central-90% proper width $/R_e$	28.83	0.59	27.16	0.65	25.40	0.96
Peak radial NEC violation $\times R_e^2$	0.0885	0.3318	0.0887	0.3041	0.0889	0.2189
Ray-integrated NEC magnitude $\times R_e$	0.304	0.431	0.311	0.432	0.320	0.425
Peak tides $\times R_e^2$, $v = 0.5$	0.528	3.469	0.521	3.088	0.511	1.980
Light transit time $/R_e$	48.59	57.31	49.30	57.51	50.10	57.41




FIG. 7. Resolution study of D21* at $N = 96, 128, 192$. Top two rows: Hamiltonian and momentum constraint norms through $t = 50$; grey dashed and dotted curves multiply the finest-grid norms by 2^2 and 2^4. Bottom row: fractional equatorial- and areal-radius changes, the change in $r\,\mathrm{Re}\,\Psi_4^{20}$ at $r = 14$, and minimum lapse over each recorded history. Near-throat constraints scale approximately at fourth order early on; the later contraction begins at resolution-dependent times.

ACKNOWLEDGMENTS

This work was supported by Gravity Frontiers (https://www.gravityfrontiers.org/en), which funded the underlying research and the development of the numerical-simulation software. The authors acknowledge the GRTL Collaboration for `GRChombo`, `GRTresna`, and `GRTeclyn`.

Appendix A: Background checks and search definition

The massless limit provides a check against the established wormhole solutions. Setting $\mathcal{V} = 0$ reduces the spectral equations to the system used in Ref. [16]. Newton iteration reproduces the saved solutions' global charges, $B = 1$, and exoticity values to machine precision.

For nonzero potential, we use two Einstein equations

that do not enter the Newton solve: the $\eta\theta$ component and the generalization of the scalar-charge identity of Ref. [12]. Their residuals, denoted E_{19} and E_c, assess the consistency of the computed background independently of the equations solved. We report maximum absolute residuals over interior nodes with $(1+\hat{\eta}^2)^{-1} \geq 0.05^2$, where $\hat{\eta}$ is the radial coordinate in the solver's unit-throat-parameter grid. The scalar-charge residual is divided by the squared charge used to normalize that calculation; the angular constraint uses its regularized compact-coordinate form. The archived checks retain these conventions for each selected background. For D21* on 210×42, they are $E_{19} = 1.6 \times 10^{-10}$ and $E_c = 4.3 \times 10^{-10}$, below the selection threshold of Sec. IV C.

For the stationary diagnostics of Sec. VI, doubling the radial quadrature and profile sampling, tightening null-ray integration, and varying the angular quadrature leave the quoted values unchanged at the displayed precision. Exact Ellis limits test the gradient, null NEC, transit and tidal normalizations. An independent proper-distance calculation of the spherical tidal tensor also agrees with the D0* measurement.

1. Search scores and dynamical interpretation

For reproducibility, the earlier S27, S50 and S85 searches used $S = X + P(C)$ on converged backgrounds, where $P = 0, 0.1, 1$ in the three residual intervals of Eq. (13); an inadmissible trial or failed solve received $W+1$, where W was the largest previous converged score, initialized to the starting $X + 1$. In historical searches the total score also included $P_{\rm rad}$, a nominal estimate of $\eta_0|\omega_I|$ with unit weight, where $|\omega_I|$ denotes an exponential growth rate. It was set to 0.25 if the static counterpart could not be solved. The potential-bearing eigenvalues from that screen are unvalidated and do not enter the stationary selection or the dynamical claims here. The resolution-screened searches use $P_{\rm rad} = 0$ by construction. The numerical background checks test the stationary equations, not stability.

A physical radial eigenmode must satisfy the linearized Einstein constraints as well as its eigenvalue equations and must converge under refinement. The known massless instability [5, 13, 14] is a benchmark for that separate calculation. The measured $3+1$ rates in Sec. V B are fits to the evolved radius, not outputs of the radial screen.

Appendix B: Exact static benchmarks

The static massless solution supplies analytical checks of the volume, local and null-ray measures. Define $a = -\gamma/(2\pi)$ and $y = \arctan(\eta/\eta_0)$. In the convention of Eq. (6), $B = 1$, $\nu = 0$, $\omega = 0$, and

$$f = -2a(\pi/2 - y), \qquad (\partial_\eta\psi)^2 = \frac{\eta_0^2(1+a^2)}{h^2}. \tag{B1}$$

In particular, $f(-\infty) = \gamma$, as required. Minimizing the sphere radius $e^{-f/2}\sqrt{h}$ gives

$$\eta_{\rm th} = a\eta_0, \qquad R_e = \eta_0\sqrt{1+a^2}\, e^{a(\pi/2-\arctan a)}. \tag{B2}$$

The angular integral in Eq. (10) cancels the 4π in the radial NEC deficit. Changing variables to y leaves

$$\begin{aligned} X &= \frac{\eta_0(1+a^2)}{R_e}\int_{-\pi/2}^{\pi/2} e^{a(\pi/2-y)}\, dy \\ &= \frac{2\sqrt{1+a^2}}{a}\, e^{a\arctan a}\sinh(\pi a/2). \end{aligned} \tag{B3}$$

The continuous $a \to 0$ limit gives the symmetric result

$$X_{\rm S0,V=0} = \frac{1}{\eta_0}\int_{-\infty}^{\infty} \frac{\eta_0^2}{\eta^2+\eta_0^2}\, d\eta = \pi. \tag{B4}$$

For D0, $a = 1/2$, so

$$X_{\rm D0,V=0} = \sqrt{5}\, e^{\frac{1}{2}\arctan(1/2)}\left(e^{\pi/4} - e^{-\pi/4}\right). \tag{B5}$$

These give 3.14159 and 4.89835, respectively, agreeing with the massless entries of Table II.

For D0 the local radial deficit is $|\rho + p_\eta| = e^f(\partial_\eta\psi)^2/(4\pi)$. Its derivative vanishes at $\eta = \eta_0/4$, giving

$$R_e^2 \max|\rho + p_\eta| = \frac{1}{4\pi}\left(\frac{20}{17}\right)^2 e^{\arctan(1/4)-\arctan(1/2)}. \tag{B6}$$

For a radial null geodesic with Killing energy $E = 1$ at positive infinity, $\dot{t} = e^{-f}$ and $|\dot{\eta}| = 1$, where a dot denotes an affine derivative. Consequently the complete ray integral is

$$\begin{aligned} R_e\int |T_{\mu\nu}k^\mu k^\nu|\, d\lambda &= \frac{R_e}{4\pi}\int (\partial_\eta\psi)^2\, d\eta \\ &= \frac{5\sqrt{5}}{64} e^{\pi/4-\frac{1}{2}\arctan(1/2)}. \end{aligned} \tag{B7}$$

Equations (B6) and (B7) yield 0.08851 and 0.30387. Both reproduce the D0 reference entries of Table IV at its displayed precision. The different peak and ray normalizations are thus checked independently of the potential-bearing solutions.

Appendix C: Convergence of the D21 evolution

The resolution study used base grids with $N = 96, 128, 192$ points across a domain of length $L = 64$. Three additional AMR levels refine the throat region, and the timestep is $\Delta t = 0.02\,\Delta x$. The runs use the same potential and numerical settings; only the base resolution changes.

Figure 7 compares the constraint norms over the common early interval through $t = 50$. The lower panels

TABLE V. Constraint orders from Eq. (C1), using linearly interpolated norms. Each cell gives 96 → 128 / 128 → 192 at the listed time and level. Finest-level orders stay near four; coarser levels vary.

constraint	$t = 10$	$t = 20$	$t = 30$	$t = 40$	$t = 50$
Ham L0	2.67 / 2.77	1.09 / 3.13	4.66 / 4.35	4.39 / 4.27	4.28 / 4.30
Mom L0	2.45 / 2.60	0.47 / 3.25	4.98 / 4.84	4.42 / 4.46	4.32 / 4.41
Ham L1	2.12 / 2.38	2.80 / 2.49	3.95 / 2.46	2.33 / 2.47	2.15 / 2.39
Mom L1	2.00 / 2.33	4.99 / 3.30	7.08 / 2.71	3.70 / 3.02	4.28 / 2.51
Ham L2	2.17 / 2.43	2.21 / 2.49	2.35 / 2.50	3.45 / 3.12	3.78 / 3.90
Mom L2	2.39 / 2.47	2.23 / 2.50	2.62 / 2.94	3.19 / 3.63	3.19 / 3.53
Ham L3	4.03 / 4.18	4.10 / 4.21	4.14 / 4.23	4.20 / 4.24	4.33 / 4.25
Mom L3	3.93 / 4.26	3.94 / 4.28	3.91 / 4.24	3.86 / 4.11	3.93 / 3.93

show the later diagnostic histories. The areal radius is estimated from the minimum area of coordinate spheres. The wave diagnostic is the change in $r\,\mathrm{Re}\,\Psi_4^{20}$ at extraction radius $r = 14$, where Ψ_4^{20} is the quadrupolar, axisymmetric mode of the Newman–Penrose curvature scalar.

Refining the grid should reduce the constraint errors, since the continuum constraints vanish. To measure how quickly they decrease, we compare norms at the same time and refinement level. For two base resolutions N_a and N_b, the measured order is

$$p_{a\to b}(t) = \frac{\ln[C_{N_a}(t)/C_{N_b}(t)]}{\ln(N_b/N_a)}, \tag{C1}$$

where C_N is the Hamiltonian or momentum constraint norm. An order near four means that halving the grid spacing reduces the error by roughly a factor of sixteen. Table V gives both neighboring-resolution comparisons. The finest level stays close to this behavior, while intermediate levels show mixed orders and transients.

The radius histories give a separate test of the dynamics. Using a one-percent decrease in equatorial radius to mark contraction, the first recorded crossings occur at $t = 58.3$, 67.3, and 80.7 in order of increasing resolution.

To interpret this shift, suppose a single unstable mode with growth time τ is excited with amplitude $a_N = a_0 N^{-p_{\mathrm{seed}}}$. In its linear regime, the fractional radius change is $\delta_N(t) \simeq a_N e^{t/\tau}$. For a fixed threshold δ_*, the crossing time then satisfies

$$\begin{aligned} t_*(N) &= \tau \ln(\delta_*/|a_0|) + p_{\mathrm{seed}}\tau \ln N, \\ t_*(N_b) - t_*(N_a) &= p_{\mathrm{seed}}\tau \ln(N_b/N_a). \end{aligned} \tag{C2}$$

The two adjacent-resolution slopes from the equatorial-radius crossings are 31.4 and 33.0. Using the independently fitted $N = 128$ areal-radius growth time $\tau = 9.49$ gives effective amplitude orders 3.31 and 3.48. Their proximity supports truncation-error seeding as an explanation of the later onset. This inference assumes that both radius diagnostics follow the same mode with a common τ; three crossing times alone do not establish a converged growth rate. Initial-data interpolation, AMR operations and the evolution discretization all contribute to the perturbation amplitude, whose order need not equal the constraint order in Table V.

[1] H. G. Ellis, "Ether flow through a drainhole: A particle model in general relativity," J. Math. Phys. **14**, 104 (1973).

[2] K. A. Bronnikov, "Scalar-tensor theory and scalar charge," Acta Phys. Polon. B **4**, 251 (1973).

[3] M. S. Morris and K. S. Thorne, "Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity," Am. J. Phys. **56**, 395 (1988).

[4] M. Visser, *Lorentzian Wormholes: From Einstein to Hawking* (AIP Press, Woodbury, NY, 1995).

[5] J. A. González, F. S. Guzmán, and O. Sarbach, "Instability of wormholes supported by a ghost scalar field. I. Linear stability analysis," Class. Quantum Grav. **26**, 015010 (2009).

[6] J. A. González, F. S. Guzmán, and O. Sarbach, "Instability of wormholes supported by a ghost scalar field. II. Nonlinear evolution," Class. Quantum Grav. **26**, 015011 (2009).

[7] M. Visser, S. Kar, and N. Dadhich, "Traversable wormholes with arbitrarily small energy condition violations," Phys. Rev. Lett. **90**, 201102 (2003).

[8] C. J. Fewster and T. A. Roman, "On wormholes with arbitrarily small quantities of exotic matter," Phys. Rev. D **72**, 044023 (2005).

[9] P. K. F. Kuhfittig, "Viable models of traversable wormholes supported by small amounts of exotic matter," arXiv:0806.1194 [gr-qc] (2008).

[10] V. Dzhunushaliev, V. Folomeev, C. Hoffmann, B. Kleihaus, and J. Kunz, "Boson Stars with Nontrivial Topology," Phys. Rev. D **90**, 124038 (2014), arXiv:1409.6978 [gr-qc].

[11] B. Kleihaus and J. Kunz, "Rotating Ellis wormholes in four dimensions," Phys. Rev. D **90**, 121503(R) (2014).

[12] X. Y. Chew, B. Kleihaus, and J. Kunz, "Geometry of spinning Ellis wormholes," Phys. Rev. D **94**, 104031 (2016).

[13] B. Azad, J. L. Blázquez-Salcedo, F. S. Khoo, and J. Kunz, "Are slowly rotating Ellis–Bronnikov wormholes stable?," Phys. Lett. B **848**, 138349 (2024).

[14] B. Azad, J. L. Blázquez-Salcedo, F. S. Khoo, and J. Kunz, "Radial perturbations of Ellis–Bronnikov wormholes in slow rotation up to second order," Phys. Rev. D **109**, 124051 (2024).

[15] N. M. Shirokov, "Wormhole dynamics: nonlinear collapse and gravitational-wave emission," arXiv:2604.00071 [gr-qc] (2026).

[16] N. M. Shirokov and I. Nachevsky, "Rotation slows the instability of Ellis–Bronnikov wormholes," arXiv:2610.29941 [gr-qc] (2026).

[17] C. Helmerich, J. Fuchs, A. Bobrick, L. Sellers, B. Melcher, and G. Martire, "Analyzing warp drive spacetimes with Warp Factory," Class. Quantum Grav. **41**, 095009 (2024).

[18] N. Hansen and A. Ostermeier, "Completely derandomized self-adaptation in evolution strategies," Evol. Comput. **9**, 159 (2001).

[19] J. Bradbury, R. Frostig, P. Hawkins, M. J. Johnson, C. Leary, D. Maclaurin, G. Necula, A. Paszke, J. VanderPlas, S. Wanderman-Milne, and Q. Zhang, `JAX`: composable transformations of Python+NumPy programs, https://github.com/jax-ml/jax (2018).

[20] D. Alic, C. Bona-Casas, C. Bona, L. Rezzolla, and C. Palenzuela, "Conformal and covariant formulation of the Z4 system with constraint-violation damping," Phys. Rev. D **85**, 064040 (2012).

[21] C. Bona, J. Massó, E. Seidel, and J. Stela, "New formalism for numerical relativity," Phys. Rev. Lett. **75**, 600 (1995).

[22] M. Alcubierre, B. Brügmann, P. Diener, M. Koppitz, D. Pollney, E. Seidel, and T. Takahashi, "Gauge conditions for long-term numerical black hole evolutions without excision," Phys. Rev. D **67**, 084023 (2003).

[23] GRTL Collaboration, `GRTeclyn`, https://github.com/GRTLCollaboration/GRTeclyn.

[24] W. Zhang *et al.*, "AMReX: a framework for block-structured adaptive mesh refinement," J. Open Source Softw. **4**, 1370 (2019).

[25] T. Andrade *et al.*, "GRChombo: an adaptable numerical relativity code for fundamental physics," J. Open Source Softw. **6**, 3703 (2021).

[26] N. M. Shirokov and I. Nachevsky, `GRTeclyn` research fork, https://github.com/Nikchik-coder/GRTeclyn, doi:10.5281/zenodo.23245933 (2026).