

isolated throat;
dash = sign of ε
ε = −0.01, level 4
ε = +0.01, level 4
unperturbed, level 4
stopped by hand
(a)
$L^2\mathcal{H}$
t
same runs;
weight = |ε|
|ε| = 0.001, level 3
|ε| = 0.1, level 3
failure (NaN)
(b)
$L^2\mathcal{M}$
t
pure quadrupole,
ε2 = 0.01, level 4
ℋ
ℳ
MOTS, t = 33
reliable to t = 60
(c)
L^2 norms
t
inflation, ε = −0.01,
L = 512 octant,
finest Δx = 1/16
ℋ
ℳ
(d)
L^2 norms
t
head-on, ℋ
used, levels 5-6-4
level 5, run past use
failure (NaN)
MOTS, t = 18
(e)
5
6
4
$L^2\mathcal{H}$
t
merger, d = 6, ℋ
used, levels 5-4
levels 6, 7, failed
segments run past use
ℳ, last segment
MOTS, t = 13
(f)
5
4
L^2 norms
t
plunge, p = 0.60, ℋ;
χ floor from t = 50
used
level-5 restart
no χ floor
reliable to t = 80
(g)
level 4
$L^2\mathcal{H}$
t
fly-by, p = 0.45, level 4
ℋ
ℳ
reliable to t = 67.6
waves at R = 20
to t = 87.6
(h)
L^2 norms
t