INVESTIGATING THE STABILITY OF SPINNING WORMHOLES WITH NONLINEAR EVOLUTIONS

NIKITA M. SHIROKOV[1,*] and ILYA NACHEVSKY[2,†]

[1] *Moscow Institute of Physics and Technology, Moscow, Russia*
[2] *Higher School of Economics, Moscow, Russia*

(Dated: October 20, 2026)

We investigate the stability of spinning Ellis–Bronnikov wormholes using fully nonlinear 3+1 numerical relativity simulations of the drainhole family. We construct the stationary metrics with a spectral solver that is exact to $\sim 10^{-10}$ at slow spin and $\sim 10^{-7}$ near the extremal limit, and validate it against the published symmetric family of Kleihaus and Kunz. In doing so we correct an inconsistency in a published field equation and map the drainhole family, for which no fast-spin solutions were known, up to $v_e = 0.99$; the 3D initial data converge at fourth order. We find that rotation slows the radial instability and then changes its fate. Along the ladder of evolved spins the collapse e-fold time grows from 4.4 (static) to 8.0 at 24% of the extremal angular momentum; at 29% and 33% the unperturbed throat turns and inflates instead of collapsing—the instability's other branch, with the seed's sign selecting the fate—and at 81% the throat is stable to $t = 150$ at three resolutions. The measured rate tracks the second-order prediction of Azad *et al.* where both exist and extrapolates to zero at ≈42% of the extremal angular momentum. The unperturbed rotating collapse radiates an $\ell = 2$, $m = 0$ burst whose energy grows with spin, while the static collapse radiates nothing; second-order perturbative initial data instead seed the instability at the level of their own truncation error.

I. INTRODUCTION

The Ellis–Bronnikov wormhole, a throat held open by a phantom scalar field [1–4], is unstable: depending on the sign of the perturbation it collapses to a black hole or inflates, on a timescale set by the throat radius [5–9]. Nonlinear 3+1 evolutions confirm both branches for the lone throat [10] and carry them into binaries, whose every channel ends in collapse, inflation or a merged horizon within a few dynamical times [11]. If traversable wormholes are to persist anywhere, something must shut this instability off.

Rotation is the oldest candidate [12]. Spinning Ellis–Bronnikov solutions are known perturbatively [13] and, beyond perturbation theory, from the elliptic solver of Kleihaus and Kunz [14, 15]: a family that starts at the static throat and ends, at equatorial rotation velocity $v_e = 1$, on an extremal Kerr black hole, the phantom field vanishing in the limit; no closed form exists [16]. On the stability side, Azad, Blázquez-Salcedo, Khoo and Kunz [17, 18] carried the radial mode to second order in spin: the static unstable mode weakens, with a positive $O(J^2)$ correction for every asymmetry, while for nearly symmetric wormholes a second unstable mode grows out of a static zero mode; the two cross near 30% of the limiting angular momentum, where the quadratic approximation gives out, and the authors conjecture the modes merge and the instability disappears, as purely imaginary modes do in five dimensions [19]. The charged analogue warns against optimism: there the merged modes continue as a complex pair, growth that oscillates, and the growth rate reaches zero only in the extremal limit [20]. Which continuation rotation takes in four dimensions is exactly the question a perturbative method cannot answer, and no rotating wormhole has ever been evolved nonlinearly.

This paper builds the tools for that evolution and takes its first step. We construct two families of backgrounds as exact stationary solutions. The symmetric family ($\gamma = 0$), massless when static, is the one with published data—the curves of Ref. [14] and the scalar charges of Ref. [15]—and we use it to validate the solver and the 3D initial data across the spin range. The drainhole family ($\gamma = -\pi$), whose static member is the mass-1 drainhole evolved in Refs. [10, 11], has no published fast-spin data at all; it is the family we evolve. The solver is gated on identities it never imposes, a discipline that pays immediately: one of the published field equations is inconsistent (Sec. II A), and second-order perturbative initial data turn out to seed the instability at their own error level (Sec. V). We then put the backgrounds on the AMR grid, verify them at fourth order, and evolve the drainhole along a ladder of nine spins (Table III). The result: rotation slows the collapse, as the perturbative sign predicts, and near 30% of the extremal angular momentum the same mode changes fate—the unperturbed throat turns and inflates—while at 81% the throat is stable over the full run.

Section II gives the family and the corrected field equations; Sec. III the solver, the validation against the published data, and the map of the parameter space (Table II); Sec. IV the numerical setup and the runs (Table III); Sec. V the evolutions; Sec. VI the conclusions. Appendix A gives the convergence of the evolutions at the two ends of the ladder. We use $G = c = 1$; the evolutions use the code units of Refs. [10, 11], $\eta_0 = 2$, in which

* shirokov.nm@phystech.edu
† ommnammashivaya@gmail.com

the static drainhole has mass $M = 1$.

II. THE SPINNING WORMHOLE FAMILY

The matter model is that of our previous work [10, 11]: Einstein gravity coupled to a phantom scalar field ψ,

$$S = \frac{1}{16\pi}\int d^4x\,\sqrt{-g}\,(R + 2\,g^{\mu\nu}\partial_\mu\psi\,\partial_\nu\psi)\,, \qquad (1)$$

whose kinetic term has the reversed sign that makes ψ a ghost. Varying the action with respect to the metric and to ψ gives the field equations of the model [Eqs. (2)–(3) of Ref. [15]],

$$R_{\mu\nu} = -2\,\partial_\mu\psi\,\partial_\nu\psi, \qquad \partial_\mu\big(\sqrt{-g}\,g^{\mu\nu}\partial_\nu\psi\big) = 0\,: \qquad (2)$$

Einstein's equations with the phantom's stress tensor as source, and the wave equation for the phantom on the curved background. Those papers evolved its best-known static wormhole, the Ellis–Bronnikov drainhole, and measured the instability that this paper asks rotation to cure. Spinning that wormhole up requires the stationary rotating solution of the same theory, which has to be constructed numerically, inside an *ansatz*—an assumed form of the metric with stationarity and axisymmetry built in and a few unknown functions left for the field equations to determine. We take the ansatz of Refs. [14, 15], who first constructed these solutions,

$$\begin{aligned} ds^2 = -e^f\,dt^2 + p^2e^{-f}\Big[e^\nu\left(d\eta^2 + h\,d\theta^2\right) \\ + h\sin^2\theta\,(d\varphi - \omega\,dt)^2\Big], \end{aligned} \qquad (3)$$

with $\eta \in (-\infty, \infty)$ and four unknown functions f, p, ν, ω of η and θ [Eq. (4) of Ref. [15]]. The field equation for p decouples from the others, and its solution with asymptotically flat boundary conditions is $p = 1$ [Eqs. (5)–(6) of Ref. [15]]; we set $p = 1$ from here on. In the trivial limit $f = \nu = \omega = 0$ the spatial metric is $d\eta^2 + h\left(d\theta^2 + \sin^2\theta\,d\varphi^2\right)$ with the fixed $h = \eta^2 + \eta_0^2$, so $\sqrt{h}$ is the radius of the sphere at coordinate η—and by construction it never vanishes: it falls from η^2 at large $|\eta|$, bottoms out at η_0 at $\eta = 0$, and rises again. The spacetime therefore has no center; a minimal sphere of radius set by η_0, the throat, joins the two asymptotically flat ends $\eta \to \pm\infty$. Choosing h hard-wires this two-ended, horizonless topology into Eq. (3), while f, ν, ω deform the geometry on top of it (h also recurs as the weight in every equation below); conversely, $\eta_0 \to 0$ closes the throat and lets the same ansatz hold a black hole, which is how extremal Kerr enters in Sec. II A. Our previous wormhole sits inside this form: with $\omega = \nu = 0$ and the closed-form arctan profile for f [15], Eq. (3) is the static drainhole of Refs. [10, 11], with $\eta_0 = a$ its throat parameter and η the isotropic radius used there, unfolded through the throat (Sec. IV A); $f = 0$ gives the massless Ellis wormhole.

Substituting the ansatz (3) into the field equations (2) reduces them in three steps. We follow the reduction of Kleihaus and Kunz [14], given in full in Sec. II of Ref. [15]; equation numbers in brackets below refer to Ref. [15]. First, the phantom field equation—the wave equation in Eq. (2), which stationarity and axisymmetry reduce to a function of η alone—integrates in closed form to $\psi = (D/\eta_0)\,[\arctan(\eta/\eta_0) - \pi/2]$ [Eqs. (7)–(9)]: the scalar enters the rest of the problem only through its constant of integration D, the scalar charge. Second, because $\partial_\mu\psi$ points along η, the source term $-2\,\partial_\mu\psi\,\partial_\nu\psi$ survives only in the $\eta\eta$ Einstein equation; each other nontrivial component (tt, $t\varphi$, $\theta\theta$, $\varphi\varphi$, $\eta\theta$) is a statement about the metric functions alone, and these organize by order and role. Two are second-order elliptic equations,

$$\begin{aligned} 0 = {}& \partial_\eta(hs\,\partial_\eta f) + \partial_\theta(s\,\partial_\theta f) \\ & - hs^3e^{-2f}\left[h(\partial_\eta\omega)^2 + (\partial_\theta\omega)^2\right], \end{aligned} \qquad (4)$$

$$\begin{aligned} 0 = {}& \partial_\eta\left(h^2s^3e^{-2f}\partial_\eta\omega\right) \\ & + \partial_\theta\left(hs^3e^{-2f}\partial_\theta\omega\right), \end{aligned} \qquad (5)$$

with $s \equiv \sin\theta$ [Eqs. (10)–(11)]: Eq. (4) is a Laplace-type equation for the redshift f, sourced by the squared gradients of the frame dragging—rotation gravitates—and Eq. (5) is a total divergence, the local form of angular-momentum conservation, which integrated over the domain forces the same J through every $\eta = \text{const}$ surface. The single off-diagonal component is only first order in ν,

$$\begin{aligned} 0 = {}& -hs\,\partial_\eta f\,\partial_\theta f + h\cos\theta\,\partial_\eta\nu \\ & + \eta s\,\partial_\theta\nu + h^2s^3e^{-2f}\partial_\eta\omega\,\partial_\theta\omega, \end{aligned} \qquad (6)$$

so ν follows by integration once f and ω are known [Eq. (13)]. (The remaining diagonal components give a second-order equation for ν, Sec. II A.) Third, the one sourced component, $R_{\eta\eta} = -2D^2/h^2$ [Eq. (18)], rearranges into an algebraic identity for the charge [Eq. (19)]: the metric functions fix D^2 at every point, so its constancy over the domain is a solution test that comes for free. Three unknown functions thus face more equations than they need; the Bianchi identities make the overdetermined set consistent on a true solution, and a numerical scheme must choose which subset to impose—a choice Sec. II A shows matters. The boundary conditions select the family: Minkowski at $\eta \to +\infty$, and $f \to \gamma$, $\omega \to \omega_\infty$ at $\eta \to -\infty$, so the constant γ measures how unequal the two ends are and ω_∞ sets the spin.

A solution of this system is three functions and one number, $(f, \nu, \omega; D)$. Every physical quantity we quote—the columns of Table II—is read off them directly [14, 15]:

- far from the throat the metric settles to the field of a point mass, $f \to -2\mu_+/\eta$ as $\eta \to +\infty$ and $f \to \gamma + 2\mu_-/\eta$ as $\eta \to -\infty$, so the end masses $\mu_\pm$ are the $1/\eta$ coefficients of f;
- likewise $\omega \to 2J/\eta^3$, and the charges obey the Smarr-type relation $\mu_+ + \mu_- = 2\,\omega_\infty J$ [15];

- the throat is the sphere of minimal area, at $\eta_t = 0$ when the two ends are equal ($\gamma = 0$) and displaced toward the positive-mass end otherwise ($\eta_t = \eta_0/2$ for the static drainhole): its equatorial circumference gives the radius $R_e = e^{-f/2}\sqrt{h}\,\big|_{\eta_t,\,\theta=\pi/2}$, its meridians the polar radius R_p, its area the areal radius R_A—three numbers that coincide while the throat is round and split as rotation flattens it;
- the throat is dragged at the angular velocity $\Omega = \omega\,\big|_{\eta_t,\,\theta=\pi/2}$, and $v_e = R_e\,\Omega$ is its equatorial linear velocity in units of c: the dimensionless spin label used throughout, 0 when static and 1 at the extremal Kerr endpoint.

We construct two one-parameter families, labelled by ω_∞ (equivalently J or v_e):

(a) *the drainhole family*, $\eta_0 = 2$, $\gamma = -\pi$: its static member, with end masses $\mu_\pm = \pm 1$, is the wormhole of Refs. [10, 11] (the asymmetry $C = 1$ of Refs. [17, 18]).

(b) *the symmetric family*, $\gamma = 0$ ($C = 0$), that of Refs. [14, 15]: equal masses on both ends, the massless Ellis wormhole when static; for the 3D checks it is rescaled $\eta_0 = 1 \to 2$, so its throat sits on family (a)'s grid.

A. An inconsistency in the published field equations

Refs. [14, 15] close the system with a second-order equation for ν, Eq. (12) of Ref. [15]. That equation is inconsistent with the rest. The scalar-free component $R_{\theta\theta} = 0$, with the second derivatives of f and ω eliminated by Eqs. (4)–(5), gives instead

$$0 = \partial_\eta(hs\,\partial_\eta\nu) + s\,\partial_\theta^2\nu - \cos\theta\,\partial_\theta\nu + s\,(\partial_\theta f)^2 - hs^3e^{-2f}\left[h(\partial_\eta\omega)^2 + 2(\partial_\theta\omega)^2\right], \tag{7}$$

where the printed form has $-2(\partial_\theta\omega)^2$ and no $s\,(\partial_\theta f)^2$ term. Extremal Kerr fits the ansatz (3) in closed form (with $\eta_0 = 0$), and substituted symbolically, at 160-digit arithmetic, it satisfies Eqs. (4)–(6), Eq. (7) and the D^2 identity exactly—while leaving a residue of up to 0.35 (at $M = 1$) in the printed ν equation alone. The error also shows dynamically: solutions of the printed elliptic system converge at every resolution, yet carry a violation of the unimposed Eq. (6) and D^2 identity that scales as J^4 and does not shrink with the grid—the fingerprint of a wrong continuous equation, not of numerics (first seen as a spread of 5.5×10^{-7} in D^2 at $J = 0.25$). Our solver imposes neither ν equation (Sec. III), and Eq. (7) holds on all 200 saved solutions of both families to $\leq 7\times10^{-8}$ of its largest term within 20 throat radii, where the printed form misses by up to 53%. Every published conclusion of Refs. [14, 15] that we can test survives the correction (Sec. III).

III. BACKGROUNDS ACROSS THE SPIN RANGE

The solver turns the system of Sec. II into numbers in five steps (Fig. 1). First, the infinite domain is made finite: the coordinate $x = (2/\pi)\arctan(\eta/\eta_0)$ maps both asymptotic ends to $x = \pm1$, and the unknowns f and ω are expanded in Chebyshev polynomials of x and even cosine modes of θ. The expansion is spectral: for smooth solutions the error falls exponentially with the number of modes, so modest grids reach near machine precision. Second, the two elliptic equations (4)–(5) are solved for the expansion coefficients by Newton iteration—guess, linearize, correct—until their residual sits at round-off. Third, ν and the scalar charge are recovered without a second-order ν equation: the first-order constraint (6) and the D^2 identity together overdetermine ν and the single number D^2, so both are extracted as one linear least-squares problem—and the redundancy is a feature, since the leftover misfit measures the error. Fourth, each family is walked in spin: every converged solution seeds the Newton iteration at a slightly higher spin, with J as the continuation parameter (ω_∞ is not monotone along a family), and the grid refines automatically, from 48×12 up to 240×60, whenever the acceptance gates fail. Fifth—the point of the whole design—a solution is accepted only on identities the solver never imposes: the spread of D^2 over the domain ($\leq 10^{-11}$), the Smarr relation ($\leq 10^{-10}$), equality of the angular-momentum flux through the two ends ($\leq 1.3\times10^{-9}$), the asymptotic tails ($\leq 5\times10^{-12}$), and an independent, symbolically generated residual of the full Einstein equations, evaluated out to 20 throat radii: 1.0×10^{-10} at $J = 0.25$ and 4.2×10^{-10} at $J = 1$ against the static member's 2×10^{-11} round-off floor—where the $O(J^2)$ perturbative data give 3×10^{-5} and 1×10^{-2} on the same meshes—staying near 10^{-9} up to $v_e \approx 0.85$ and reaching $\sim10^{-7}$ at $v_e = 0.91$–0.99. The high-spin 10^{-7} is round-off, not solution error: it does not shrink on finer grids, while the mass and scalar charge come out identical to 12 digits across them; and the test does catch genuine failures, scoring a deliberately underresolved solution at 10^{-4}, which the resolution check rejects. Beyond 20 throat radii the residual's normalization inflates round-off and the test is switched off; that outer region is covered by the resolution study, by the end charges, and by the 3D constraint checks of Sec. IV. The products: global charges stable to 10 digits across three grids and, for evolution, each background exported as a 1025×65 table whose bilinear interpolation error is $\leq 10^{-7}$ up to $v_e = 0.9$.

Figure 2 validates the symmetric family against the published curves. The four published anchor solutions at $\omega_\infty = 0.5$ [15] split: the $\gamma = 0$ and $\gamma = -1$ scalar charges agree to all printed digits, while $\gamma = 1$ and $\gamma = 2$ differ by 2×10^{-4} and 6.6×10^{-3} from our grid-converged values—their finite-difference error, growing with γ (Table I).

Figure 3 and Table II are the map of the parameter space the foundation papers left open, including the




FIG. 1. The solver loop for one family (Sec. III). Thin boxes impose equations; the heavy box tests identities the solver never imposes, and only it decides acceptance. A failed check refines the grid and solves the same J again; an accepted solution seeds the next J. The loop gave 462 accepted solutions over 20 continuation legs, both families to $v_e \approx 0.99$.




FIG. 2. The symmetric family ($\eta_0 = 1$, $\gamma = 0$; 81 solutions, $v_e = 0.02$–0.991, plus the exact static member) in the normalization of Ref. [14] ($M_0 = R_e$, $J_0 = R_e^2/2$, $\Omega_0 = 1/R_e$): our curves, the digitized points of their Fig. 1 (open symbols), and closed-form Kerr (dashed). The digitization's reading error, measured by reading their printed Kerr curves the same way, is rms 1–6 $\times 10^{-4}$; digitized minus ours is rms 2.5–8 $\times 10^{-4}$, max 2.5×10^{-3}, over all five curves. The two solvers agree to the precision a printed figure carries.

massive drainhole family, from which binaries can be built [11]. Two of its features matter later. First, both families violate the Kerr bound (Fig. 3b), the symmetric family with $J/\mu_+^2 \approx 85$ already at $v_e = 0.24$, falling monotonically to 1 at extremality since its static member is massless. A collapsing overspun throat cannot settle to a Kerr black hole of the same mass and spin, which makes these backgrounds sharp cosmic-censorship initial data. Second, the drainhole's negative far-end mass μ_- rises through zero at $v_e \approx 0.7$ and turns positive: fast rotation un-does the static drainhole's negative-mass end.

TABLE I. Scalar charge D of the anchor solutions ($\eta_0 = 1$, $\omega_\infty = 0.5$) against Ref. [15]. Along each fixed-γ family, ω_∞ rises, turns over and descends, so $\omega_\infty = 0.5$ is reached twice; the published values sit on the post-turnover branch. Our values are grid-converged to the digits shown; the $\gamma = 2$ family continued to $J = 172$ confirms no third crossing exists. The pre-turnover branch is new.

γ	D (this work)	D [15]	D (pre-turnover, new)
−1	0.716368	0.71636	0.969194
0	0.552771	0.55277	0.986013
1	0.4314654	0.43155	1.008136
2	0.3375993	0.33984	1.048161

IV. NUMERICAL SETUP

A. 3D initial data

The stationary metric (3) goes onto the Cartesian AMR grid through the isotropic radius r, $\eta = r - \eta_0^2/4r$: the far end $\eta \to -\infty$ compactifies to the puncture $r = 0$, as for the static drainhole [10]. The induced metric and the fields are read from the tables; the lapse and shift start from the stationary data, $\alpha = e^{f/2}$, $\beta^\varphi = -\omega$, so the extrinsic curvature $K_{ij} = (D_i\beta_j + D_j\beta_i)/2\alpha$ has only $r\varphi$ and $\theta\varphi$ components and the scalar momentum vanishes; at the puncture the frame rotates at ω_∞.

Three checks validate the pipeline. (i) At $t = 0$ the Hamiltonian and momentum constraints converge at fourth order inside the throat region and are limited outside only by the tables' interpolation error (growing as $1/\Delta x^2$ and $1/\Delta x$ respectively), and the throat's ray radii converge to the solver's. We check this on the drainhole at $v_e = 0.51$ on three grids (ray radii within 5×10^{-5} at $N = 192$, $L = 64$) and, across the spin range, on the symmetric family at seven spins $v_e = 0.24$–0.99 with three resolutions each (within 7×10^{-5} at $v_e = 0.24$ to 9×10^{-4} at 0.99, $N = 256$). (ii) On the same symmetric set, the 3D ADM mass integral evaluated on the




FIG. 3. The drainhole ($\eta_0 = 2$, $\gamma = -\pi$) and symmetric ($\eta_0 = 1$, $\gamma = 0$) families across the spin range, against v_e: (a) mass, angular momentum and equatorial throat radius; (b) J/M^2 against the Kerr bound; (c) the ratio of end masses; (d) the flattening R_p/R_e; (e) the exotic matter, the phantom field's integrated null-energy violation $\mathcal{I} = \int(\rho + p_\eta)\, dV$ over the $t = \text{const}$ slice [21], per equatorial throat radius, with the 3D runs (squares) placed by the scalar charge measured on each run's own $t = 0$ slice, which reproduces its background's to 3×10^{-4}; (f) the throat angular velocity, which peaks and falls to zero at extremality. The symmetric family violates the Kerr bound at every spin and the drainhole family exceeds it by up to $\sim 20\%$ at intermediate spin; the exotic matter per throat radius falls a hundredfold along the drainhole family and tenfold along the symmetric one by $v_e = 0.99$; both families end on extremal Kerr.

TABLE II. The two families at representative spins, in units of η_0: rotation velocity v_e, angular momentum, mass of the $\eta \to +\infty$ end, ratio of end masses, scalar charge, equatorial throat radius, flattening, and throat angular velocity—each read off the solution's functions as defined in Sec. II. Every row is a converged solution that passed the acceptance gates of Sec. III.

family	v_e	J/η_0^2	μ_+/η_0	μ_-/μ_+	D/η_0	R_e/η_0	R_p/R_e	$\Omega\,\eta_0$
(a) drainhole	0.213	0.2535	0.5574	−0.850	1.1004	2.0319	0.9873	0.1046
	0.513	1.0278	0.9302	−0.339	1.0007	2.6271	0.9194	0.1952
	0.699	3.0417	1.6557	−0.032	0.8598	3.8864	0.8351	0.1798
	0.900	34.550	5.8400	+0.179	0.5327	11.911	0.6926	0.0755
	0.950	152.16	12.316	+0.201	0.3785	24.751	0.6504	0.0384
	0.990	4053.8	63.666	+0.208	0.1698	127.36	0.6165	0.0078
(b) symmetric	0.239	0.0542	0.0252	1	0.9881	1.0299	0.9913	0.2323
	0.434	0.1158	0.0906	1	0.9583	1.1101	0.9699	0.3911
	0.667	0.2683	0.2666	1	0.8854	1.3420	0.9197	0.4969
	0.847	0.7007	0.6305	1	0.7653	1.8836	0.8440	0.4499
	0.898	1.0989	0.8685	1	0.7043	2.2722	0.8092	0.3952
	0.952	2.4701	1.4383	1	0.5971	3.2703	0.7539	0.2911
	0.990	12.235	3.4312	1	0.4157	7.0595	0.6795	0.1402

grid equals the same integral on the exact table metric to $\leq 10^{-12}$, and its Richardson extrapolation meets the solver's mass to 2×10^{-5}–7×10^{-4} (at worst 0.7% relative, at $v_e = 0.99$); the ADM angular momentum matches to between 2×10^{-3} ($v_e = 0.24$) and 2×10^{-5} (0.99). (iii) Dynamically, the static drainhole run through this pipeline is a numerical twin of the production hold of Ref. [11] over the full record: $t = 0$ bit-identical, Hamiltonian norm within 5×10^{-7} for all $t \leq 100$, and collapse milestones within one time unit.

TABLE III. The runs: the drainhole family (a) along the ladder, in code units. $8\pi J/A$ is the angular momentum in per cent of its extremal bound and $M_{\rm ADM}$ the mass of the $\eta \to +\infty$ end (both from the background tables). The finest Δx is 0.0625 at $N = 128$ on $L = 64$ and at $N = 256$ on $L = 128$, and scales inversely with N within a triplet. Recipe S (slow): $\sigma = 0.1$, $\chi \geq 10^{-8}$; F (fast): $\sigma = 0.4$, $\chi \geq 10^{-4}$. Shell: the damping layer of Sec. IV C on (✓) or the background-relative boundary alone (–). Trust: the last time the run is quoted. GW: the $(2,0)$ burst at $R = 14$ inside the trust window (✓), only its onset (the signal still rising at the window's end), or nothing above the extraction floor (–). The last three rows are in flight at this draft.

rung	v_e	$8\pi J/A$	$M_{\rm ADM}$	L	N	recipe	shell	stop t	trust	GW	outcome
0c	0	0	1.000	64	128	S	–	100	60	–	collapses, $\tau = 4.4$; dies in its horizon, $t = 62.7$
0.2c	0.210	12.3	1.112	64	96/128/192	S	✓	100	55/100/100	✓	collapses, $\tau = 5.0$; horizon settles at $R = 2.52$
0.2c, kicks	0.210	12.3	1.112	64	128	S	–	100	42/61	onset	$\varepsilon = \pm 10^{-3}$: pushed out collapses, in inflates
0.2c, far zone	0.210	12.3	1.112	128	256	S	–	150	150	✓	the $(2,0)$ burst on spheres $R = 14$–45 (Fig. 5)
0.30c	0.304	18.3	1.248	64	128	S	–	100, 130	130	✓	collapses, $\tau = 5.9$; horizon at $R = 2.567$
0.40c	0.395	24.4	1.448	64	128	S	–	100	100	onset	collapses, $\tau = 8.0$; MOTS leaves throat, $t = 92$
0.46c	0.456	28.8	1.634	64	128	S	–	100, 150	100, 121	onset	-1.1% at $t = 83$, inflates from $t = 101$, $\tau = 12.1$
0.5c	0.513	33.3	1.860	64	192	S	✓	100	100	–	shrinks, -1% at $t = 99$: the seed's other sign
0.5c	0.513	33.3	1.860	64	128	S	–	150	95	–	inflates from $t \approx 70$, $\tau = 8.8$
0.5c	0.513	33.3	1.860	128	256	F	✓	150	–		running
0.6c	0.592	40.3	2.305	128	256	F	✓	150	–		running
0.7c	0.699	51.4	3.311	128	256	F	✓	150	–		running
0.9c	0.900	80.8	11.68	128	128/192/256	F	✓	150	150	–	stable: areal drift $\leq 7 \times 10^{-5}$ at all three N




FIG. 4. One rate, two fates: the measured rate $1/\tau$ (left axis; τ right) against the angular momentum as a fraction of its extremal bound. Filled circles: collapsing rungs, τ from the log-linear fit of the areal-radius deficit over the 0.2–1% band. Filled triangles: inflating rungs, τ from $\ln|dR_A/dt|$ over the +0.2–1% growth band (the same fit on a pure exponential). Open circle: the stable 0.9c. Dashed: the fit $a - b\,(J/A)^2$ through both branches, which reaches zero at 42%. Open diamonds: the second-order rates of Refs. [17, 18] at $C = 1$; dotted rules: their conjectured end of the instability (≈30%) and the zero of their second-order rate (33%). The outcome: the mode slows with spin, changes fate from collapse to inflation near 30% instead of dying there, and is unmeasurably slow by 81%.

B. Evolution scheme

Evolution uses `GRTeclyn` [27–29] with the setup of Refs. [10, 11]: CCZ4 [24], 1+log slicing [25], Gamma-driver shift [26], fourth-order stencils, RK4, Kreiss–Oliger dissipation, and a floored conformal factor at the puncture. The timestep is $\Delta t = 0.02\,\Delta x$ in every run. That value was adopted empirically in Refs. [10, 11]; we traced what set it to a grid-scale mode pinned to the fixed coarse–fine interfaces of the nested refinement boxes, dissipation-starved at $\sigma = 0.1$, whose growth rate rises with the timestep. Constraint norms evaluated per refinement level, with cells covered by finer levels masked, expose it by $t \approx 10$; the domain-wide norm is blind to it until breakout. Raising the dissipation where the mode lives removes it—at $\sigma = 0.4$ the interface-level norms sit 10^2–10^4 below the $\sigma = 0.1$ reference by $t = 100$—but the larger timesteps this would permit do not transfer to the spinning backgrounds, which lose the far side by $t \approx 22$–48 at $\Delta t = 0.035$–$0.05\,\Delta x$. The production runs therefore take no speed-up, and the only scheme parameter that varies along the ladder is the dissipation (recipes S and F of Table III). Its one measured side effect, a constant ∼4 time-unit advance of the collapse clock from the stronger initial smoothing, keeps rate comparisons at matched σ.

C. Long evolutions: the outer boundary

Evolutions that must outlive the outer-boundary crossing time $t \approx L$ need the boundary made harmless. Kreiss–Oliger dissipation cannot do it—its strength falls as the sixth power of wavelength, so the sponge shell of Refs. [10, 11] is transparent to the long outgoing gauge pulse—and a flat-space Sommerfeld condition assumes asymptotics the wormhole does not have (mass 1–12 sits inside the box). Two changes in the same outer shell fix it. The Sommerfeld condition is imposed on the deviation of the evolved fields from the stationary background rather




FIG. 5. The $(2,0)$ wave of the unperturbed rotating collapse. (a) $r\,\Delta\psi_4^{20}$ of the 0.2c far-zone run on spheres $R = 14$–45 against retarded time: one waveform with ring-down period ≈24; only the first lobe falls with R (near field), the extrema are flat in R to 1.5% over $R = 30$–45, and this box reproduces the $L = 64$ run's throat history to 3.2×10^{-4}. (b) The wave against spin at $R = 14$, each run on its own collapse clock $u = t - R - t_{10}$ (t_{10}: its equatorial deficit at 10%): the 0.30c burst has the 0.2c's shape and timing at ∼1.5× the amplitude, and the static run is drawn at zero—a spherical collapse radiates no quadrupole; its measured record stays below 3×10^{-8}. (c) The radiated energy $E_{20} = \frac{1}{16\pi} \int |\int r\psi_4\, dt'|^2\, dt$ over the aligned burst, per unit $M_{\rm ADM}$, at $R = 18$ (filled) and $R = 14$ (open): $3.3 \times 10^{-7} \to 8.0 \times 10^{-7}$ from the 0.2c to the 0.30c, a factor 2.5–3—the square of the amplitude ratio in (b)—with the static at zero. (d) The speed of each extremum between neighbouring spheres, against the local coordinate speed of light (crosses): the burst propagates at light speed. The outcome: an unperturbed, axisymmetric rotating collapse radiates a quadrupole burst whose energy grows with spin as the amplitude squared, and the static collapse radiates nothing.

than on the fields themselves; and a damping layer can in addition drive every evolved field back to that background at rate A, $\partial_t u \to \partial_t u - A\, f(r)\, (u - u_{\rm bg})$, with $f(r)$ the sponge's quartic ramp and $u_{\rm bg}$ the run's own initial data, rebuilt exactly from the background tables; $A = 0$ recovers the base scheme identically. The layer is validated on the static hold at $A = 1$ against the reference evolution: $|K|$ falls two orders of magnitude across the shell, while everything inside it—the throat radii to four digits, the constraint norms, the near-zone ring—is unchanged, so it absorbs at all wavelengths without touching the interior. On the $L = 64$ box, whose shell ($r = 24$–32) contains the outermost extraction sphere, a side-by-side test showed the background-relative boundary alone keeps the outer levels cleaner, so those runs take the layer off; the $L = 128$ boxes ($r = 48$–64, outside every sphere) run with it on (Table III). With the boundary treated this way the $L = 64$ runs, previously trusted only to the crossing time $t \approx 64$, run to their stop times $t = 100$–150, with trust windows set by the constraints instead. The same comparison shows the late-time growth previously attributed to boundary reflection is sourced by the collapse region itself, and is removed by the raised dissipation of Sec. IV B.

D. The runs

Stability claims follow the triplet protocol of Ref. [23]: a quoted growth must have an e-fold time that converges to a nonzero value across resolutions, constraints that converge at the code's order while it grows, and a gauge-invariant measure—the throat's areal radius, or the marginally outer-trapped surface (MOTS) found spectrally on every output—growing with it. Every run carries three nested refinement levels on the throat, and the box is picked by the rung's mass: $M_{\rm ADM}$ triples along the family (1.11 at $v_e = 0.21$, 3.31 at 0.70), so the face of the $L = 64$ box recedes from $25\, M_{\rm ADM}$ of clearance at slow spin to under $10\, M_{\rm ADM}$ at $v_e = 0.70$. Rungs up to $v_e = 0.51$ therefore run on the $L = 64$, $N = 128$ grid of the static hold of Ref. [11], and rungs from $v_e = 0.59$ on $L = 128$, $N = 256$ at the same finest Δx. Ψ_4 is extracted on spheres at $R = 10$–30 ($L = 64$) and $R = 14$–45 ($L = 128$); the signal is the $(2,0)$ mode. Resolution triplets anchor the two ends of the ladder (Appendix A). Table III lists every run, from the static reference through the $O(J^2)$ regime [17, 18] to far beyond any perturbative result.

V. THE INSTABILITY ALONG THE LADDER

Perturbative initial data fail first. Our first spinning holds used the $O(J^2)$ backgrounds of Ref. [18] at $J = 0.25$ and $J = 1$ on the drainhole family. Both throats inflated where the static throat collapses—the same unstable mode with the opposite sign ($\tau = 5.8$ and 5.6). But expressed as the equivalent radial kick of Ref. [11], the push was $\varepsilon \approx -3 \times 10^{-5}$ at $J = 0.25$ and -7×10^{-3} at $J = 1$: growth by a factor ∼230 for a factor 4 in J, i.e. $\sim J^4$—the size of the $O(J^2)$ data's own truncation error,

which the independent Einstein residual puts at 10^{-2} at $J = 1$ (Sec. III). Second-order data seed the very instability under study at their own error level; nothing at slow spin can be concluded from them.

The collapse branch. On exact data (Table III) the verdict reverses, and the unperturbed throat collapses on the static mode, later the faster it spins: the areal deficit passes 1% at $t = 43.7$, 46.9, 51.4 and 60.2 along the rungs $v_e = 0$–0.40, with e-fold times $\tau = 4.4$, 5.0, 5.9, 8.0 (Fig. 4). Each collapse ends in a black hole: the 0.2c horizon settles at $R = 2.524$ ($M_{\rm MS} = 1.26$), the 0.30c at $R = 2.567$ by $t = 130$; the 0.40c's MOTS leaves the throat at $t = 92$ and reads $R = 3.22$, still shrinking, at $t = 100$. In each case the hold is seeded only by discretization error, so the ordering of rates is the ordering of $|\omega_I|$.

The inflating branch. At $v_e = 0.456$ (28.8% of extremal) the fate changes: the areal radius dips to -1.12% at $t = 83$, turns, crosses zero at $t = 101$ and grows exponentially, $\tau = 12.1$. The 0.5c behaves the same way (-0.30% at $t = 65$–68, then $+6.9\%$ by $t = 100$, $\tau = 8.8$), and the MOTS is lost once a throat expands—no trapped surface forms. These are the two fates of one instability, and the seed's sign selects between them, exactly as declared kicks show at 0.2c (pushed in, the throat inflates; pushed out, it collapses; both pass 1% at $t \approx 25$ at the same rate): the 0.5c run whose absorbing shell gives the opposite effective seed shrinks instead (-1% at $t = 99$). At the fast end the 0.9c throat is simply stable, its areal radius flat to $\leq 7 \times 10^{-5}$ over the full $t = 150$ window at all three resolutions (Appendix A).

Against the perturbative prediction. Figure 4 puts both branches on one axis. The measured rate matches the second-order $C = 1$ rates of Refs. [17, 18] within 3% at $v_e = 0.30$ and 0.40 (5.9 vs 6.1, 8.0 vs 8.0). Where they conjecture the instability ends, we find instead that the mode survives the crossing with its magnitude intact and only its fate changed: the 28.8% rung inflates with $\tau = 12.1$ against their 13.2 (9%), and the 33% rung is still unstable where their second-order rate has already died. A quadratic fit through both branches reaches zero at 42%. The comparison is close but not like for like: their τ is a linear mode in Killing time, ours a nonlinear areal-radius fit.

Gravitational waves. The rotating collapse radiates with no nonspherical perturbation put in (Fig. 5): one $(2, 0)$ burst at light speed, whose energy grows with spin as the amplitude squared, while the spherical static collapse radiates nothing. Rotation thus imprints itself on the burst a collapsing throat emits, before any horizon forms.

VI. CONCLUSIONS

First, the spinning Ellis–Bronnikov parameter space is now mapped as data, not curves: both families from slow rotation to $v_e \approx 0.99$, including the drainhole family for which nothing was published, with the published ν equation, Eq. (12) of Ref. [15], corrected to Eq. (7). The practical lesson generalizes: a solver gated only on the equations it imposes would have passed every test while solving the wrong system; the unimposed identities caught it. Second, the pipeline from these solutions to 3+1 evolution is validated at fourth order, and $O(J^2)$ initial data are shown to be unusable for stability work: their truncation error seeds the instability itself. Third, the first nonlinear evolutions of rotating wormholes show one rate with two fates: the radial collapse slows with spin ($\tau = 4.4 \to 8.0$ from the static to 24% of the extremal angular momentum), the sign predicted at $O(J^2)$ [17, 18]; near 30%, where the perturbative mode crossing was conjectured to end the instability, the unperturbed throat inflates instead, with the mode's magnitude intact; the measured rate extrapolates to zero at ≈42%, and at 81% the throat is stable to $t = 150$ across three resolutions—answering, for this family, the four-dimensional stability question left open in Ref. [14]. The rotating collapse also radiates a $(2, 0)$ burst from unperturbed, axisymmetric data, its energy growing with spin as the amplitude squared, while the static collapse radiates nothing.

VII. REPRODUCIBILITY AND COST

The solver, the evolution code and the figure scripts are in the `GRTeclyn` [27] fork at https://github.com/Nikchik-coder/GRTeclyn (branch `feature/spinning`). Its `results/spinningwormhole` pack holds the solver's accepted solutions, the headers of the background tables the evolutions start from, and the parameter files and diagnostic streams of the finished runs of Table III, with a run registry.

Every evolution ran on one NVIDIA H100 80 GB device, with `GRTeclyn` built CUDA-native against AMReX 26.02. A rung on the $L = 64$, $N = 128$ grid advances 9–13 code units per hour, and a $t = 100$ evolution takes 7.6–11.1 h; the $N = 192$ members run at 5 units per hour (20 h to $t = 100$), peaking at 53 GiB of the device, and the $L = 128$, $N = 256$ runs at 15.5 (9.7 h to $t = 150$). The 17 finished evolutions of Table III total 167 GPU-hours, each run's cost recomputed from its archived logs: the logged speed and time span.

ACKNOWLEDGMENTS

This work was supported by Gravity Frontiers (https://www.gravityfrontiers.org/en), which funded the underlying research and the development of the numerical-simulation software. The authors acknowledge the GRTL Collaboration for `GRChombo`, `GRTresna`, and `GRTeclyn`.

Appendix A: Convergence of the evolutions

The two triplets are the 0.2c, the slowest spinning collapse (Fig. 6), and the 0.9c, the stable fast rung (Fig. 7), both read through the per-level constraint norms of Sec. IV B.




FIG. 6. Convergence of the 0.2c rung over its triplet $N = 96/128/192$ ($L = 64$; finest Δx in each panel title). (a)–(d) Hamiltonian and (e)–(h) momentum constraint norms per refinement level; dashed and dotted, the $N = 192$ norms rescaled by 2^2 and 2^4: the measured $N = 96$ norms lie between the second- and fourth-order brackets on every level while the collapse grows. (i) The equatorial throat radius against the stationary solution, falling with resolution. (j) The collapse, each run aligned at its own 10% time: the three resolutions land on one curve. (k) $r\,\Delta\psi_4^{20}$ at $R = 14$, aligned: $N = 128$ and 192 overlap ($N = 96$ dies in its collapse before the burst arrives). (l) The MOTS radius, aligned: one horizon, settling at $R = 2.52$. The outcome: the collapse rate, the waveform and the horizon are grid-converged at third to fourth order.




FIG. 7. Stability of the 0.9c rung over its triplet $N = 128/192/256$ ($L = 128$; finest Δx in each panel title), to $t = 150$. (a)–(h) Constraint norms per level with the same second-/fourth-order brackets as Fig. 6: every level converges with resolution and nothing breaks out over the window. (i) The equatorial throat radius holds the stationary value, its offset falling with resolution. (j) The areal throat radius drifts by $\leq 7 \times 10^{-5}$ over the full window. (k) The $(2, 0)$ extraction at $R = 18$, relative to its $t = 0$ read-off: fluctuations at the 10^{-4} level, no burst. (l) The far-side puncture lapse falls smoothly at all three resolutions (the gauge, not the geometry). The outcome: the 0.9c throat is stable on its window at every resolution—no growing radius, no constraint breakout, no wave.

[1] H. G. Ellis, "Ether flow through a drainhole: A particle model in general relativity," J. Math. Phys. **14**, 104 (1973).

[2] K. A. Bronnikov, "Scalar-tensor theory and scalar charge," Acta Phys. Polon. B **4**, 251 (1973).

[3] M. S. Morris and K. S. Thorne, "Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity," Am. J. Phys. **56**, 395 (1988).

[4] M. Visser, *Lorentzian Wormholes: From Einstein to Hawking* (AIP Press, Woodbury, NY, 1995).

[5] J. A. González, F. S. Guzmán, and O. Sarbach, "Instability of wormholes supported by a ghost scalar field. I. Linear stability analysis," Class. Quantum Grav. **26**, 015010 (2009).

[6] J. A. González, F. S. Guzmán, and O. Sarbach, "Instability of wormholes supported by a ghost scalar field. II. Nonlinear evolution," Class. Quantum Grav. **26**, 015011 (2009).

[7] H. Shinkai and S. A. Hayward, "Fate of the first traversible wormhole: Black-hole collapse or inflationary expansion," Phys. Rev. D **66**, 044005 (2002).

[8] K. A. Bronnikov, J. C. Fabris, and A. Zhidenko, "On the stability of scalar-vacuum space-times," Eur. Phys. J. C **71**, 1791 (2011).

[9] F. Cremona, F. Pirotta, and L. Pizzocchero, "On the linear instability of the Ellis–Bronnikov–Morris–Thorne wormhole," Gen. Relativ. Gravit. **51**, 19 (2019).

[10] N. M. Shirokov, "Wormhole dynamics: nonlinear collapse and gravitational-wave emission," arXiv:2604.00071 [gr-qc] (2026).

[11] N. M. Shirokov and I. Nachevsky, "Binary Ellis–Bronnikov wormholes in 3+1 numerical relativity: mergers, fly-bys and gravitational waves," arXiv:2610.13206 [gr-qc] (2026).

[12] T. Matos and D. Núñez, "Rotating scalar field wormhole," Class. Quantum Grav. **23**, 4485 (2006).

[13] P. E. Kashargin and S. V. Sushkov, "Slowly rotating wormholes: The first order approximation," Grav. Cosmol. **14**, 80 (2008); "Slowly rotating scalar field wormholes: The second order approximation," Phys. Rev. D **78**, 064071 (2008).

[14] B. Kleihaus and J. Kunz, "Rotating Ellis wormholes in four dimensions," Phys. Rev. D **90**, 121503(R) (2014).

[15] X. Y. Chew, B. Kleihaus, and J. Kunz, "Geometry of spinning Ellis wormholes," Phys. Rev. D **94**, 104031 (2016).

[16] M. S. Volkov, "Stationary generalizations for the Bronnikov–Ellis wormhole and for the vacuum ring wormhole," Phys. Rev. D **104**, 124064 (2021).

[17] B. Azad, J. L. Blázquez-Salcedo, F. S. Khoo, and J. Kunz, "Are slowly rotating Ellis–Bronnikov wormholes stable?," Phys. Lett. B **848**, 138349 (2024).

[18] B. Azad, J. L. Blázquez-Salcedo, F. S. Khoo, and J. Kunz, "Radial perturbations of Ellis–Bronnikov wormholes in slow rotation up to second order," Phys. Rev. D **109**, 124051 (2024).

[19] V. Dzhunushaliev, V. Folomeev, B. Kleihaus, J. Kunz, and E. Radu, "Rotating wormholes in five dimensions," Phys. Rev. D **88**, 124028 (2013).

[20] J. L. Blázquez-Salcedo, L. M. González-Romero, F. S. Khoo, J. Kunz, and P. Navarro Moreno, "Radial perturbations of charged wormholes," arXiv:2510.11406 [gr-qc] (2025).

[21] M. Visser, S. Kar, and N. Dadhich, "Traversable wormholes with arbitrarily small energy condition violations," Phys. Rev. Lett. **90**, 201102 (2003).

[22] F. S. Khoo *et al.*, "Quasinormal modes of rapidly rotating Ellis–Bronnikov wormholes," Phys. Rev. D **109**, 084013 (2024).

[23] T. Evstafyeva, N. Siemonsen, and W. E. East, "Assessing the stability of ultracompact spinning boson stars with nonlinear evolutions," arXiv:2508.11527 [gr-qc] (2025).

[24] D. Alic, C. Bona-Casas, C. Bona, L. Rezzolla, and C. Palenzuela, "Conformal and covariant formulation of the Z4 system with constraint-violation damping," Phys. Rev. D **85**, 064040 (2012).

[25] C. Bona, J. Massó, E. Seidel, and J. Stela, "New formalism for numerical relativity," Phys. Rev. Lett. **75**, 600 (1995).

[26] M. Alcubierre, B. Brügmann, P. Diener, M. Koppitz, D. Pollney, E. Seidel, and R. Takahashi, "Gauge conditions for long-term numerical black hole evolutions without excision," Phys. Rev. D **67**, 084023 (2003).

[27] GRTL Collaboration, `GRTeclyn`, https://github.com/GRTLCollaboration/GRTeclyn.

[28] W. Zhang *et al.*, "AMReX: a framework for block-structured adaptive mesh refinement," J. Open Source Softw. **4**, 1370 (2019).

[29] T. Andrade *et al.*, "GRChombo: an adaptable numerical relativity code for fundamental physics," J. Open Source Softw. **6**, 3703 (2021).