Rotation slows the instability of Ellis–Bronnikov wormholes

Nikita M. Shirokov[1, *] and Ilya Nachevsky[2, †]

[1] *Moscow Institute of Physics and Technology, Moscow, Russia*
[2] *Higher School of Economics, Moscow, Russia*

(Dated: October 20, 2026)

We present nonlinear 3+1 evolutions of rotating Ellis–Bronnikov wormholes. We construct stationary rotating drainhole solutions with a spectral solver up to an equatorial throat velocity of $0.99c$, validate the solver against published symmetric solutions, and correct an inconsistency in a published field equation. Perturbed only by truncation error, the rotating throats show that rotation slows the radial instability of the static wormhole. Near 30% of the extremal angular momentum the instability switches from collapse to expansion, rather than disappearing as second-order perturbation theory suggests. At 81% we find no evidence of instability over the evolved time at three resolutions. Collapsing rotating throats emit an $\ell = 2$, $m = 0$ gravitational-wave burst whose energy grows with spin. We also show that second-order perturbative initial data seed the instability at the level of their own truncation error, so they are unsuitable for stability studies.

I. INTRODUCTION

The Ellis–Bronnikov wormhole, a throat held open by a phantom scalar field [1–4], is unstable: depending on the sign of the perturbation it collapses to a black hole or expands without bound, on a timescale set by the throat radius [5–9]. Nonlinear 3+1 evolutions confirm both branches for the lone throat [10] and carry them into binaries, whose every channel ends in collapse, expansion or a merged horizon within a few dynamical times [11]. If traversable wormholes are to persist anywhere, something must shut this instability off.

Rotation is the oldest candidate [12]. Spinning Ellis–Bronnikov solutions are known perturbatively [13] and, beyond perturbation theory, from the elliptic solver of Kleihaus and Kunz [14, 15]: a family that starts at the static throat and ends, at equatorial rotation velocity $v_e = 1$, on an extremal Kerr black hole, the phantom field vanishing in the limit; no closed form exists [16]. On the stability side, Azad, Blázquez-Salcedo, Khoo and Kunz [17, 18] carried the radial mode to second order in spin: the static unstable mode weakens, with a positive $O(J^2)$ correction for every asymmetry, while for nearly symmetric wormholes a second unstable mode grows out of a static zero mode; the two cross near 30% of the limiting angular momentum, where the quadratic approximation gives out, and the authors conjecture the modes merge and the instability disappears, as purely imaginary modes do in five dimensions [19]. Quasinormal modes of the rapidly rotating family have also been computed [22]. The charged analogue warns against optimism: there the merged modes continue as a complex pair, growth that oscillates, and the growth rate reaches zero only in the extremal limit [20]. Which continuation rotation takes in four dimensions is exactly the question a perturbative method cannot answer, and no rotating wormhole has ever been evolved nonlinearly.

This paper builds the tools for that evolution and takes its first step. We construct two families of backgrounds as exact stationary solutions, distinguished by the asymmetry γ between their two ends (defined in Sec. II). The symmetric family ($\gamma = 0$), massless when static, is the one with published data, the curves of Ref. [14] and the scalar charges of Ref. [15], and we use it to validate the solver and the 3D initial data across the spin range. The drainhole family ($\gamma = -\pi$), whose static member is the mass-1 drainhole evolved in Refs. [10, 11], has no published fast-spin data at all; it is the family we evolve. The solver is gated on identities it never imposes, a discipline that pays immediately: one of the published field equations requires a correction (Sec. II A), and second-order perturbative initial data turn out to seed the instability at their own error level (Sec. V). We then put the backgrounds on the adaptive-mesh-refinement (AMR) evolution grid, verify them at fourth order, and evolve the drainhole at nine spins (Table III). We find that rotation slows the collapse, as the perturbative sign predicts; that near 30% of the extremal angular momentum the instability changes its end state, the unperturbed throat turning to expansion; and that at 81% the throat shows no growth over the full run.

Section II gives the family and the corrected field equations; Sec. III the solver, the validation against the published data, and the map of the parameter space (Table II); Sec. IV the numerical setup and the runs (Table III); Sec. V the evolutions; Sec. VI the conclusions. Appendix A gives the convergence of the evolutions at the two ends of the ladder. We use $G = c = 1$; the evolutions use the code units of Refs. [10, 11], with throat parameter $\eta_0 = 2$ (defined in Sec. II), in which the static drainhole has mass $M = 1$.

* shirokov.nm@phystech.edu
† ommnammashivaya@gmail.com

II. THE SPINNING WORMHOLE FAMILY

The matter model is that of our previous work [10, 11]: Einstein gravity coupled to a phantom scalar field ψ,

$$S = \frac{1}{16\pi}\int d^4x\,\sqrt{-g}\,(R + 2\,g^{\mu\nu}\partial_\mu\psi\,\partial_\nu\psi)\,, \tag{1}$$

whose kinetic term has the reversed sign that makes ψ a ghost. Varying the action with respect to the metric and to ψ gives the field equations of the model [Eqs. (2)–(3) of Ref. [15]],

$$R_{\mu\nu} = -2\,\partial_\mu\psi\,\partial_\nu\psi, \qquad \partial_\mu\big(\sqrt{-g}\,g^{\mu\nu}\partial_\nu\psi\big) = 0\,: \tag{2}$$

Einstein's equations with the phantom's stress tensor as source, and the wave equation for the phantom on the curved background. Spinning the wormhole of Refs. [10, 11] up requires the stationary rotating solution of the same theory, constructed numerically inside an ansatz with stationarity and axisymmetry built in. We take the ansatz of Refs. [14, 15], who first constructed these solutions,

$$\begin{aligned} ds^2 = &-e^f dt^2 + p^2 e^{-f}\Big[e^\nu\left(d\eta^2 + h\,d\theta^2\right) \\ &+ h\sin^2\theta\,(d\varphi - \omega\,dt)^2\Big], \end{aligned} \tag{3}$$

with $\eta \in (-\infty, \infty)$ and four unknown functions f, p, ν, ω of η and θ [Eq. (4) of Ref. [15]]. The field equation for p decouples from the others, and its solution with asymptotically flat boundary conditions is $p = 1$ [Eqs. (5)–(6) of Ref. [15]]; we set $p = 1$ from here on. In the trivial limit $f = \nu = \omega = 0$ the spatial metric is $d\eta^2 + h\left(d\theta^2 + \sin^2\theta\,d\varphi^2\right)$ with the fixed $h = \eta^2 + \eta_0^2$: $\sqrt{h}$, the radius of the sphere at coordinate η, never vanishes, bottoming out at η_0 at $\eta = 0$, so a minimal sphere, the throat, joins the two asymptotically flat ends $\eta \to \pm\infty$ and the spacetime has no center. Choosing h hard-wires this two-ended, horizonless topology into Eq. (3); conversely, $\eta_0 \to 0$ closes the throat and lets the same ansatz hold a black hole, which is how extremal Kerr enters in Sec. II A. Our previous wormhole sits inside this form: with $\omega = \nu = 0$ and the closed-form arctan profile for f [15], Eq. (3) is the static drainhole of Refs. [10, 11], with $\eta_0 = a$ its throat parameter and η the isotropic radius used there, unfolded through the throat (Sec. IV A); $f = 0$ gives the massless Ellis wormhole. Substituting the ansatz (3) into the field equations (2) reduces them in three steps. We follow the reduction of Kleihaus and Kunz [14], given in full in Sec. II of Ref. [15]; equation numbers in brackets below refer to Ref. [15]. First, the phantom field equation—the wave equation in Eq. (2), which stationarity and axisymmetry reduce to a function of η alone—integrates in closed form to $\psi = (D/\eta_0)\,[\arctan(\eta/\eta_0) - \pi/2]$ [Eqs. (7)–(9)]: the scalar enters the rest of the problem only through its constant of integration D, the scalar charge. Second, because $\partial_\mu\psi$ points along η, the source term $-2\,\partial_\mu\psi\,\partial_\nu\psi$ survives only in the $\eta\eta$ Einstein equation; each other nontrivial component (tt, $t\varphi$, $\theta\theta$, $\varphi\varphi$, $\eta\theta$) is a statement about the metric functions alone, and these organize by order and role. Two are second-order elliptic equations,

$$\begin{aligned} 0 = &\,\partial_\eta(hs\,\partial_\eta f) + \partial_\theta(s\,\partial_\theta f) \\ &- hs^3 e^{-2f}\left[h(\partial_\eta\omega)^2 + (\partial_\theta\omega)^2\right], \end{aligned} \tag{4}$$

$$\begin{aligned} 0 = &\,\partial_\eta\big(h^2 s^3 e^{-2f}\partial_\eta\omega\big) \\ &+ \partial_\theta\big(hs^3 e^{-2f}\partial_\theta\omega\big)\,, \end{aligned} \tag{5}$$

with $s \equiv \sin\theta$ [Eqs. (10)–(11)]: Eq. (4) is a Laplace-type equation for the redshift f, sourced by the squared gradients of the frame dragging (rotation gravitates), and Eq. (5) is a total divergence, angular-momentum conservation, forcing the same J through every $\eta = \text{const}$ surface. The single off-diagonal component is only first order in ν,

$$\begin{aligned} 0 = &-hs\,\partial_\eta f\,\partial_\theta f + h\cos\theta\,\partial_\eta\nu \\ &+ \eta s\,\partial_\theta\nu + h^2 s^3 e^{-2f}\partial_\eta\omega\,\partial_\theta\omega, \end{aligned} \tag{6}$$

so ν follows by integration once f and ω are known [Eq. (13)]. (The remaining diagonal components give a second-order equation for ν, Sec. II A.) Third, the one sourced component, $R_{\eta\eta} = -2D^2/h^2$ [Eq. (18)], rearranges into an algebraic identity for the charge [Eq. (19)]: the metric functions fix D^2 at every point, so its constancy over the domain is a solution test that comes for free. Three unknown functions thus face more equations than they need; the Bianchi identities make the overdetermined set consistent on a true solution, and a numerical scheme must choose which subset to impose, a choice Sec. II A shows matters. The boundary conditions select the family: Minkowski at $\eta \to +\infty$, and $f \to \gamma$, $\omega \to \omega_\infty$ at $\eta \to -\infty$, so the constant γ measures how unequal the two ends are and ω_∞ sets the spin.

A solution of this system is three functions and one number, $(f, \nu, \omega; D)$. Every physical quantity we quote—the columns of Table II—is read off them directly [14, 15]:

- far from the throat the metric settles to the field of a point mass, $f \to -2\mu_+/\eta$ as $\eta \to +\infty$ and $f \to \gamma + 2\mu_-/\eta$ as $\eta \to -\infty$, so the end masses $\mu_\pm$ are the $1/\eta$ coefficients of f;
- likewise $\omega \to 2J/\eta^3$, and the charges obey the Smarr-type relation $\mu_+ + \mu_- = 2\,\omega_\infty J$ [15];
- the throat is the sphere of minimal area, at $\eta_t = 0$ when the two ends are equal ($\gamma = 0$) and displaced toward the positive-mass end otherwise ($\eta_t = \eta_0/2$ for the static drainhole): its equatorial circumference gives the radius $R_e = e^{-f/2}\sqrt{h}\,\big|_{\eta_t,\,\theta=\pi/2}$, its meridians the polar radius R_p, its area the areal radius R_A—three numbers that coincide while the throat is round and split as rotation flattens it;
- the throat is dragged at the angular velocity $\Omega = \omega\,\big|_{\eta_t,\,\theta=\pi/2}$, and $v_e = R_e\,\Omega$ is its equatorial linear

velocity in units of c: the dimensionless spin label used throughout, 0 when static and 1 at the extremal Kerr endpoint.

We construct two one-parameter families, labelled by ω_∞ (equivalently J or v_e):

(a) *the drainhole family*, $\eta_0 = 2$, $\gamma = -\pi$: its static member, with end masses $\mu_\pm = \pm 1$, is the wormhole of Refs. [10, 11]; its end-mass asymmetry is $C = 1$ in the notation of Refs. [17, 18].

(b) *the symmetric family*, $\gamma = 0$ ($C = 0$), that of Refs. [14, 15]: equal masses on both ends, the massless Ellis wormhole when static; for the 3D checks it is rescaled $\eta_0 = 1 \to 2$, so its throat sits on family (a)'s grid.

A. A correction to the published ν equation

Refs. [14, 15] close the system with a second-order equation for ν, Eq. (12) of Ref. [15]. We find that this equation requires a correction. The scalar-free component $R_{\theta\theta} = 0$, with the second derivatives of f and ω eliminated by Eqs. (4)–(5), gives

$$0 = \partial_\eta(hs\,\partial_\eta\nu) + s\,\partial_\theta^2\nu - \cos\theta\,\partial_\theta\nu + s\,(\partial_\theta f)^2 - hs^3 e^{-2f}\left[h(\partial_\eta\omega)^2 + 2(\partial_\theta\omega)^2\right], \tag{7}$$

where the printed form has $-2(\partial_\theta\omega)^2$ and no $s\,(\partial_\theta f)^2$ term. Extremal Kerr fits the ansatz (3) in closed form (with $\eta_0 = 0$), and substituted symbolically, at 160-digit arithmetic, it satisfies Eqs. (4)–(6), Eq. (7) and the D^2 identity exactly, while leaving a residue of up to 0.35 (at $M = 1$) in the printed ν equation alone. The difference also shows dynamically: solutions of the printed elliptic system converge at every resolution, yet carry a violation of the unimposed Eq. (6) and D^2 identity that scales as J^4 and does not shrink with the grid, the signature of a difference in the continuous equations rather than in the numerics. Our solver imposes neither ν equation (Sec. III), and Eq. (7) holds on all 200 saved solutions of both families to $\leq 7 \times 10^{-8}$ of its largest term within 20 throat radii, where the printed form differs by up to 53%. Every published conclusion of Refs. [14, 15] that we can test survives the correction (Sec. III).

III. BACKGROUNDS ACROSS THE SPIN RANGE

The solver turns the system of Sec. II into numbers in five steps (Fig. 1). First, the infinite domain is made finite: the coordinate $x = (2/\pi)\arctan(\eta/\eta_0)$ maps both asymptotic ends to $x = \pm 1$, and the unknowns f and ω are expanded in Chebyshev polynomials of x and even cosine modes of θ. Second, the two elliptic equations (4)–(5) are solved for the expansion coefficients by Newton iteration (guess, linearize, correct) until their residual sits at round-off. Third, ν and the scalar charge are recovered without a second-order ν equation: the first-order constraint (6) and the D^2 identity together overdetermine ν and the single number D^2, so both are extracted as one linear least-squares problem; the redundancy is a feature, since the leftover misfit measures the error. Fourth, each family is walked in spin: every converged solution seeds the Newton iteration at a slightly higher spin, with J as the continuation parameter (ω_∞ is not monotone along a family), and the grid refines automatically, from 48×12 up to 240×60, whenever the acceptance gates fail. Fifth, and this is the point of the whole design, a solution is accepted only on identities the solver never imposes: the spread of D^2 over the domain ($\leq 10^{-11}$), the Smarr relation ($\leq 10^{-10}$), equality of the angular-momentum flux through the two ends ($\leq 1.3 \times 10^{-9}$), the asymptotic tails ($\leq 5 \times 10^{-12}$), and an independent, symbolically generated residual of the full Einstein equations, evaluated out to 20 throat radii. That residual is 1.0×10^{-10} at $J = 0.25$ and 4.2×10^{-10} at $J = 1$, against the static member's 2×10^{-11} round-off floor; the $O(J^2)$ perturbative data give 3×10^{-5} and 1×10^{-2} on the same meshes. It stays near 10^{-9} up to $v_e \approx 0.85$ and reaches $\sim 10^{-7}$ at $v_e = 0.91$–0.99. The high-spin 10^{-7} is round-off, not solution error: it does not shrink on finer grids, while the global charges agree to 12 digits across them; a deliberately under-resolved solution scores 10^{-4} and is rejected. Beyond 20 throat radii the residual's normalization inflates round-off and the test is switched off; the outer region is covered by the resolution study and the 3D constraint checks of Sec. IV. The products: global charges stable to 10 digits across three grids and, for evolution, each background exported as a 1025×65 table whose bilinear interpolation error is $\leq 10^{-7}$ up to $v_e = 0.9$.

TABLE I. Scalar charge D of the anchor solutions ($\eta_0 = 1$, $\omega_\infty = 0.5$) against Ref. [15]. Along each fixed-γ family, ω_∞ rises, turns over and descends, so $\omega_\infty = 0.5$ is reached twice; the published values sit on the post-turnover branch. Our values are grid-converged to the digits shown; the $\gamma = 2$ family continued to $J = 172$ confirms no third crossing exists. The pre-turnover branch is new.

γ	D (this work)	D [15]	D (pre-turnover, new)
-1	0.716368	0.71636	0.969194
0	0.552771	0.55277	0.986013
1	0.4314654	0.43155	1.008136
2	0.3375993	0.33984	1.048161

Figure 2 validates the symmetric family against the published curves. The four published anchor solutions at $\omega_\infty = 0.5$ [15] split: the $\gamma = 0$ and $\gamma = -1$ scalar charges agree to all printed digits, while $\gamma = 1$ and $\gamma = 2$ differ by 2×10^{-4} and 6.6×10^{-3} from our grid-converged values, consistent with their finite-difference error growing with




FIG. 1. The solver loop for one family (Sec. III). Thin boxes impose equations; the heavy box tests identities the solver never imposes, and only it decides acceptance. A failed check refines the grid and solves the same J again; an accepted solution seeds the next J. The loop gave 462 accepted solutions in 20 continuation sweeps, both families to $v_e \approx 0.99$.




FIG. 2. The symmetric family ($\eta_0 = 1$, $\gamma = 0$; 81 solutions, $v_e = 0.02$–0.991, plus the exact static member) in the normalization of Ref. [14] ($M_0 = R_e$, $J_0 = R_e^2/2$, $\Omega_0 = 1/R_e$): our curves, the digitized points of their Fig. 1 (open symbols), and closed-form Kerr (dashed). The digitization's reading error, measured by reading their printed Kerr curves the same way, is rms 1–6 $\times 10^{-4}$; digitized minus ours is rms 2.5–8 $\times 10^{-4}$, max 2.5×10^{-3}, over all five curves. The two solvers agree to the precision a printed figure carries.

γ (Table I).

Figure 3 and Table II are the map of the parameter space that Refs. [14, 15] left open, including the massive drainhole family, from which binaries can be built [11]. Two of its features matter later. First, both families violate the Kerr bound (Fig. 3b), the symmetric family with $J/\mu_+^2 \approx 85$ already at $v_e = 0.24$, falling monotonically to 1 at extremality since its static member is massless. A collapsing overspun throat cannot settle to a Kerr black hole of the same mass and spin, which makes these backgrounds sharp cosmic-censorship initial data. Second, the drainhole's negative far-end mass μ_- rises through zero at $v_e \approx 0.7$ and turns positive: fast rotation undoes the static drainhole's negative-mass end.

IV. NUMERICAL SETUP

A. 3D initial data

The stationary metric (3) goes onto the Cartesian AMR grid through the isotropic radius r, $\eta = r - \eta_0^2/4r$: the far end $\eta \to -\infty$ compactifies to the puncture $r = 0$, as for the static drainhole [10]. The induced metric and the fields are read from the tables; the lapse and shift start from the stationary data, $\alpha = e^{f/2}$, $\beta^\varphi = -\omega$, so the extrinsic curvature $K_{ij} = (D_i\beta_j + D_j\beta_i)/2\alpha$ has only $r\varphi$ and $\theta\varphi$ components and the scalar momentum vanishes; at the puncture the frame rotates at ω_∞.

Three checks validate the pipeline. (i) At $t = 0$ the Hamiltonian and momentum constraints converge at fourth order inside the throat region and are limited outside only by the tables' interpolation error (growing as $1/\Delta x^2$ and $1/\Delta x$ respectively), and the throat's ray radii converge to the solver's. We check this on the drainhole at $v_e = 0.51$ on three grids (ray radii within 5×10^{-5} at $N = 192$, $L = 64$) and, across the spin range, on the symmetric family at seven spins $v_e = 0.24$–0.99 with three resolutions each (to 9×10^{-4} at worst, $N = 256$). (ii) On the same symmetric set, the 3D ADM mass integral evaluated on the grid equals the same integral on the exact table metric to $\leq 10^{-12}$, and its Richardson extrapolation meets the solver's mass to at worst 0.7% relative (at $v_e = 0.99$); the ADM angular momentum matches to 2×10^{-3} or better. (iii) Dynamically, evolving the static drainhole through this pipeline reproduces the evolution of Ref. [11] over the full record: the $t = 0$ data are bit-identical, the Hamiltonian norm stays within 5×10^{-7} for all $t \leq 100$, and the collapse milestones agree within one time unit.




FIG. 3. The drainhole ($\eta_0 = 2$, $\gamma = -\pi$) and symmetric ($\eta_0 = 1$, $\gamma = 0$) families across the spin range, against v_e: (a) mass, angular momentum and equatorial throat radius; (b) J/M^2 against the Kerr bound; (c) the ratio of end masses; (d) the flattening R_p/R_e; (e) the exotic matter, the phantom field's integrated null-energy violation $\mathcal{I} = \int(\rho + p_\eta)\, dV$ over the $t = \text{const}$ slice [21] (ρ: energy density, p_η: radial pressure), per equatorial throat radius, with the 3D runs (squares) placed by the scalar charge measured on each run's own $t = 0$ slice, which reproduces its background's to 3×10^{-4}; (f) the throat angular velocity, which peaks and falls to zero at extremality. The symmetric family violates the Kerr bound at every spin and the drainhole family exceeds it by up to $\sim 20\%$ at intermediate spin; the exotic matter per throat radius falls a hundredfold along the drainhole family and tenfold along the symmetric one by $v_e = 0.99$; both families end on extremal Kerr.

TABLE II. The two families at representative spins, in units of η_0: rotation velocity v_e, angular momentum, mass of the $\eta \to +\infty$ end, ratio of end masses, scalar charge, equatorial throat radius, flattening, and throat angular velocity—each read off the solution's functions as defined in Sec. II. Every row is a converged solution that passed the acceptance gates of Sec. III.

family	v_e	J/η_0^2	μ_+/η_0	μ_-/μ_+	D/η_0	R_e/η_0	R_p/R_e	$\Omega\,\eta_0$
(a) drainhole	0.213	0.2535	0.5574	−0.850	1.1004	2.0319	0.9873	0.1046
	0.513	1.0278	0.9302	−0.339	1.0007	2.6271	0.9194	0.1952
	0.699	3.0417	1.6557	−0.032	0.8598	3.8864	0.8351	0.1798
	0.900	34.550	5.8400	+0.179	0.5327	11.911	0.6926	0.0755
	0.950	152.16	12.316	+0.201	0.3785	24.751	0.6504	0.0384
	0.990	4053.8	63.666	+0.208	0.1698	127.36	0.6165	0.0078
(b) symmetric	0.239	0.0542	0.0252	1	0.9881	1.0299	0.9913	0.2323
	0.434	0.1158	0.0906	1	0.9583	1.1101	0.9699	0.3911
	0.667	0.2683	0.2666	1	0.8854	1.3420	0.9197	0.4969
	0.847	0.7007	0.6305	1	0.7653	1.8836	0.8440	0.4499
	0.898	1.0989	0.8685	1	0.7043	2.2722	0.8092	0.3952
	0.952	2.4701	1.4383	1	0.5971	3.2703	0.7539	0.2911
	0.990	12.235	3.4312	1	0.4157	7.0595	0.6795	0.1402

B. Evolution scheme

Evolution uses `GRTeclyn` [27–29] with the setup of Refs. [10, 11]: the CCZ4 formulation of the Einstein equations [24], 1+log slicing [25], Gamma-driver shift [26], fourth-order stencils, fourth-order Runge–Kutta time integration with $\Delta t = 0.02\,\Delta x$ in every run, Kreiss–Oliger dissipation of strength σ, and a floored conformal factor at the puncture. The only scheme parameter that varies

TABLE III. The simulations: the drainhole family (a), in code units. Model Dn is the drainhole background with $v_e \approx n/100$ (D0: static). $8\pi J/A$ is the angular momentum as a percentage of its extremal value $J_{\rm ext} = A/8\pi$ at the throat area A, and $M_{\rm ADM}$ the mass of the $\eta \to +\infty$ end (both from the background tables). The finest Δx is 0.0625 at $N = 128$ on $L = 64$ and at $N = 256$ on $L = 128$, and scales inversely with N within a triplet. σ is the Kreiss–Oliger dissipation strength (Sec. IV B); the conformal factor is floored at 10^{-8}, except at 10^{-4} in the D90 runs. Damping: the boundary damping layer of Sec. IV C on, or the background-relative boundary alone (off). Conv. t: the time up to which the results are converged and quoted. τ: the e-fold time of the unstable mode (Sec. V); MOTS: the marginally outer-trapped surface (Sec. IV D). GW: the $(2,0)$ burst at $R = 14$ arrives within the converged span (burst), is still rising at its end (onset), or stays below the extraction floor (none). The D59 and D70 runs are still in progress.

model	v_e	$8\pi J/A$	$M_{\rm ADM}$	L	N	σ	damping	final t	conv. t	τ	GW	end state
D0	0	0	1.000	64	128	0.1	off	100	60	5.4	none	collapses; dies inside its horizon, $t = 62.7$
D21	0.210	12.3	1.112	64	96/128/192	0.1	on	100	55/100/100	6.6	burst	collapses; horizon settles at $R = 2.52$
D21	0.210	12.3	1.112	64	128	0.1	off	100	42/61	–	onset	perturbation $\lvert\varepsilon\rvert = 10^{-3}$: outward collapses, inward expands
D21	0.210	12.3	1.112	128	256	0.1	off	150	150	–	burst	collapses; the $(2,0)$ burst on spheres $R = 14$–45 (Fig. 5)
D30	0.304	18.3	1.248	64	128	0.1	off	100, 130	130	8.3	burst	collapses; horizon at $R = 2.567$
D40	0.395	24.4	1.448	64	128	0.1	off	100	100	13.3	onset	collapses; MOTS leaves the throat, $t = 92$
D46	0.456	28.8	1.634	64	128	0.1	off	100, 150	100, 121	12.1	onset	-1.1% at $t = 83$, expands from $t = 101$
D51	0.513	33.3	1.860	64	192	0.1	on	100	100	–	none	shrinks, -1% at $t = 99$: the opposite perturbation sign
D51	0.513	33.3	1.860	64	128	0.1	off	150	95	8.8	none	expands from $t \approx 70$
D59	0.592	40.3	2.305	128	256	0.4	on	150	–			in progress
D70	0.699	51.4	3.311	128	256	0.4	on	150	–			in progress
D90	0.900	80.8	11.68	128	128/192/256	0.4	on	150	150	–	none	no growth: areal drift $\leq 7 \times 10^{-5}$ at all three N

between runs is σ (Table III): the larger value suppresses a grid-scale mode that grows at the fixed mesh-refinement interfaces (Appendix B). Its one measured side effect, a constant advance of the collapse clock by about 4 time units from the stronger initial smoothing, keeps rate comparisons at matched σ.

C. Long evolutions: the outer boundary

Evolutions that must outlive the outer-boundary crossing time $t \approx L$ need the boundary made harmless. Kreiss–Oliger dissipation cannot do it, since its strength falls as the sixth power of wavelength, so the sponge layer of Refs. [10, 11] is transparent to the long outgoing gauge pulse; and a flat-space Sommerfeld condition assumes asymptotics the wormhole does not have (mass 1–12 sits inside the box). Two changes in the same outer layer fix it. The Sommerfeld condition is imposed on the deviation of the evolved fields from the stationary background rather than on the fields themselves; and a damping layer can in addition drive every evolved field back to that background at rate λ, $\partial_t u \to \partial_t u - \lambda\, f(r)\,(u - u_{\rm bg})$, with $f(r)$ the sponge's quartic ramp and $u_{\rm bg}$ the run's own initial data, rebuilt exactly from the background tables; $\lambda = 0$ recovers the base scheme identically. The layer is validated at $\lambda = 1$ on the static drainhole against the reference evolution: the trace K of the extrinsic curvature falls two orders of magnitude across the layer, while everything inside it (the throat radii to four digits, the constraint norms, the near-zone ring) is unchanged, so it absorbs at all wavelengths without touching the interior. On the $L = 64$ box, whose layer ($r = 24$–32) contains the outermost extraction sphere, a side-by-side test showed the background-relative boundary alone keeps the outer levels cleaner, so those runs take the damping off; the $L = 128$ boxes ($r = 48$–64, outside every sphere) run with it on (Table III). With the boundary treated this way the $L = 64$ runs, previously reliable only to the crossing time $t \approx 64$, run to their final times $t = 100$–150, with the converged span set by the constraint norms instead.

D. The runs

Stability claims follow the criteria of Sec. II of Ref. [23]: a gauge-invariant measure of the growth, for which we use the throat's areal radius and the marginally outer-trapped surface (MOTS) found spectrally on every output; constraint norms that converge at the code's order while the growth proceeds; an e-fold time that converges to a nonzero value across resolutions; and a varied perturbation amplitude, which we impose at $v_e = 0.21$ (the $\varepsilon = \pm 10^{-3}$ runs of Table III). Every run carries three nested refinement levels on the throat, and the box is picked by the model's mass: $M_{\rm ADM}$ triples along the family (1.112 at $v_e = 0.21$, 3.311 at 0.70), so the face of the $L = 64$ box recedes from $25\,M_{\rm ADM}$ of clearance




FIG. 4. The inverse e-fold time $1/\tau$ of the throat's areal radius (left axis; τ on the right) against the angular momentum as a fraction of its extremal value. Filled circles: collapsing models; filled triangles: expanding models; open circle: the stable D90; every τ is the offset-free fit of Sec. V and a row of Table III. Dashed: the fit $a - b\,(J/A)^2$ through the collapse branch alone, zero at 32%. Open diamonds: the second-order rates of Refs. [17, 18] at $C = 1$; dotted rules: their conjectured end of the instability (≈30%) and the zero of their second-order rate (33%). The collapse branch dies where second-order theory predicts, and past the crossing the unperturbed throat expands with a rate that rises with spin.

at slow spin to under $10\,M_{\rm ADM}$ at $v_e = 0.70$. Models up to $v_e = 0.51$ therefore run on the $L = 64$, $N = 128$ grid of the static evolution of Ref. [11], and models from $v_e = 0.59$ on $L = 128$, $N = 256$ at the same finest Δx. The Newman–Penrose scalar Ψ_4, whose multipoles carry the outgoing gravitational wave, is extracted on spheres at $R = 10$–30 ($L = 64$) and $R = 14$–45 ($L = 128$); the signal is its $(\ell, m) = (2, 0)$ mode. Resolution triplets anchor the two ends of the spin sequence (Appendix A). Table III lists every run, from the static reference through the $O(J^2)$ regime [17, 18] to far beyond any perturbative result.

V. THE INSTABILITY ACROSS THE SPIN RANGE

Perturbative initial data. We first evolved the second-order $O(J^2)$ backgrounds of Ref. [18] at $J = 0.25$ and $J = 1$ on the drainhole family. Both throats expanded where the static throat collapses: the same unstable mode with the opposite sign ($\tau = 5.8$ and 5.6). Expressed as the equivalent radial perturbation amplitude of Ref. [11], however, the push was $\varepsilon \approx -3 \times 10^{-5}$ at $J = 0.25$ and -7×10^{-3} at $J = 1$: growth by a factor ∼230 for a factor 4 in J, i.e. $\sim J^4$, the size of the $O(J^2)$ data's own truncation error, which the independent Einstein residual puts at 10^{-2} at $J = 1$ (Sec. III). Second-order data seed the very instability under study at their own error level; nothing at slow spin can be concluded from them.

Measuring the growth. The diagnostic for every unperturbed evolution is the throat's areal radius $R_A(t)$, the size of the $\ell = 0$ mode the perturbative work computes. The e-fold time is an offset-free fit: the slope of $\ln|dR_A/dt|$ over the span where the areal deficit $1 - R_A(t)/R_A(0)$ (collapse) or growth (expansion) first passes 0.2–1%. Fitting the rate of change rather than the deficit itself removes the initial transient's baseline, which biases a direct log fit of the deficit by a resolution-dependent amount. The band sits on a plateau: the measured τ moves by under 4% between the 0.2–1% and 0.5–2% bands. So measured, τ self-converges across the D21 resolution triplet, 7.8, 6.6 and 6.1 at $N = 96$, 128 and 192, at third to fourth order ($Q = 2.3$; Appendix A), with a continuum value $\tau = 5.5$–5.7. The finite-resolution bias is close to a common factor on every rate, so it cancels in the zero of the fit below.

The collapse branch. On exact data (Table III) the behaviour reverses, and the unperturbed throat collapses on the static mode, later the faster it spins: the areal deficit passes 1% at $t = 43.7$, 46.9, 51.4 and 60.2 along the models D0–D40, with e-fold times $\tau = 5.4$, 6.6, 8.3, 13.3 (Fig. 4). Each collapse ends in a black hole: the D21 horizon settles at $R = 2.524$ (Misner–Sharp mass $M_{\rm MS} = 1.26$), the D30 at $R = 2.567$ by $t = 130$; the D40's MOTS leaves the throat at $t = 92$ and reads $R = 3.22$, still shrinking, at $t = 100$. In each case the evolution is seeded only by discretization error, so the ordering of rates is the ordering of the unstable mode's growth rate.

The expanding branch. At $v_e = 0.456$ (28.8% of extremal) the end state changes: the areal radius dips to -1.12% at $t = 83$, turns, crosses zero at $t = 101$ and grows exponentially, $\tau = 12.1$. The D51 model behaves the same way (-0.30% at $t = 65$–68, then exponential growth with $\tau = 8.8$), and the MOTS is lost once a throat expands: no trapped surface forms. Collapse and expansion are the instability's two end states, and the sign of the initial perturbation selects between them, exactly as the imposed radial perturbations at D21 show (pushed inward, the throat expands; pushed outward, it collapses; both pass 1% at $t \approx 25$ at the same rate): the D51 run whose absorbing layer gives the opposite effective perturbation shrinks instead (-1% at $t = 99$). At the fast end, the D90 throat radius shows no growth up to $t = 150$, about 28 e-folding times of the static mode, at all three resolutions (Appendix A).

Against the perturbative prediction. Figure 4 puts both branches on one axis. Where second order should hold, the two methods agree: the D21 continuum rate matches the second-order $C = 1$ value of Refs. [17, 18] to 5% ($\tau = 5.5$–5.7 against 5.46), and the static member lies beside its linear mode (5.4 at $N = 128$, extrapolating toward 5.1). At intermediate spin the measured mode is slower than their second-order values by more than the resolution bias ($\tau = 8.3$ against 6.1 at $v_e = 0.30$, 13.3 against 8.0 at 0.40), a gap that grows with spin as the neglected $O(J^4)$ terms must. The collapse branch itself




FIG. 5. The $(2,0)$ wave of the unperturbed rotating collapse. (a) $r\,\Delta\psi_4^{20}$ of the D21 run on the $L = 128$ box, on spheres $R = 14$–45 against retarded time $t - R$; only the first lobe falls with R (near field). (b) The wave against spin at $R = 14$, each run on its own collapse clock $u = t - R - t_{10}$ (t_{10}: the time its equatorial deficit passes 10%); the static run (D0) is drawn at zero, its measured record below 3×10^{-8}. (c) The radiated energy $E_{20} = \frac{1}{16\pi}\int |\int r\psi_4\, dt'|^2\, dt$ over the aligned burst, per unit $M_{\rm ADM}$, at $R = 18$ (filled) and $R = 14$ (open). (d) The speed of each extremum between neighbouring spheres, against the local coordinate speed of light (crosses). The burst propagates at light speed and its energy grows with spin as the amplitude squared, while the static collapse radiates nothing.

follows $a - b\,(J/A)^2$ to rms 2×10^{-3} in $1/\tau$ and reaches zero at 32% of the extremal angular momentum, beside the zero of their second-order rate (33%) and their conjectured mode crossing (≈30%). The instability nevertheless does not end there: the 28.8% and 33% models expand with rates that rise with spin ($\tau = 12.1$, then 8.8), consistent with their second unstable mode taking over rather than the two modes merging away. Their τ is a linear mode in Killing time, ours a nonlinear areal-radius fit, so the pointwise comparison is indicative.

Gravitational waves. The rotating collapse radiates with no nonspherical perturbation put in (Fig. 5): a single $(2,0)$ burst with ring-down period ≈24, propagating at the local light speed, its extrema flat in extraction radius to 1.5% over $R = 30$–45. Aligned on each model's own collapse, the D30 burst has the D21's shape and timing at about 1.5 times the amplitude, and the radiated energy per unit mass grows as the amplitude squared, $E_{20}/M_{\rm ADM} = 3.3 \times 10^{-7} \to 8.0 \times 10^{-7}$. The spherical static collapse radiates nothing: its measured record stays below 3×10^{-8}. Rotation thus imprints itself on the burst a collapsing throat emits, before any horizon forms. (The $L = 128$ box reproduces the $L = 64$ throat history to 3×10^{-4}.)

VI. CONCLUSIONS

First, the spinning Ellis–Bronnikov parameter space is now mapped as data, not curves: both families from slow rotation to $v_e \approx 0.99$, including the drainhole family for which nothing was published, with the published ν equation, Eq. (12) of Ref. [15], corrected to Eq. (7). The practical lesson generalizes: a solver gated only on the equations it imposes would have passed every test while solving a different system; the unimposed identities caught the difference. Second, the pipeline from these solutions to 3+1 evolution is validated at fourth order, and $O(J^2)$ initial data are shown to be unsuitable for stability work: their truncation error seeds the instability itself. Third, these first nonlinear evolutions of rotating wormholes show the instability slowing and then changing branch: the radial collapse slows with spin ($\tau = 5.4 \to 13.3$ from the static model to 24% of the extremal angular momentum), its rate following $a - b\,(J/A)^2$ to a zero at 32%, beside the second-order prediction of 33% [17, 18]; near 30%, where the perturbative mode crossing was conjectured to end the instability, the unperturbed throat expands instead, with a rate that rises with spin; and at 81% the throat radius shows no growth up to $t = 150$, about 28 e-folding times of the static mode, at three resolutions, addressing, for this family and timescale, the four-dimensional stability question left open in Ref. [14]. The rotating collapse also radiates a $(2,0)$ burst from unperturbed, axisymmetric data, its energy growing with spin as the amplitude squared, while the static collapse radiates nothing.

VII. REPRODUCIBILITY AND COST

The solver, the evolution code and the figure scripts are in the `GRTeclyn` [27] fork at https://github.com/Nikchik-coder/GRTeclyn (branch `feature/spinning`), whose `results/spinningwormhole` pack holds the solver's accepted solutions, the background-table headers the evolutions start from, and the finished runs' parameter files and diagnostic streams, with a run registry.

Every evolution ran on one NVIDIA H100 80 GB device, with `GRTeclyn` built CUDA-native against AM-

ReX 26.02. A run on the $L = 64$, $N = 128$ grid takes 7.6–11.1 h to $t = 100$; the $N = 192$ members 20 h, peaking at 53 GiB of the device; the $L = 128$, $N = 256$ runs 9.7 h to $t = 150$. The 17 finished evolutions of Table III total 167 GPU-hours, recomputed from their archived logs.

ACKNOWLEDGMENTS

This work was supported by Gravity Frontiers (https://www.gravityfrontiers.org/en), which funded the underlying research and the development of the numerical-simulation software. The authors acknowledge the GRTL Collaboration for `GRChombo`, `GRTresna`, and `GRTeclyn`.

Appendix A: Convergence of the evolutions

The two resolution triplets are D21, the slowest spinning collapse (Fig. 6), and D90, the stable fast model (Fig. 7), both read through the per-level constraint norms of Appendix B. Rescaling constraint norms between resolutions is a valid convergence test because their exact value is zero. For the D21 triplet's grid spacings ($h \propto 1/N$, $N = 96/128/192$) the expected self-convergence factors $Q_n = (h_1^n - h_2^n)/(h_2^n - h_3^n)$ are $Q_3 = 1.95$ and $Q_4 = 2.69$; the offset-free growth rate of Sec. V measures $Q = 2.3$, converging at the same order as the constraints.




FIG. 6. Convergence of the D21 model over its triplet $N = 96/128/192$ ($L = 64$; finest Δx in each panel title). (a)–(d) Hamiltonian and (e)–(h) momentum constraint norms per refinement level; dashed and dotted, the $N = 192$ norms rescaled by 2^2 and 2^4: the measured $N = 96$ norms lie between the second- and fourth-order brackets on every level while the collapse grows. (i) The equatorial throat radius against the stationary solution, falling with resolution. (j) The collapse, each run aligned at its own 10% time: the three resolutions land on one curve. (k) $r\,\Delta\psi_4^{20}$ at $R = 14$, aligned: $N = 128$ and 192 overlap ($N = 96$ dies in its collapse before the burst arrives). (l) The MOTS radius, aligned: one horizon, settling at $R = 2.52$. The collapse rate, the waveform and the horizon are grid-converged at third to fourth order.




FIG. 7. Stability of the D90 model over its triplet $N = 128/192/256$ ($L = 128$; finest Δx in each panel title), to $t = 150$. (a)–(h) Constraint norms per level with the same second-/fourth-order brackets as Fig. 6: every level converges with resolution and nothing breaks out over the run. (i) The equatorial throat radius holds the stationary value, its offset falling with resolution. (j) The areal throat radius drifts by $\leq 7 \times 10^{-5}$ over the full run. (k) The $(2, 0)$ extraction at $R = 18$, relative to its $t = 0$ read-off: fluctuations at the 10^{-4} level, no burst. (l) The lapse at the $\eta \to -\infty$ puncture falls smoothly at all three resolutions (the gauge, not the geometry). The D90 throat shows no growing radius, no constraint breakout and no wave at any resolution over the run.

Appendix B: The refinement-interface mode and the timestep

The timestep $\Delta t = 0.02\,\Delta x$ was adopted empirically in Refs. [10, 11]. We traced what sets it to a grid-scale mode pinned to the fixed coarse–fine interfaces of the nested refinement boxes, dissipation-starved at $\sigma = 0.1$, whose growth rate rises with the timestep. Constraint norms evaluated per refinement level, with cells covered by finer levels masked, expose the mode by $t \approx 10$; the domain-wide norm is blind to it until breakout. Raising the dissipation where the mode lives removes it: at $\sigma = 0.4$ the interface-level norms sit 10^2–10^4 below the $\sigma = 0.1$ reference by $t = 100$. The larger timesteps this would permit do not transfer to the spinning backgrounds, which lose the $\eta \to -\infty$ region (the puncture) by $t \approx 22$–48 at $\Delta t = 0.035$–$0.05\,\Delta x$, so the production runs take no speed-up.

[1] H. G. Ellis, "Ether flow through a drainhole: A particle model in general relativity," J. Math. Phys. **14**, 104 (1973).

[2] K. A. Bronnikov, "Scalar-tensor theory and scalar charge," Acta Phys. Polon. B **4**, 251 (1973).

[3] M. S. Morris and K. S. Thorne, "Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity," Am. J. Phys. **56**, 395 (1988).

[4] M. Visser, *Lorentzian Wormholes: From Einstein to Hawking* (AIP Press, Woodbury, NY, 1995).

[5] J. A. González, F. S. Guzmán, and O. Sarbach, "Instability of wormholes supported by a ghost scalar field. I. Linear stability analysis," Class. Quantum Grav. **26**, 015010 (2009).

[6] J. A. González, F. S. Guzmán, and O. Sarbach, "Instability of wormholes supported by a ghost scalar field. II. Nonlinear evolution," Class. Quantum Grav. **26**, 015011 (2009).

[7] H. Shinkai and S. A. Hayward, "Fate of the first traversible wormhole: Black-hole collapse or inflationary expansion," Phys. Rev. D **66**, 044005 (2002).

[8] K. A. Bronnikov, J. C. Fabris, and A. Zhidenko, "On the stability of scalar-vacuum space-times," Eur. Phys. J. C **71**, 1791 (2011).

[9] F. Cremona, F. Pirotta, and L. Pizzocchero, "On the linear instability of the Ellis–Bronnikov–Morris–Thorne wormhole," Gen. Relativ. Gravit. **51**, 19 (2019).

[10] N. M. Shirokov, "Wormhole dynamics: nonlinear collapse and gravitational-wave emission," arXiv:2604.00071 [gr-qc] (2026).

[11] N. M. Shirokov and I. Nachevsky, "Binary Ellis–Bronnikov wormholes in 3+1 numerical relativity: mergers, fly-bys and gravitational waves," arXiv:2610.13206 [gr-qc] (2026).

[12] T. Matos and D. Núñez, "Rotating scalar field wormhole," Class. Quantum Grav. **23**, 4485 (2006).

[13] P. E. Kashargin and S. V. Sushkov, "Slowly rotating wormholes: The first order approximation," Grav. Cosmol. **14**, 80 (2008); "Slowly rotating scalar field wormholes: The second order approximation," Phys. Rev. D **78**, 064071 (2008).

[14] B. Kleihaus and J. Kunz, "Rotating Ellis wormholes in four dimensions," Phys. Rev. D **90**, 121503(R) (2014).

[15] X. Y. Chew, B. Kleihaus, and J. Kunz, "Geometry of spinning Ellis wormholes," Phys. Rev. D **94**, 104031 (2016).

[16] M. S. Volkov, "Stationary generalizations for the Bronnikov–Ellis wormhole and for the vacuum ring wormhole," Phys. Rev. D **104**, 124064 (2021).

[17] B. Azad, J. L. Blázquez-Salcedo, F. S. Khoo, and J. Kunz, "Are slowly rotating Ellis–Bronnikov wormholes stable?," Phys. Lett. B **848**, 138349 (2024).

[18] B. Azad, J. L. Blázquez-Salcedo, F. S. Khoo, and J. Kunz, "Radial perturbations of Ellis–Bronnikov wormholes in slow rotation up to second order," Phys. Rev. D **109**, 124051 (2024).

[19] V. Dzhunushaliev, V. Folomeev, B. Kleihaus, J. Kunz, and E. Radu, "Rotating wormholes in five dimensions," Phys. Rev. D **88**, 124028 (2013).

[20] J. L. Blázquez-Salcedo, L. M. González-Romero, F. S. Khoo, J. Kunz, and P. Navarro Moreno, "Radial perturbations of charged wormholes," arXiv:2510.11406 [gr-qc] (2025).

[21] M. Visser, S. Kar, and N. Dadhich, "Traversable wormholes with arbitrarily small energy condition violations," Phys. Rev. Lett. **90**, 201102 (2003).

[22] F. S. Khoo *et al.*, "Quasinormal modes of rapidly rotating Ellis–Bronnikov wormholes," Phys. Rev. D **109**, 084013 (2024).

[23] T. Evstafyeva, N. Siemonsen, and W. E. East, "Assessing the stability of ultracompact spinning boson stars with nonlinear evolutions," arXiv:2508.11527 [gr-qc] (2025).

[24] D. Alic, C. Bona-Casas, C. Bona, L. Rezzolla, and C. Palenzuela, "Conformal and covariant formulation of the Z4 system with constraint-violation damping," Phys. Rev. D **85**, 064040 (2012).

[25] C. Bona, J. Massó, E. Seidel, and J. Stela, "New formalism for numerical relativity," Phys. Rev. Lett. **75**, 600 (1995).

[26] M. Alcubierre, B. Brügmann, P. Diener, M. Koppitz, D. Pollney, E. Seidel, and R. Takahashi, "Gauge conditions for long-term numerical black hole evolutions without excision," Phys. Rev. D **67**, 084023 (2003).

[27] GRTL Collaboration, `GRTeclyn`, https://github.com/GRTLCollaboration/GRTeclyn.

[28] W. Zhang *et al.*, "AMReX: a framework for block-structured adaptive mesh refinement," J. Open Source Softw. **4**, 1370 (2019).

[29] T. Andrade *et al.*, "GRChombo: an adaptable numerical relativity code for fundamental physics," J. Open Source Softw. **6**, 3703 (2021).