


Central Line Operating Room Checklist

Pre-Operative	Administer periop ancef 30 min prior to incision (Clindamycin if PCN allg)	
	Bair Hugger off prior to prep	
	Alcohol skin cleanse prior to prep	
	Formal prep with Chlorohexadine; drape at 3 minutes	
	Ioban/Steri Drapes for secure sterile field	
Intra-Operative	Line opened at time of readiness for insertion	
	US guidance used for access	
	RIJ access attempted (exceptions documented)	
	Fluoroscopic image saved	
	CVL secured to skin with suture	
	Dress CVL exit site with Statseal Disc	
	Line Packed with Heparin per Guidelines (below)	
Post-Operative	CVL and PAC dressed as shown below	
	CXR in PACU if concerns for complex placement (document reason)	

Flush protocol

- Flush central line with 3cc of 10U/ml Heparinized saline, 2.5cc per lumen for DLCL (total 5cc)
- Flush accessed port with 5cc of 10U/ml Heparinized saline
- Flush unaccessed port with 5cc of 100U/ml Heparinized saline (for children under 1 year of age, flush with 5cc of 10U/ml heparinized saline)



Angiodynamics Port (all detachable)	Right internal jugular vein size
Mini port, 5Fr	3.3 - 4.3mm
Low profile, 6Fr*	≥ 4.4mm
Standard profile, 8Fr	≥ 5.3mm

CVL type (all cuffed lines)	Right internal jugular vein size
SLCL, Bard Broviac 4.2Fr (white)	2.8 – 4.3mm
SLCL, Bard Broviac 6.6Fr (white)*	≥ 4.4mm
DLCL, Bard Hickman 6Fr (purple)	4 – 4.5mm
DLCL, Bard Hickman 7Fr (white)*	≥ 4.6mm