

Index

Unit – 5 \Rightarrow Applied Statistics	3
1) Method – 1 \Rightarrow Basic Concepts of Tests of Significance	5
Test for Variables for Large Samples	9
2) Method – 2 \Rightarrow Test of Significance of Mean.....	9
3) Method – 3 \Rightarrow Test of Significance of Difference Between Two Means	11
4) Method – 4 \Rightarrow Test of Significance of Difference Between Two Standard Deviations	14
Tests of Significance for Small Samples	18
5) Method – 5 \Rightarrow Test of Significance of Mean.....	18
6) Method – 6 \Rightarrow Test of Significance of Difference Between Two Means	20
7) Method – 7 \Rightarrow F-test for Equality of Two Population Variances	23
8) Method – 8 \Rightarrow Chi-square Test: for Goodness of Fit	28

Unit – 5 \rightsquigarrow Applied Statistics

Introduction

- Many problems in engineering required to decide which of two competing claims for statements about parameter is true. Statements are known as **hypothesis**, and the decision-making procedure is known as **hypothesis testing**.
- This is one of the most useful aspects of statistical inference, because many types of decision-making problems, tests or experiments in the engineering world can be formulated as hypothesis testing problems.

Population OR Universe

- An aggregate of objects under study is known as **Population OR Universe**.
- It is a collection of individuals or of their attributes (qualities) or of results of operations which can be numerically specified.
- A universe containing a finite number of individuals or members is known as a **finite universe**.
 - For Example: The universe of the weights of students in a particular class or the universe of smokes in Rothay district.
- A universe with infinite number of members is known as an **infinite universe**.
 - For Example: The universe of pressures at various points in the atmosphere.
- In some cases, we may be even ignorant whether or not a particular universe is infinite, e.g., the universe of stars.
- The collection of all possible ways in which a specified event can happen is known as a **hypothetical universe**.
 - For Example: The universe of heads and tails obtained by tossing an infinite number of times is a hypothetical universe.

Sampling

- A finite subset of a universe or population is known as a **sample**.
- A sample is a small portion of the universe.
- The number of individuals in a sample is known as **sample size**.
- Sample size is denoted by "**n**" and population size is denoted by "**N**".

Unit - 5 Applied Statistics

- If sample size $n \geq 30$, then it is known as **Large Sample**, otherwise it known as **Small Sample**.
- The process of selecting a sample from a universe is known as **sampling**.
- The theory of sampling is a study of relationship between a population and samples drawn from the population.
- The fundamental object of sampling is to get as much information as possible of the whole universe by examining only a part of it.
- Sampling is quite often used in our day-to-day practical life.

For example: in a shop we assess the quality of sugar, rice or any commodity by taking only a handful of it from the bag and then decide whether to purchase it or not.

Parameter and Statistics

- The statistical constants of **population** are known as **parameters**.
- The statistical constants of **sample** are known as **statistics**.
- Notations for statistical constants are as below:

Statistical Constants	For Sample	For Population
No. of elements (Size)	n	N
Proportion	p	P
Mean	\bar{x}	μ
Standard Deviation	s	σ
Variance	s^2	σ^2
Correlation Coefficient	r	ρ

Method – 1 \Rightarrow Basic Concepts of Tests of Significance

Test of Significance

- An important aspect of the sampling theory is to study the test of significance, which will enable us to decide, on the basis of the results of the sample.
- For applying the tests of significance, we first set up a hypothesis which is a definite statement about the population parameter known as **null hypothesis**, denoted by H_0 .
- Note that, Null hypothesis H_0 is always in equality.
- Any hypothesis which is complementary to the null hypothesis is known as an **alternative hypothesis** denoted by H_1 .
- For example, if we want to test the null hypothesis that the population has a specified mean μ_0 , then we have $H_0 : \mu = \mu_0$
- Alternative hypothesis will be
 - **Two tailed alternative hypothesis** : $H_1 : \mu \neq \mu_0$.
 - **Right (One) tailed alternative hypothesis** : $H_1 : \mu > \mu_0$.
 - **Left (One) tailed alternative hypothesis** : $H_1 : \mu < \mu_0$.
- Hence alternative hypothesis helps to know whether the test is two tailed or one tailed test.
- Note that, Alternate hypothesis H_1 is either in the form of **less than** OR **greater than** OR **not equal**.

Errors In Sampling

- The main aim of the sampling theory is to draw a valid conclusion about the population parameters. In doing this we may commit the following two type of errors.
 - Type I error: When H_0 is true, we may reject it.
 $P(\text{Reject } H_0 \text{ when } H_0 \text{ is true}) = \alpha$
Where, α is called the size of the type I error also referred to as product's risk.
 - Type II error: When H_0 is wrong we may accept it.
 $P(\text{Accept } H_0 \text{ when } H_1 \text{ is true}) = \beta$
where, β is called the size of the type II error, also referred to as consumer's risk.


Unit - 5 Applied Statistics

Level of Significance

- The maximum probability of making a type I error is known as **level of significance**.
- It is denoted by α .
i.e., **$P(\text{Rejecting } H_0 \text{ when } H_0 \text{ is true}) = \alpha$** .
- The commonly used level of significance in practice are 5% (0.05) and 1% (0.01).
- For 5% level of significance means that there is a probability of making 5 out of 100 type I error. Similarly, 1% level of significance.
- If no level of significance is given, α is taken as 5% = 0.05.

Critical Region

- The region of the standard normal curve corresponding to predetermined level of significance α is known as **Critical Region**.
- It is also known as **Rejection Region**.
- The region under the normal curve which is not covered by the rejection region is known as **Acceptance Region**.
- Thus, the statistic which leads to rejection of null hypothesis H_0 gives rejection region or critical region.
- The value of the test statistic calculated to test the null hypothesis H_0 is known as **Critical Value**. Thus, the critical value separates the rejection region from the acceptance region.



Unit - 5 Applied Statistics

Critical value(Z_{α})	Level of significance (α)		
	1%	5%	10%
Two Tailed Test	$ Z_{\alpha} = 2.58$	$ Z_{\alpha} = 1.96$	$ Z_{\alpha} = 1.645$
Right Tailed Test	$Z_{\alpha} = 2.33$	$Z_{\alpha} = 1.645$	$Z_{\alpha} = 1.28$
Left Tailed Test	$Z_{\alpha} = -2.33$	$Z_{\alpha} = -1.645$	$Z_{\alpha} = -1.28$

- Note that, **Null Hypothesis H_0 is rejected when $|z| > 3$** without mentioning any level of significance.

Standard Error

- The standard deviation of the sampling distribution of a statistic is known as the **Standard Error**.
- It is denoted by **SE(t)** and read standard error of t.
- It plays an important role in the theory of large samples and it forms a basis of testing of hypothesis.

Test Statistics

- If t is any statistic, for large sample, then test statistics

$$Z = \frac{t - E(t)}{SE(t)} \sim N(0, 1)$$

is normally distributed with mean 0 and variance 1.

Where, $E(t)$ = Expected value of t

$SE(t)$ = Standard Error of t

Confidence Limits OR Confidence Interval

- The limits within which a hypothesis should lie with specified probability are known as **confidence limits**.
- Generally, the confidence limits are set up with 5% or 1% level of significance (α).
- If the sample value **lies between the confidence limit**, the hypothesis is **accepted**, otherwise **rejected**.

Unit - 5 Applied Statistics

Steps for Testing of Statistical Hypothesis

- **Step 1:** Set up null hypothesis H_0 .
- **Step 2:** Set up alternative hypothesis H_1 .
- **Step 3:** Set up level of significance α .
- **Step 4:** Apply test statistic.
- **Step 5:** Set up critical region. (Given in data OR to find from statistical tabular table).
- **Step 6:** Conclusion.
 - Compare the computed value of Z with critical value Z_α .
 - If $|Z| > |Z_\alpha|$, we **reject H_0** and conclude that there is significant difference.
 - If $|Z| < |Z_\alpha|$, we **accept H_0** and conclude that there is no significant difference.

Test for Variables for Large Samples

Method – 2 \rightsquigarrow Test of Significance of Mean

Test of Significance of Mean

- Consider a sample X size n with mean \bar{x} and SD s taken from population with mean μ and SD σ .
- This test is used to find significant difference between sample **mean \bar{x}** and **population mean μ** .
- In this test,

$$SE(t) = \begin{cases} \frac{\sigma}{\sqrt{n}} & ; \sigma \text{ is known} \\ \frac{s}{\sqrt{n}} & ; \sigma \text{ is unknown} \end{cases}$$

→ Formula for Test Statistics

- When population SD **σ is known**

$$Z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

where, $Q = 1 - P$

- When population SD **σ is not known**

$$Z = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

where, $q = 1 - p$

→ Confidence Limits

$$\text{Confidence Limits} = \bar{x} \pm Z_{\alpha} SE(t)$$

where, Z_{α} = Critical value at level of significance α

Example of Method-2: Test of Significance of Mean

C	1	<p>Let X be the length of a life of certain computer is approximately normally distributed with mean 800 days and standard deviation 40 days. If a random sample of 30 computers have an average life of 788 days, test the null hypothesis that $\mu \neq 800$ days at 5 % and 15% level of significance. $(Z_{0.05} = 1.96 ; Z_{0.15} = 1.45)$</p> <p>Answer: Null hypothesis accepted at 5% and rejected at 15%.</p>
C	2	<p>The mean IQ of a sample of 1600 children was 99. Is it likely that this was a random sample from a population with mean IQ 100 and SD 15? $(Z_{0.05} = 1.96)$</p> <p>Answer: Sample was not drawn from a population with mean 100 and SD 15.</p>
C	3	<p>An insurance agent has claimed that the average age of policy-holders who insure through him is less than the average for all agent, which is 30.5 years. This sample was drawn from a sample whose mean is 28.8 km and a standard deviation of 6.35 km. Test the significance at 0.05 level. $(Z_{0.05} = -1.645)$</p> <p>Answer: An insurance agent's claim is valid.</p>
C	4	<p>A college claims that its average class size is 35 students. A random sample of 64 students from class has a mean of 37 with a standard deviation of 6. Test at the $\alpha = 0.05$ level of significance if the claimed value is too low. $(Z_{0.05} = 1.645)$</p> <p>Answer: The true mean class size is likely to be more than 35.</p>

Method – 3 \rightarrow Test of Significance of Difference Between Two Means

Test of Significance of Difference Between Two Means

- \rightarrow Consider two samples X_1 and X_2 of sizes n_1 & n_2 , mean \bar{x}_1 & \bar{x}_2 and SD s_1 and s_2 respectively taken from two different population of sizes N_1 & N_2 , mean μ_1 & μ_2 and SD σ_1 and σ_2 .
- \rightarrow This test is used to find the significant difference between **sample means \bar{x}_1 & \bar{x}_2** .
- \rightarrow In this test,

$$SE(t) = \begin{cases} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} & ; \sigma \text{ is known} \\ \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} & ; \sigma \text{ is unknown} \end{cases}$$

\rightarrow Formula for Test Statistics

- When population SD **σ is known**

$$Z = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

where, $Q = 1 - P$

- When population SD **σ is not known**

$$Z = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

where, $q = 1 - p$

\rightarrow Confidence Limits

$$\text{Confidence Limits} = (\bar{x}_1 - \bar{x}_2) \pm Z_\alpha SE(t)$$

where, Z_α = Critical value at level of significance α

Example of Method-3: Test of Significance of Difference Between Two Means

C	1	<p>Test the significance of the difference between the means of two normal population with the same standard deviation from the following data:</p> <table><tr><td></td><td>Size</td><td>Mean</td><td>SD</td></tr><tr><td>Sample I</td><td>100</td><td>64</td><td>6</td></tr><tr><td>Sample II</td><td>200</td><td>67</td><td>8</td></tr></table> <p>($Z_{0.05} = 1.96$)</p> <p>Answer: There is significant difference between the means.</p>		Size	Mean	SD	Sample I	100	64	6	Sample II	200	67	8
	Size	Mean	SD											
Sample I	100	64	6											
Sample II	200	67	8											
C	2	<p>In a random sample of 100 light bulbs manufactured by a company A, the mean lifetime of light bulb is 1190 hours with standard deviation of 90 hours. Also, in a random sample of 75 light bulbs manufactured by company B, the mean lifetime of light bulb is 1230 hours with standard deviation of 120 hours. Is there a difference between the mean lifetime of the two brands of light bulbs at a significance level of 0.05 and 0.01?</p> <p>($Z_{0.05} = 1.96$; $z_{0.01} = 2.58$)</p> <p>Answer: There is difference between the mean lifetimes at 5%.</p> <p>There is no difference between the mean lifetimes at 1%.</p>												
C	3	<p>The means of simple samples of size 1000 and 2000 are 67.5 and 68 cm respectively. Can the sample be regarded as drawn from the same proportion of SD 2.5 cm. ($Z_{0.05} = 1.96$)</p> <p>Answer: Samples are not drawn from population having SD 2.5 cm.</p>												

Unit - 5 Applied Statistics

C	4	<p>A company A manufactured tube lights and claims that its tube lights are superior than its main competitor company B. The study showed that a sample of 40 tube lights manufactured by company A has a mean lifetime of 647 hours of continuous use with a standard deviation of 27 hours, while a sample of 40 tube lights manufactured by company B had a mean lifetime 638 hours of continuous use with a standard deviation of 31 hours. Does this substantiate the claim of company A that their tube lights are superior than manufactured by company B at 0.05 and 0.01 level of significance?</p> <p>($Z_{0.05} = 1.645$; $Z_{0.01} = 2.33$)</p> <p>Answer: The claim of company A is not valid at 5%.</p> <p>The claim of company A is not valid at 1%.</p>
C	5	<p>The average of marks scored by 32 boys is 72 with standard deviation 8 while that of 36 girls is 70 with standard deviation 6. Test at 1% level of significance whether the boys perform better than the girls. ($Z_{0.05} = 1.645$)</p> <p>Answer: The boys do not perform better than the girls.</p>

Method – 4 \rightsquigarrow Test of Significance of Difference Between Two Standard Deviations

Test of Significance of Difference Between Two Standard Deviations

- Consider two samples X_1 and X_2 of sizes n_1 & n_2 , and SD s_1 and s_2 respectively taken from two different population of sizes N_1 & N_2 , and SD σ_1 and σ_2 .
- This test is used to find the significant difference between **sample SDs s_1 & s_2** .
- In this test,

$$SE(t) = \begin{cases} \sqrt{\frac{\sigma_1^2}{2n_1} + \frac{\sigma_2^2}{2n_2}} & ; \sigma \text{ is known} \\ \sqrt{\frac{s_1^2}{2n_1} + \frac{s_2^2}{2n_2}} & ; \sigma \text{ is unknown} \end{cases}$$

→ Formula for Test Statistics

- When population SD **σ is known**

$$Z = \frac{s_1 - s_2}{\sqrt{\frac{\sigma_1^2}{2n_1} + \frac{\sigma_2^2}{2n_2}}}$$

- When population SD **σ is not known**

$$Z = \frac{s_1 - s_2}{\sqrt{\frac{s_1^2}{2n_1} + \frac{s_2^2}{2n_2}}}$$

→ Confidence Limits

$$\text{Confidence Limits} = (s_1 - s_2) \pm Z_\alpha SE(t)$$

where, Z_α = Critical value at level of significance α

Unit - 5 Applied Statistics

Example of Method-4: Test of Significance of Difference Between Two Standard Deviations

C	1	<p>The SD of a random sample of 900 members is 4.6 and that of another independent sample of 1600 members is 4.8. Examine if the two samples could have been drawn from a population with SD 4? ($Z_{0.05} = 1.96$)</p> <p>Answer: Two samples could have been drawn from a population with SD 4.</p>									
C	2	<p>Random samples drawn from two countries gave the following data relating to the heights of adult males:</p> <table border="1"> <thead> <tr> <th></th><th>Country A</th><th>Country B</th></tr> </thead> <tbody> <tr> <td>Standard Deviation</td><td>2.58</td><td>2.5</td></tr> <tr> <td>Number in samples</td><td>1000</td><td>1200</td></tr> </tbody> </table> <p>Is the difference between the standard deviation significant? ($Z_{0.05} = 1.96$)</p> <p>Answer: There is no significant difference between sample SDs.</p>		Country A	Country B	Standard Deviation	2.58	2.5	Number in samples	1000	1200
	Country A	Country B									
Standard Deviation	2.58	2.5									
Number in samples	1000	1200									

Unit - 5 Applied Statistics

Introduction

- If the sample are large ($n > 30$) then the sampling distribution of a statistics is normal. But if the samples are small ($n < 30$) then method that we have seen in previous unit does not hold good.
- For estimating of the parameter as well as for testing a hypothesis, following distributions are used:
 - Student's t – distribution
 - Snedecor's F – distribution
 - Chi – Square distribution

Degree of Freedom

- The number of independent pieces of information used to calculate a statistic is known as **degrees of freedom(df)**.
- In other words, they are the number of values that are able to be changed in a data set.
- For Example:

- Let two numbers: x, y. The mean of those numbers is m.
- In this data set of three variables (x, y, m), if you choose the values of any two variables, the third one is already determined.
- Let, x = 2, y = 4. You can't choose any mean you like as it's already determined as below:

$$m = \frac{x + y}{2} = 3$$

- Similarly, If you take, x = 3, m = 6, then y's value is automatically set. It is not free to change.

$$m = \frac{x + y}{2}$$

$$\Rightarrow 6 = \frac{3 + y}{2}$$

$$\Rightarrow y = 9$$

- Any time you assign some two values, the third has no **"freedom to change"**. Hence, there are **2** degrees of freedom in this example.

Unit - 5 Applied Statistics

Working Rule for Hypothesis Testing

- **Step 1:** Set up null hypothesis H_0 .
- **Step 2:** Set up alternative hypothesis H_1 .
- **Step 3:** Set up level of significance α .
- **Step 4:** Apply test statistic.
- **Step 5:** Find appropriate degree of freedom and set up critical region. (Given in data OR to find from statistical tabular table).
- **Step 6:** Conclusion.
 - Compare the computed value of t with critical value $t_{\alpha, v}$.
 - If $|t| > |t_{\alpha, v}|$, we **reject H_0** and conclude that there is significant difference.
 - If $|t| < |t_{\alpha, v}|$, we **accept H_0** and conclude that there is no significant difference.

Tests of Significance for Small Samples

Method – 5 \Rightarrow Test of Significance of Mean

Test of Significance of Mean

- Consider a sample X of size n with mean \bar{x} taken from normal population with mean μ of size N.
- This test is used to find the significant difference between **mean of sample \bar{x}** & **mean of population μ** when variance of the population is unknown.
- In this test,

$$SE(t) = f(x) = \begin{cases} \frac{s}{\sqrt{n-1}} & ; \quad \text{when SD is given in data} \\ \frac{S}{\sqrt{n}} & ; \quad \text{when SD is not given in data} \end{cases}$$

$$df(v) = n - 1$$

$$\text{Where, } s = \text{sample SD} = \sqrt{\frac{1}{n} \cdot \sum_{i=1}^n (x_i - \bar{x})^2}$$

$$S = \text{unbiased estimation of sample SD} = \sqrt{\frac{1}{n-1} \cdot \sum_{i=1}^n (x_i - \bar{x})^2}$$

- Relationship between sample SD “s” & estimates of SD “S” is **$n \cdot s^2 = (n - 1) \cdot S^2$**

$$\Rightarrow \frac{s^2}{n-1} = \frac{S^2}{n}$$

$$\Rightarrow \frac{s}{\sqrt{n-1}} = \frac{S}{\sqrt{n}}$$

- **Formula for Test Statistics**

$$t = \frac{\bar{x} - \mu}{SE(t)}$$

- **Confidence Limits**

$$\text{Confidence Limits} = \bar{x} \pm t_{\alpha, v} SE(t)$$

where, $t_{\alpha, v}$ = Critical value at level of significance α at degree of freedom (v).

Unit - 5 Applied Statistics

Example of Method-5: Test of Significance of Mean

C	1	<p>A random sample of size 16 has 53 as mean. The sum of squares of the derivation from mean is 135. Can this sample be regarded as taken from population having 56 as mean? ($t_{0.01,15} = 2.1314$)</p> <p>Answer: The sample mean has not come from a population mean 56.</p>
C	2	<p>The heights of 10 males of a given locality are found to be 175, 168, 155, 170, 152, 170, 175, 160, 160 and 165 cm. Based on this sample, find the 95% confidence limits for the heights of males in that locality.</p> <p>($t_{0.05,9} = 2.2622$)</p> <p>Answer: 159.2671 and 170.7329</p>
C	3	<p>Ten individuals were chosen random from a normal population and their heights were found to be in inches 63, 63, 66, 67, 68, 69, 70, 70, 71 and 71. Test the hypothesis that the mean height of the population is 66 inches.</p> <p>($t_{0.05,9} = 2.2622$)</p> <p>Answer: There is no significant difference between population mean and sample mean.</p>
C	4	<p>A soap manufacturing company was distributing a particular brand of soap through large number of retail shops. Before a heavy advertisement campaign, the mean sales per week per shop was 140 dozen. After the campaign, a sample of 26 shops was taken and the mean sales was found to be 147 dozen with standard deviation 16. Can you consider the advertisement effective? ($t_{0.05,25} = 1.7081$)</p> <p>Answer: The advertisement is effective.</p>

Method – 6 \rightsquigarrow Test of Significance of Difference Between Two Means

Test of Significance of Difference Between Two Means

- Consider two samples X_1 and X_2 of sizes n_1 & n_2 and of mean \bar{x} & \bar{y} respectively taken from two different population of sizes N_1 & N_2 and mean μ_X & μ_Y .
- This test is used to find the significant difference between **sample means \bar{x} & \bar{y}** under the assumption that the population variance are equal. ($\sigma_X = \sigma_Y = \sigma$).
- In this test,

$$SE(t) = S \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

$$df(v) = n_1 + n_2 - 2$$

Where,

$$S = \sqrt{\frac{1}{n_1 + n_2 - 2} \cdot \left[\sum_{i=1}^n (x_i - \bar{x})^2 + \sum_{i=1}^n (y_i - \bar{y})^2 \right]} ; \text{ when SD is not given in data}$$

$$= \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2}{n_1 + n_2 - 2}} ; \text{ when SD is given in data}$$

- **Formula for Test Statistics**

$$t = \frac{\bar{x} - \bar{y}}{SE(t)}$$

- **Confidence Limits**

$$\text{Confidence Limits} = (\bar{x} - \bar{y}) \pm t_{\alpha, v} SE(t)$$

where, $t_{\alpha, v}$ = Critical value at level of significance α at degree of freedom (v).

Example of Method-6: Test of Significance of Difference Between Two Means

C	1	<p>Two types of batteries are tested for their length of life and the following data are obtained:</p> <table><tr><td></td><td>No. of samples</td><td>Mean life in hours</td><td>Variance</td></tr><tr><td>Type A</td><td>9</td><td>600</td><td>121</td></tr><tr><td>Type B</td><td>8</td><td>640</td><td>144</td></tr></table> <p>Is there a significant difference in the two means? Find 95% confidence limits for the difference in means. ($t_{0.05,15} = 2.1314$)</p> <p>Answer: There is significant difference in the two means.</p> <p>Confidence Limits: – 27.3411 and – 52.65889</p>		No. of samples	Mean life in hours	Variance	Type A	9	600	121	Type B	8	640	144		
	No. of samples	Mean life in hours	Variance													
Type A	9	600	121													
Type B	8	640	144													
C	2	<p>The following data represents the biological values of protein from cow's milk and buffalo's milk at a certain level:</p> <table><tr><td>Cow's milk</td><td>1.82</td><td>2.02</td><td>1.88</td><td>1.61</td><td>1.81</td><td>1.54</td></tr><tr><td>Buffalo's milk</td><td>2.00</td><td>1.83</td><td>1.86</td><td>2.03</td><td>2.19</td><td>1.88</td></tr></table> <p>Examine if the average values of protein in the two samples significantly differ. ($t_{0.05,10} = 2.2281$)</p> <p>Answer: There is no significant difference in average values of proteins in two milk samples.</p>	Cow's milk	1.82	2.02	1.88	1.61	1.81	1.54	Buffalo's milk	2.00	1.83	1.86	2.03	2.19	1.88
Cow's milk	1.82	2.02	1.88	1.61	1.81	1.54										
Buffalo's milk	2.00	1.83	1.86	2.03	2.19	1.88										
C	3	<p>The mean height and SD height of 8 randomly chosen soldiers are 166.9 cm and 2.29 cm respectively. The corresponding values of 6 randomly chosen sailors are 107.3 cm and 8.50 cm respectively. Based on this data, can we conclude that soldiers are, in general, shorter than sailors?</p> <p>($t_{0.05,12} = -1.7823$)</p> <p>Answer: The soldiers are not shorter than sailors.</p>														

- C** **4** Samples of two types of electric bulbs were tested for length of life and the following data were obtained.

	Size	Mean	SD
Sample 1	8	1234	36 hr.
Sample 2	7	1036	40 hr.

Is the difference in the means sufficient to warrant that type 1 bulbs are superior to type 2 bulbs? ($t_{0.05,13} = 1.7709$)

Answer: The type 1 bulbs are superior to type 2 bulbs.

Method – 7 \rightsquigarrow F-test for Equality of Two Population Variances

F-test for Equality of Two Population Variances

- Consider two samples X_1 and X_2 of sizes n_1 & n_2 , mean \bar{x}_1 & \bar{x}_2 respectively taken from two different population of sizes N_1 & N_2 , mean μ and SD σ .
- This test is used to find the significance of the difference between **population standard deviations** σ_1 and σ_2 using unbiased estimates of SDs **S_1 and S_2** .

- In this test,

$$df(v) = n - 1$$

- **Formula for Test Statistics**

- If **$S_1 > S_2$** ,

$$F = \frac{(S_1)^2}{(S_2)^2}$$

- If **$S_2 > S_1$** ,

$$F = \frac{(S_2)^2}{(S_1)^2}$$

$$\text{where, } S_1 = \sqrt{\frac{1}{n_1 - 1} \cdot \sum_{i=1}^n (x_i - \bar{x})^2}$$

$$S_2 = \sqrt{\frac{1}{n_2 - 1} \cdot \sum_{i=1}^n (y_i - \bar{y})^2}$$

- Relationship between sample SD “s” & estimates of SD “S” is **$n \cdot s^2 = (n - 1) \cdot S^2$**

$$\Rightarrow S^2 = \frac{n \cdot s^2}{n - 1}$$

$$\text{So, } S_1^2 = \frac{n_1 \cdot s_1^2}{n_1 - 1} \quad ; \quad S_2^2 = \frac{n_2 \cdot s_2^2}{n_2 - 1}$$

Example of Method-7: F-test for Equality of Two Population Variances

C	1	<p>In a laboratory experiment two samples gave the following results:</p> <table><tr><th>Sample no.</th><th>Size</th><th>Mean</th><th>Variance</th></tr><tr><td>I</td><td>10</td><td>15</td><td>90</td></tr><tr><td>II</td><td>12</td><td>14</td><td>108</td></tr></table> <p>Test the equality of sample variances at 5 % level of significance. ($F_{0.05}(11, 9) = 3.10$)</p> <p>Answer: The two population have the same variances.</p>	Sample no.	Size	Mean	Variance	I	10	15	90	II	12	14	108				
Sample no.	Size	Mean	Variance															
I	10	15	90															
II	12	14	108															
C	2	<p>The time taken by workers in performing a job by method I and method II is given below.</p> <table><tr><td>Method I</td><td>20</td><td>16</td><td>26</td><td>27</td><td>22</td><td>-</td><td>-</td></tr><tr><td>Method II</td><td>27</td><td>33</td><td>42</td><td>35</td><td>32</td><td>34</td><td>38</td></tr></table> <p>Do the data show that the variances of time distribution in a population from which these samples are drawn do not differ significantly? ($F_{0.05}(6, 4) = 6.16$)</p> <p>Answer: The variances of time distribution in a population from which samples are drawn do not differ significantly.</p>	Method I	20	16	26	27	22	-	-	Method II	27	33	42	35	32	34	38
Method I	20	16	26	27	22	-	-											
Method II	27	33	42	35	32	34	38											
C	3	<p>Two random samples gave the following data:</p> <table><tr><th>Sample no.</th><th>Size</th><th>Mean</th><th>Variance</th></tr><tr><td>I</td><td>16</td><td>9.6</td><td>40</td></tr><tr><td>II</td><td>25</td><td>16.5</td><td>42</td></tr></table> <p>Can we conclude that the two samples have been drawn from the same normal population? ($F_{0.05}(24, 15) = 2.29$; $t_{0.05, 39} = 2.0227$)</p> <p>Answer: The two samples are not drawn from the same normal population.</p>	Sample no.	Size	Mean	Variance	I	16	9.6	40	II	25	16.5	42				
Sample no.	Size	Mean	Variance															
I	16	9.6	40															
II	25	16.5	42															

Unit - 5 Applied Statistics

C

4

Two nicotine contents in two random samples of tobacco are given below:

Sample I	21	24	25	26	27	-
Sample II	22	27	28	30	31	36

Can we say that two samples came from the same population?

($F_{0.05}(5, 4) = 6.26$; $|t_{0.05,9}| = 2.2622$)

Answer: Two samples came from the same population.

Unit - 5 Applied Statistics

Chi – Square Test: Introduction

- The chi-square (χ^2) test is a useful measure of comparing experimentally obtained results with those expected theoretically and based on hypothesis.
- Symbol χ^2 is read as “**Ky Square**”.
- It is used as a test statistic in testing a hypothesis that provides a set of theoretical frequencies with which observed frequencies are compared.
- The magnitude of discrepancy between observed and theoretical frequencies is given by the quantity χ^2 .
- If $\chi^2 = 0$, the observed and expected frequencies completely coincides. As the value of χ^2 increases, the discrepancy between the observed and theoretical frequency decreases.
- If $f_{o_1}, f_{o_2}, \dots, f_{o_n}$ be a set of observed frequencies and $f_{e_1}, f_{e_2}, \dots, f_{e_n}$ be the corresponding set of expected frequencies χ^2 then it is defined by,

$$\chi^2 = \frac{(f_{o_1} - f_{e_1})^2}{f_{e_1}} + \frac{(f_{o_2} - f_{e_2})^2}{f_{e_2}} + \dots + \frac{(f_{o_n} - f_{e_n})^2}{f_{e_n}} = \sum_{i=1}^n \frac{(f_{o_i} - f_{e_i})^2}{f_{e_i}}$$

- **Conditions for validity of χ^2 test:**
 - The sample observations should be independent.
 - The total frequency $N = \sum f_i$ should be reasonably large, say, greater than 50.
 - Each expected frequency $f_{e_i} \geq 5$. If not, then it is pooled with preceding or succeeding frequency so that the pooled frequency is more than 5. In this case, we need to adjust degree of freedom lost in pooling.
- **Applications of χ^2 test:**
 - Goodness of Fit
 - Independence of Attributes

Unit - 5 Applied Statistics

Working Rule for Chi- Square Test

- **Step 1:** Set up null hypothesis H_0 .
- **Step 2:** Set up alternative hypothesis H_1 .
- **Step 3:** Set up level of significance α .
- **Step 4:** Find appropriate f_e and apply test statistic.
- **Step 5:** Find appropriate degree of freedom and set up critical region. (Given in data OR to find from statistical tabular table).
- **Step 6:** Conclusion.
 - Compare the computed value of χ^2 with critical value $\chi_{\alpha, v}$.
 - If $\chi^2 > \chi_{\alpha, v}$, we **reject H_0** and conclude that there is significant difference.
 - If $\chi^2 < \chi_{\alpha, v}$, we **accept H_0** and conclude that there is no significant difference.

Method – 8 \rightsquigarrow Chi-square Test: for Goodness of Fit

Chi-square Test: for Goodness of Fit

→ The values of χ^2 is used to test whether the deviations of the observed frequencies from the expected frequencies are significant or not.

→ It is also used test to check fitting a set of observations to given distribution or not.

→ **Degree of freedom (df)**

- If the data is given in a series of n numbers, $v = n - 1$
- In case of binomial distribution, $v = n - 1$
- In case of poisson distribution, $v = n - 2$
- In case of normal distribution, $v = n - 3$

→ **Expected Frequency (f_e)**

- If the data is given in a series of n numbers,

$$f_e = \text{mean} = \frac{\sum x_i}{n}$$

Where, n = Total observation

- In case of binomial distribution,

$$f_e = N \cdot p(x)$$

Where, $p(x) = {}_n C_x \cdot p^x \cdot q^{n-x}$

x = Observed data (x_i)

n = Total observation

N = Sum of observed frequency

- In case of poisson distribution,

$$f_e = N \cdot p(x)$$

Where, $p(x) = \frac{e^{-\lambda} \cdot \lambda^x}{x!}$

x = Observed data (x_i)

N = Sum of observed frequency

$$\lambda = \frac{\sum x_i}{N}$$

Unit - 5 Applied Statistics

- In case of normal distribution,

$$f_e = N \cdot \text{Difference of area under the curve up to } Z_{x_i} \text{ and } Z_{x_i+1}$$

$$\text{Where, } Z_x = \frac{\bar{x} - \mu}{\sigma}$$

\bar{x} = Sample Mean (Given in Data OR Manually calculate from data)

μ = Population Mean (Given in Data)

σ = Population SD (Given in Data)

→ **Formula for Test of Significance**

$$\chi^2 = \sum_{i=1}^n \frac{(f_{o_i} - f_{e_i})^2}{f_{e_i}}$$

Example of Method-8: Chi-square Test: for Goodness of Fit

C	1	<p>Suppose that a die is tossed 120 times and the recorded data is as follows:</p> <table><tr><td>Face Observed(x)</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr><tr><td>Frequency</td><td>20</td><td>22</td><td>17</td><td>18</td><td>19</td><td>24</td></tr></table> <p>Test the hypothesis that the die is unbiased at $\alpha = 0.05$.</p> <p>$(\chi^2_{0.05,5} = 11.070)$</p> <p>Answer: The die is unbiased.</p>	Face Observed(x)	1	2	3	4	5	6	Frequency	20	22	17	18	19	24
Face Observed(x)	1	2	3	4	5	6										
Frequency	20	22	17	18	19	24										
C	2	<p>Theory predicts that the proportion of beans in the four group A, B, C, D should be 9 : 3 : 3 : 1. In an experiment among 1600 beans, the numbers in the four groups were 882, 313, 287 and 118. Do the experimental results support the theory? $(\chi^2_{0.05,3} = 7.815)$</p> <p>Answer: Experimental results support the theory.</p>														

Unit - 5 Applied Statistics

C	3	<p>Records taken of the number of male and female births in 830 families having four children are as follows:</p> <table><tr><td>No. of male births</td><td>0</td><td>1</td><td>2</td><td>3</td><td>4</td></tr><tr><td>No. of female births</td><td>4</td><td>3</td><td>2</td><td>1</td><td>0</td></tr><tr><td>No. of families</td><td>32</td><td>178</td><td>290</td><td>236</td><td>94</td></tr></table> <p>Test whether data are consistent with hypothesis that the binomial law holds and the chance of male birth is equal to that of female birth, namely.</p> <p>$(\chi^2_{0.05,4} = 9.488)$</p> <p>Answer: The data are not consistence with the hypothesis.</p>	No. of male births	0	1	2	3	4	No. of female births	4	3	2	1	0	No. of families	32	178	290	236	94		
No. of male births	0	1	2	3	4																	
No. of female births	4	3	2	1	0																	
No. of families	32	178	290	236	94																	
C	4	<p>Suppose that during 400 five-minute intervals the air-traffic control of an airport received 0, 1, 2, ... or 13 radio messages with respective frequencies of 3, 15, 47, 76, 68, 74, 46, 39, 15, 9, 5, 2, 0 and 1. Test at 0.05 level of significance, the hypothesis that the number of radio messages received during 5 minute interval follows Poisson distribution with $\lambda = 4.6$.</p> <p>$(\chi^2_{0.05,8} = 15.507)$</p> <p>Answer: Poisson distribution with $\lambda = 4.6$ provides a good fit.</p>																				
C	5	<p>The following table indicates (a) the frequencies of a given distribution with (b) the frequencies of the normal distribution having the same mean, standard deviation and the total frequency as in (a).</p> <table><tr><td>(a)</td><td>1</td><td>5</td><td>20</td><td>28</td><td>42</td><td>22</td><td>15</td><td>5</td><td>2</td></tr><tr><td>(b)</td><td>1</td><td>6</td><td>18</td><td>25</td><td>40</td><td>25</td><td>18</td><td>6</td><td>1</td></tr></table> <p>Apply the χ^2 - test of goodness of fit. $(\chi^2_{0.05,4} = 9.488)$</p> <p>Answer: This normal distribution provides a good fit.</p>	(a)	1	5	20	28	42	22	15	5	2	(b)	1	6	18	25	40	25	18	6	1
(a)	1	5	20	28	42	22	15	5	2													
(b)	1	6	18	25	40	25	18	6	1													
C	6	<p>Fit the equation of the best fitting normal curve to the following data:</p> <table><tr><td>X</td><td>135</td><td>145</td><td>155</td><td>165</td><td>175</td><td>185</td><td>195</td><td>205</td></tr><tr><td>f</td><td>1</td><td>6</td><td>18</td><td>25</td><td>40</td><td>25</td><td>18</td><td>6</td></tr></table> <p>Compare the theoretical and observed frequencies. Using χ^2 - test find goodness of fit. Given that $\mu = 165.6$ and $\sigma = 15.02$. $(\chi^2_{0.05,2} = 5.991)$</p> <p>Answer: This normal distribution provides a good fit.</p>	X	135	145	155	165	175	185	195	205	f	1	6	18	25	40	25	18	6		
X	135	145	155	165	175	185	195	205														
f	1	6	18	25	40	25	18	6														

***** End of the Unit *****