Peachtree Regional Medical Center
1450 Peachtree Industrial Blvd · Atlanta, GA 30309 · Main (404) 555-0100

TALWAR, ANJALI
DOB: 03/14/1958 (68 y) · F
MRN: 004417829 · CSN: 7731-0928-44
Adm: 10/05/2026 · Disch: 10/08/2026

AFTER VISIT SUMMARY — DISCHARGE INSTRUCTIONS

Attending: Samuel Brooks, MD (Hospital Medicine)
Unit: 5 West Telemetry, Rm 5214
Discharged to: Home, self-care
PCP on file: Linda Chen, MD
Pharmacy: CVS #4417 Ponce de Leon Ave
Code status: Full code

REASON FOR HOSPITALIZATION / DIAGNOSES

Type	Diagnosis	ICD-10
Principal	Atrial fibrillation with rapid ventricular response (RVR), new diagnosis	I48.91
Secondary	Hypokalemia	E87.6
Secondary	Hyperlipidemia, unspecified	E78.5

Hospital course: Patient presented to the ED 10/05/2026 with palpitations and dyspnea; ECG showed AF with RVR (HR 142). Rate controlled with IV diltiazem then transitioned to oral metoprolol succinate. TTE: LVEF 55–60%, mild LA enlargement. CHA_2DS_2-VASc = 3; anticoagulation initiated with apixaban after shared decision-making. K 3.1 on admission, repleted. Discharged in stable condition, HR 78, BP 128/76.

YOUR MEDICATION LIST — PLEASE READ CAREFULLY

Status	Medication	How to take	Notes
NEW	Metoprolol succinate ER 25 mg tablet	Take 1 tablet by mouth once daily	Controls heart rate
NEW	Apixaban (Eliquis) 5 mg tablet	Take 1 tablet by mouth twice daily (morning and evening)	Blood thinner — do not skip doses
NEW	Potassium chloride ER 20 mEq tablet	Take 1 tablet by mouth once daily for 7 days, with food	Then stop
CONTINUE	Atorvastatin 20 mg tablet	Take 1 tablet by mouth nightly	No change
STOP	Ibuprofen (Advil, Motrin) 200 mg	Do not take	Bleeding risk with apixaban. For pain use acetaminophen.

New prescriptions were sent electronically to CVS Pharmacy, Ponce de Leon Ave, (404) 555-0163. Please pick them up today.

FOLLOW-UP CARE

You should follow up with cardiology within 2 weeks of discharge for management of your new atrial fibrillation and anticoagulation; bring your medication list to this visit. Your primary care physician should see you within 7 days of discharge to review your hospital stay and your new medicines. Fasting blood work (basic metabolic panel and lipid panel) should be completed in 1 week to recheck potassium and kidney function while on the new medications — nothing to eat or drink except water for 10–12 hours before the draw. Results will be sent to Dr. Chen.

With	Where	Phone	When
Raymond Okafor, MD — Cardiology	Peachtree Heart Associates, 2100 Peachtree Rd NW, Ste 310	(404) 555-0142	Within 2 weeks. Patient to call to schedule.
Peachtree Labs — BMP + Lipid panel (fasting)	Peachtree Labs, 1450 Peachtree Industrial Blvd, Lower Level	(404) 555-0178	In 1 week. Fasting 10–12 h. Order on file.
Linda Chen, MD — Primary Care	Midtown Family Medicine, 990 W Peachtree St	(404) 555-0115	Within 7 days. Patient to call to schedule.

WHEN TO GET HELP RIGHT AWAY

CALL 911 or go to the nearest Emergency Department if you have:

- Chest pain or pressure
- Fainting or passing out
- Heart racing or pounding that will not slow down
- Sudden weakness or numbness on one side, face drooping, or trouble speaking — **these may be signs of a stroke. Call 911.**
- Unusual bleeding, black or tarry stools, or blood in your urine
- Severe shortness of breath

ACTIVITY

- Walk daily as tolerated; increase slowly.
- No heavy lifting over 10 lb for 1 week.
- You may shower. Do not drive if you feel dizzy.
- Use a soft toothbrush and electric razor while on apixaban.

DIET

- Heart-healthy, low-sodium diet (< 2,000 mg/day).
- Limit alcohol and caffeine; these can trigger AF.
- Eat potassium-rich foods (bananas, oranges, potatoes).

ABOUT ATRIAL FIBRILLATION

Atrial fibrillation (AFib) is an irregular heartbeat. It can cause blood clots that may lead to stroke. Your blood thinner (apixaban) lowers this risk. Your heart rate medicine (metoprolol) keeps your heart from beating too fast. Take both exactly as prescribed.

YOUR VITALS AT DISCHARGE

BP	HR	SpO_2	Wt
128/76	78 irreg	97% RA	71.2 kg

PENDING RESULTS AT DISCHARGE

None. Echocardiogram final report available in MyChart.

ADDITIONAL INSTRUCTIONS

Keep a list of all your medicines with you. Tell every doctor and dentist that you take a blood thinner. Avoid NSAIDs (ibuprofen, naproxen, aspirin unless prescribed). If you miss a dose of apixaban, take it as soon as you remember the same day; do not double up. Questions about your medicines: call your pharmacy or Dr. Chen's office.

Patient / Representative signature

Discharging RN: M. Alvarez, RN · 10/08/2026 14:22

Attending: Samuel Brooks, MD (e-signed)

I have received and understand these instructions. Interpreter services were offered. Copies: Patient, PCP (Linda Chen, MD), Cardiology (Raymond Okafor, MD).