

MILLS 8A

A REVIVAL OF MONOTYPE MODERN NO. 8A FROM THE *Bulletin of the American Mathematical Society*, 1940–1948

ROMAN, ITALIC, BOLD, SMALL CAPITALS

ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz fi fl ff ffi ffl

ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz fi fl ff ffi ffl

ABCDEFGHIJKLMNOPQRSTUVWXYZ abcdefghijklmnopqrstuvwxyz

SMALL CAPITALS: LEMMA, THEOREM, PROOF

FIGURES, LINING AND OLD STYLE

0123456789 0123456789 0123456789 0123456789

TEXT, 11 ON 12 PT

A function $f(x)$ is said to be a prime-representing function if $f(x)$ is a prime number for all positive integral values of x . It will be shown that there exists a real number A such that $[A^{3x}]$ is a prime-representing function, where $[R]$ denotes the greatest integer less than or equal to R .

Footnotes and references are set in the 9 pt cut: Ingham, *On the difference between consecutive primes*, Quart. J. Math. Oxford Ser. vol. 8 (1937) pp. 255–266.

MATHEMATICS

$$p_{n+1} - p_n < K p_n^{5/8}, \quad u_{n+1} = P_{n+1}^{3-n-1} > P_n^{3-n} = u_n, \quad \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p \frac{1}{1 - p^{-s}}$$

$$\int_0^{\infty} e^{-x^2} dx = \frac{\sqrt{\pi}}{2}, \quad \alpha\beta\gamma\eta\theta\lambda\mu\xi\pi\sigma\phi\chi\psi\omega\Gamma\Sigma\Phi$$

$$x \in A \cap B, \quad a \equiv b \pmod{p}, \quad a \not\equiv b, \quad \lim_{n \rightarrow \infty} x_n$$

ACCENTS AND PUNCTUATION

Erdős, Gödel, Poincaré, École, Schönflies; “quoted”, why? — § 3, †, &, 50%.