

Interspecies comparisons of Mg/Ca ratios in limpet shells

Niklas Hausmann^{a,*}, Donna Surge^b, Francisco Zangrando^c, Angelica Tivoli^c, Ivan Briz-Godino^d

^a*Leibniz Zentrum für Archäologie, Ludwig-Lindenschmit-Forum 1, Mainz, Germany, 55116*

^b*University of North Carolina, 104 South Road, 225 Geology Building, Chapel Hill, NC, US, 27599-3315*

^c*CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), Avenida Maipú 305, Ushuaia, Argentina, V9410BJA*

^d*, Spain,*

Abstract

This document is a template demonstrating the Aps format.

Keywords: Sclerochronology, Limpets, Elemental Ratio, Mg/Ca

1. Introduction

This study provides a short reassessment of the use of Magnesium to Calcium (Mg/Ca) ratios in Atlantic limpet shells to determine past sea surface temperatures. While recent studies of particularly *Patella* sp. in the Mediterranean and Southwest Europe have provided promising results ([Hausmann et al., 2019](#); [García-Escárzaga et al., 2015, 2018](#)).

In previous studies, various limpet specimens (*Patella vulgata*, *Nacella magellanica* and *Nacella odifosdio*) have been studied using Mg/Ca with mixed results. While *Patella vulgata* along the Spanish shoreline has since then repeatedly produced reliable correlations between sst and Mg/Ca ratios, this is not the case for other species.

In this study, we present elemental maps of various such species together with stable oxygen isotope values for some of the specimens. Some of these...

*Corresponding author

Email address: niklas@palaeo.eu (Niklas Hausmann)

2. Methods

3. Results




Figure 1: Correlation graphs for all specimens

4. Discussion





Figure 2: Correlation graphs for Ferguson et al. specimens

Table 1: Overview of comparative correlations {#tab:correlations}

Species	Locality	Specimen	Correlation R ²	Study
<i>Patella depressa</i>	Northern Spain	LAN541	0.87	10.3390/app11072959
		LAN545	0.86	
		LAN554	0.78	
		LAN559	0.82	
<i>Patella caerulea</i>	Croatia	ISTPC1	0.9	10.1038/s41598-019-39959-9
		ISTPC2	0.84	
	Crete	AF1911A	0.91 ¹	
		AF3003A	0.92 ²	
	Israel	AKKPC2	0.96	
		AKKPC3	0.89	
		FRMPC1	0.84	
		FRMPC2	0.96	
	Libya	MO31A	0.83	
		MP64A	0.33	
		MP67A	0.96	
		MP68A	0.81	
	Malta	MA10	0.82	
	Tunisia	TUNPC1	0.81	
		TUNPC2	0.78	
	Turkey	ANTPC1	0.95	
		ANTPC2	0.93	
		KIZPC1	0.94	
		KIZPC2	0.86	
<i>Patella rustica</i>	Gibraltar	JL1	0.02	doi.org/10.1016/j.epsl.2011.05.054
		JL2	0.8 (0.79)	
<i>Patella caerulea</i>	Gibraltar	JM00	0.69 (0.79)	
		JM30	0.83 (0.79)	
<i>Patella vulgata</i>	Orkney	ORK-LT5	not reported, here 0.88	doi.org/10.1016/j.palaeo.2016.10.021 and this study

¹SST only, no other geochemical data available²SST only, no geochemical data available

4.1. Reanalysis of ORK-LT5



References

García-Escárzaga, A., Clarke, L.J., Gutiérrez-Zugasti, I., González-Morales, M.R., Martínez, M., López-Higuera, J.M., Cobo, A., 2018. Mg/Ca profiles within archaeological mollusc (*patella vulgata*) shells: Laser-Induced breakdown spectroscopy compared to inductively coupled Plasma-Optical emission spectrometry. *Spectrochim. Acta Part B At. Spectrosc.* 148, 8–15. doi:[10.1016/j.sab.2018.05.026](https://doi.org/10.1016/j.sab.2018.05.026).

- García-Escárzaga, A., Moncayo, S., Gutiérrez-Zugasti, I., González-Morales, M.R., Martín-Chivelet, J., Cáceres, J.O., 2015. Mg/Ca ratios measured by laser induced breakdown spectroscopy (LIBS): a new approach to decipher environmental conditions. *J. Anal. At. Spectrom.* 30, 1913–1919. doi:[10.1039/C5JA00168D](https://doi.org/10.1039/C5JA00168D).
- Hausmann, N., Prendergast, A.L., Lemonis, A., Zech, J., Roberts, P., Siozos, P., Anglos, D., 2019. Extensive elemental mapping unlocks Mg/Ca ratios as climate proxy in seasonal records of mediterranean limpets. *Scientific Reports* 9, 3698. doi:[10.1038/s41598-019-39959-9](https://doi.org/10.1038/s41598-019-39959-9).