


AID-nummer: AID-number:	2198	Datum: Date:	18 - 06 - 05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN1

Blad nummer:
Sheet number:
281

Step 3: Form a tree



Step 4: Form conditional Database

A		-
B		A : 3
C		AB : 1 , AD : 1
D		A : 3

Step 5: for each x-conditional database
 form frequent ~~set~~ pattern by
 recursively calling FP-Grow

AID-nummer: AID-number:	2198	Datum: Date:	18-06-05
Kurskod: Course code:	F32A75	Provkod: Exam code:	TEN 1

Blad nummer: Sheet number:
25

A - Conditional DB

—

B - Conditional Database

Items : A

{2}

AB

—

Support : 3

/

Sort : A

A,,,

Output : AB

Cat AB - Cond

—

C - Conditional DB

Items : A B D

{2}

AC

—

Support : 2 1 1

/

BC

A:1

Sort : A B D

A,,

DC

A:1

Output : AC, BC, DC

B, D,

BC - Cond

Items : A

{2}

ABC

—

Sup : 2

/

Orders : A

A,

Output : ABC

AID-nummer: AID-number:	2198	Datum: Date:	18-06-05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN1

Blad nummer:
Sheet number:

26

DC - Cond

items ∈ A

support ∈ I

sort ∈ A

Output ∈ ADC

{ } ADC | —

|
A,

D - Conditional Database

items ∈ A

Support ∈ B

sort ∈ A

Output ∈ AD

{ } |

AD | —

| A III

↖

AID-nummer: AID-number:	2198	Datum: Date:	18 - 06 - 05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN1

Blad nummer:
Sheet number:
27
26

Q.8 (a)

range ≤ 5 is an example of antimonotone constraint since it implies

$c(A) = \text{true}$ then $c(B) = \text{true}$,
for $B \subseteq A$

range > 5 is an example of monotone constraint as in monotone constraint

$c(A) = \text{true}$ then $c(B) = \text{true}$
for $A \subseteq B$

$\text{avg}(S) \geq 6$ is an example of convertible monotone if data is arranged in order R which does not make it monotone but it can be made monotone if order will be changed i.e. rearrange data in decreasing order.

AID-nummer: AID-number:	2198	Datum: Date:	18-06-05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN 1

Blad nummer:
Sheet number:
28

$\text{arg}(s) > c$ is also an
example of ^{convertible} \uparrow antimonotone constraint
if data is arranged in some
sequence R which does not
satisfy the condition $C(B) = \text{true}$
then $C(A) = \text{true}$ for $B \subseteq A$

It can be made antimonotone
by changing seq 'R' that is
arrange in increasing order
of value.

AID-nummer: AID-number:	2198	Datum: Date:	18 - 06 - 05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN1

Blad nummer:
Sheet number:
29

Q. 8 (b)

$$50\% = \frac{50}{100} = 0.5$$

$$\text{minconf} \geq 0.5$$

ABC

$$AB \rightarrow C \Rightarrow \frac{\text{support}(ABC)}{\text{support}(AB)} = \frac{1}{3} < 0.5 \quad (\text{i.e } 30\% < 50\%)$$

Since $AB \rightarrow C$ does not satisfy the minconf and is not a valid rule thus

$$A \rightarrow BC$$

$$B \rightarrow AC$$

will not satisfy it either and thus not a frequent rule.

$$BC \rightarrow A \Rightarrow \frac{\text{support}(ABC)}{\text{support}(BC)} = \frac{1}{1} = 1$$

Since it satisfies the minconf
thus it is a valid association rule

AID-nummer: AID-number:	2198	Datum: Date:	18 - 06 - 05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN1

Blad nummer:
Sheet number:
30
201

and so will its subsets i.e

$$\begin{array}{l} B \longrightarrow CA \\ C \longrightarrow BA \end{array}$$

$$AC \longrightarrow B \Rightarrow \frac{\text{support}(ABC)}{\text{support}(AC)} = \frac{1}{2} \\ = 0.5 \geq 0.5 \\ 50\% = 50\%$$

since it satisfies the minconf thus
a valid association rule.

2

AID-nummer: AID-number:	2198	Datum: Date:	18 - 06 - 05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN1

Blad nummer:
Sheet number:
31
30

Q. 8 (c)

Consider a scenario where X is dependent on Y and Y is dependent on Z but X is independent of Z.

If there are no hidden variables, then it could be due to

$$X \rightarrow Y \rightarrow Z$$

$$X \leftarrow Y \leftarrow Z$$

$$X \rightarrow Y \leftarrow Z$$

~~or~~ selection bias, confounding, feedback
but never due to



$$X \leftarrow Y \rightarrow Z$$

1

If we say X is dependent on Y and Y & Z are dependent but X is not caused by Y then

$$X \rightarrow Y \rightarrow Z$$

is the only explanation

AID-nummer: AID-number:	2198	Datum: Date:	18-06-05
Kurskod: Course code:	732A75	Provkod: Exam code:	TEN 1

Blad nummer: Sheet number:	32
-------------------------------	----

Here, $X \rightarrow Y$ i.e.
 $X \rightarrow Y$ shows the causal
relation

similarly
relation $Y \rightarrow Z$ shows the causal