



ANCIENT METAGENOMICS


Physalia course, online, 2-4 June 2026

Data pre-processing: quality control and adapter removal

Nikolay Oskolkov, Metabolic Research Group leader at NIRI and Lund University



Physalia
Courses



nature reviews methods primers

Explore content ▾ About the journal ▾ Publish with us ▾

[nature](#) > [nature reviews methods primers](#) > [primers](#) > [article](#)

Primer | Published: 11 February 2021

Ancient DNA analysis


[Ludovic Orlando](#)  [Robin Allaby](#), [Pontus Skoglund](#), [Clio Der Sarkissian](#), [Philipp W. Stockhammer](#), [Maria C. Ávila-Arcos](#), [Qiaomei Fu](#), [Johannes Krause](#), [Eske Willerslev](#), [Anne C. Stone](#) & [Christina Warinner](#)

[Nature Reviews Methods Primers](#) 1, Article number: 14 (2021) | [Cite this article](#)

44k Accesses | 307 Citations | 519 Altmetric | [Metrics](#)

Abstract

Although the first ancient DNA molecules were extracted more than three decades ago, the first ancient nuclear genomes could only be characterized after high-throughput sequencing was invented. Genome-scale data have now been gathered from thousands of ancient archaeological specimens, and the number of ancient biological tissues amenable to genome sequencing is growing steadily. Ancient DNA fragments are typically ultrashort molecules and carry extensive amounts of chemical damage accumulated after death. Their extraction, manipulation and authentication require specific experimental wet-laboratory and dry-laboratory procedures before patterns of genetic variation from past individuals, populations and species can be interpreted. Ancient DNA data help to address an entire array of questions in anthropology, evolutionary biology and the environmental and archaeological sciences. The data have revealed a considerably more dynamic past than previously appreciated and have revolutionized our understanding of many major prehistoric and historic events. This Primer provides an overview of concepts and state-of-the-art methods underlying ancient DNA analysis and illustrates the diversity of resulting



ASM Journals | msystems | Vol. 6, No. 6 | Ancient Metagenomic Studies: Considerations for the Wider Scientific Community

ASM GLOBAL RESEARCH SYMPOSIUM
ASM-SIAT SYMPOSIUM
ON MICROBES IN BIOTECHNOLOGY
NOVEMBER 18-21, 2021 | SHENZHEN, CHINA
ONLINE EVENT - REGISTERABLE

Open Peer Review | Conduct of Scientific Research | Opinion/Hypothesis | 21 December 2021 | f x in

Ancient Metagenomic Studies: Considerations for the Wider Scientific Community

Authors: Clio Der Sarkissian, Irina M. Velko, Anna K. Fotakis, Ashild J. Vågne, Alexander Hübner, James A. Fellows Yates

AUTHORS INFO & AFFILIATIONS

<https://doi.org/10.1093/msystems/0310-21> | Check for updates

18 / 5,320

ABSTRACT

Like modern metagenomics, ancient metagenomics is a highly data-rich discipline, with the added challenge that the DNA of interest is degraded and, depending on the sample type, in low abundance. This requires the application of specialized measures during molecular experiments and computational analyses. Furthermore, researchers often work with finite sample sizes, which impedes optimal experimental design and control of confounding factors, and with ethically sensitive samples necessitating the consideration of additional guidelines. In September 2020, early career researchers in the field of ancient metagenomics met (Standards, Precautions & Advances in Ancient Metagenomics 2 [SPAAM2] community meeting) to discuss the state of the field and how to

ANNUAL REVIEWS

Annual Review of Genomics and Human Genetics
A Robust Framework for Microbial Archaeology

Christina Warinner,^{1,2} Alexander Herbig,¹ Allison Mann,^{1,2} James A. Fellows Yates,¹ Clemens L. Weiß,³ Hernán A. Burbano,³ Ludovic Orlando,^{4,5} and Johannes Krause¹

¹Department of Archaeogenetics, Max Planck Institute for the Science of Human History, Jena 07745, Germany; email: warinner@dh.mpg.de
²Department of Anthropology, University of Oklahoma, Norman, Oklahoma 73019
³Research Group for Ancient Genomics and Evolution, Department of Molecular Biology, Max Planck Institute for Developmental Biology, Tübingen 72076, Germany
⁴Centre for GeoGenetics, Natural History Museum of Denmark, 1350 Copenhagen N, Denmark
⁵Laboratoire d'Anthropologie Moléculaire et d'Imagerie de Synthèse, CNRS UMR 5288, Université Toulouse III – Paul Sabatier, Toulouse 31000, France

Further
Click here to view this article's online features:
• Download figures as PPT slides
• Navigate linked references
• Download citations
• Explore related articles
• Search keywords


Summary

- Basic Statistics
- Per base sequence quality
- Per tile sequence quality
- Per sequence quality scores
- Per base sequence content
- Per sequence GC content
- Per base N content
- Sequence Length Distribution
- Sequence Duplication Levels
- Overrepresented sequences
- Adapter Content

Basic Statistics

Measure	Value
Filename	G69146_pe_1.fastq.gz
File type	Conventional base calls
Encoding	Sanger / Illumina 1.9
Total Sequences	8997912
Sequences flagged as poor quality	0
Sequence length	77-101
%GC	49

Per base sequence quality



Quality Scores

A quality score (or Q-score) expresses an error probability. In particular, it serves as a convenient and compact way to communicate very small error probabilities.

Given an assertion, A, the quality score, Q(A), expresses the probability that A is not true, P(~A), according to the relationship:

$$Q(A) = -10 \log_{10}(P(\sim A))$$

where P(~A) is the estimated probability of an assertion A being wrong.


The relationship between the quality score and error probability is demonstrated with the following table:

Quality score, Q(A)	Error probability, P(~A)
10	0.1
20	0.01
30	0.001

Summary

- ✓ [Basic Statistics](#)
- ✓ [Per base sequence quality](#)
- ✓ [Per tile sequence quality](#)
- ✓ [Per sequence quality scores](#)
- ! [Per base sequence content](#)
- ✓ [Per sequence GC content](#)
- ✓ [Per base N content](#)
- ! [Sequence Length Distribution](#)
- ✓ [Sequence Duplication Levels](#)
- ✓ [Overrepresented sequences](#)
- ✗ [Adapter Content](#)


✓ Per sequence quality scores



Summary

- ✓ [Basic Statistics](#)
- ✓ [Per base sequence quality](#)
- ✓ [Per tile sequence quality](#)
- ✓ [Per sequence quality scores](#)
- ! [Per base sequence content](#)
- ✓ [Per sequence GC content](#)
- ✓ [Per base N content](#)
- ! [Sequence Length Distribution](#)
- ✓ [Sequence Duplication Levels](#)
- ✓ [Overrepresented sequences](#)
- ✗ [Adapter Content](#)


! Per base sequence content




Before fastp

After fastp

✖ Per base sequence content



✖ Per base sequence content




Was NovaSeq or NextSeq used for sequencing?


Summary

- ✓ Basic Statistics
- ✓ Per base sequence quality
- ✓ Per tile sequence quality
- ✓ Per sequence quality scores
- ! Per base sequence content
- ✓ Per sequence GC content
- ✓ Per base N content
- ! Sequence Length Distribution
- ✓ Sequence Duplication Levels
- ✓ Overrepresented sequences
- ✗ Adapter Content


✓ Per sequence GC content




FastQC: Per Sequence GC Content




🚫 Adapter Content



✅ Sequence Length Distribution



⚠️ Sequence Length Distribution




Adapter removal

Summary

- ✔ [Basic Statistics](#)
- ✔ [Per base sequence quality](#)
- ✔ [Per tile sequence quality](#)
- ✔ [Per sequence quality scores](#)
- ! [Per base sequence content](#)
- ✔ [Per sequence GC content](#)
- ✔ [Per base N content](#)
- ! [Sequence Length Distribution](#)
- ✔ [Sequence Duplication Levels](#)
- ✔ [Overrepresented sequences](#)
- ✘ [Adapter Content](#)


✔ Per base N content



Summary

- ✓ [Basic Statistics](#)
- ✓ [Per base sequence quality](#)
- ✓ [Per tile sequence quality](#)
- ✓ [Per sequence quality scores](#)
- ! [Per base sequence content](#)
- ✓ [Per sequence GC content](#)
- ✓ [Per base N content](#)
- ! [Sequence Length Distribution](#)
- ✓ [Sequence Duplication Levels](#)
- ✓ [Overrepresented sequences](#)
- ✗ [Adapter Content](#)


! Sequence Length Distribution



Summary

- ✓ [Basic Statistics](#)
- ✓ [Per base sequence quality](#)
- ✓ [Per tile sequence quality](#)
- ✓ [Per sequence quality scores](#)
- ! [Per base sequence content](#)
- ✓ [Per sequence GC content](#)
- ✓ [Per base N content](#)
- ! [Sequence Length Distribution](#)
- ✓ [Sequence Duplication Levels](#)
- ✓ [Overrepresented sequences](#)
- ✗ [Adapter Content](#)

✓ Sequence Duplication Levels




Summary

- ✓ [Basic Statistics](#)
- ✓ [Per base sequence quality](#)
- ✓ [Per tile sequence quality](#)
- ✓ [Per sequence quality scores](#)
- ! [Per base sequence content](#)
- ✓ [Per sequence GC content](#)
- ✓ [Per base N content](#)
- ! [Sequence Length Distribution](#)
- ✓ [Sequence Duplication Levels](#)
- ✓ [Overrepresented sequences](#)
- ✗ [Adapter Content](#)

✓ Overrepresented sequences

No overrepresented sequences

✗ Adapter Content



A modular tool to aggregate results from bioinformatics analyses across many samples into a single report.

Report generated on 2017-09-01, 09:26 based on data in: `/pica/v11/b2017206_nobackup/private/wabi/RNAseq/multiQC_raw_vs_trimmed`

📘 Welcome! Not sure where to start?

Watch a tutorial video

(6:06)

don't show again ✕

General Statistics

📄 Copy table

⚙️ Configure Columns

📊 Plot

Showing 18/18 rows and 4/5 columns.

Sample Name	% Dups	% GC	Length	M Seqs
P8205_101_S1_L001_R1_001	75.7%	46%	126 bp	81.2
P8205_101_S1_L001_R1_001.trimmomatic	75.7%	46%	126 bp	80.5
P8205_101_S1_L001_R1_001.trimmomatic_unpaired	38.7%	47%	106 bp	0.7
P8205_101_S1_L001_R2_001	71.9%	47%	126 bp	81.2
P8205_101_S1_L001_R2_001.trimmomatic	72.0%	46%	126 bp	80.5
P8205_101_S1_L001_R2_001.trimmomatic_unpaired	7.7%	47%	126 bp	0.0
P8205_102_S2_L001_R1_001	83.5%	46%	126 bp	86.2
P8205_102_S2_L001_R1_001.trimmomatic	83.5%	46%	126 bp	85.5
P8205_102_S2_L001_R1_001.trimmomatic_unpaired	50.3%	47%	108 bp	0.7
P8205_102_S2_L001_R2_001	80.0%	47%	126 bp	86.2
P8205_102_S2_L001_R2_001.trimmomatic	80.1%	47%	126 bp	85.5
P8205_102_S2_L001_R2_001.trimmomatic_unpaired	4.0%	41%	126 bp	0.0

FastQC

FastQC is a quality control tool for high throughput sequence data, written by Simon Andrews at the Babraham Institute in Cambridge.

Sequence Quality Histograms


10 2 6

The mean quality value across each base position in the read. See the [FastQC help](#).

Y-Limits: ☒ on

Mean Quality Scores

📄 Export Plot



FastQC is a quality control tool for high throughput sequence data, written by Simon Andrews at the Babraham Institute in Cambridge.

Sequence Quality Histograms 10 2 6

The mean quality value across each base position in the read. See the [FastQC help](#).

Y-Limits: 0 95Per Sequence Quality Scores 15 3

The number of reads with average quality scores. Shows if a subset of reads has poor quality. See the [FastQC help](#).

Y-Limits: 0 95

Per Sequence GC Content


10

8


The average GC content of reads. Normal random library typically have a roughly normal distribution of GC content. See the [FastQC help](#). Y-Limits: ☒ on

Percentages

Counts



Illumina DNA construct: indexes and adapters




Zhou et al. 2015. Atlas Oral Microbiol. <https://doi.org/10.1016/B978-0-12-802234-4.00002-1>

GGTGATACGGCGACCACCACaccgacGGCCCTACACGACGCTCTTCCGATCTXXXXXXAGCACACGTCTGGGCTCCAGTCACgacactaCCGTCTTCTGCTTG

TTACTATGCCGCTGGTGGTgtggtgttGGGATGTGCTGCGAGGGGGCTAGAXXXXXTCGTGTGCAGACTTGAGGTCAGTgctgtgatGGCAGGGGACGGGC



[Adapter/Index Primer] [Index] [Target primer] [Target] [Target primer] [Index] [Adapter/Index Primer]

@K00233:37:HGHLYBBXX:3:1101:2646:1121 1:N:0:NACGCATC+NGCTGGTG
NCGCATGAGCCGCCTGTATCAGGCGCTGATCGGGCCGGGCATTGCAGTTGGGATAGATCGGGGGAGCACACGTCTG
+
#A7F<<GG<JFJFJJJJJJJFFJJJJJJJAFFJFJJJJJJJFJAFFJJAJFJJ<FJJJJJFFF<FFA--FFFJJJJJ



Results

These plots are produced by the workflow, one for benchmarking results (Snakemake runs each tool three times to assess runtime, CPU and RAM usage), and one plot with adapter removal performance.



OPEN ACCESS

EDITED BY
 Richard Allen White III,
 University of North Carolina at Charlotte,
 United States

REVIEWED BY
 Emma Fay Harding,
 University of New South Wales, Australia
 Sydney Birch,
 University of North Carolina at Charlotte,
 United States

*CORRESPONDENCE
 Gabriel Renaud,
 gabriel.reno@gmail.com

RECEIVED 17 July 2023
 ACCEPTED 21 November 2023
 PUBLISHED 07 December 2023

CITATION
 Lien A, Legori LP, Kraft L, Sackett PW and
 Renaud G (2023), Benchmarking
 software tools for trimming adapters and
 merging next-generation sequencing
 data for ancient DNA.
Front. Bioinform. 3:1260486.
 doi: 10.3389/fbinf.2023.1260486

COPYRIGHT
 © 2023 Lien, Legori, Kraft, Sackett and
 Renaud. This is an open-access article
 distributed under the terms of the
 Creative Commons Attribution License
 (CC BY). The use, distribution or
 reproduction in other forums is
 permitted, provided the original author(s)
 and the copyright owner(s) are credited
 and that the original publication in this
 journal is cited, in accordance with
 accepted academic practice. No use,
 distribution or reproduction is permitted
 which does not comply with these terms.

Benchmarking software tools for trimming adapters and merging next-generation sequencing data for ancient DNA

Annette Lien¹, Leonardo Pestana Legori², Louis Kraft¹,
 Peter Wad Sackett¹ and Gabriel Renaud^{1*}

¹Department of Health Technology, Section for Bioinformatics, Technical University of Denmark, Kongens Lyngby, Denmark, ²University of Debrecen, Debrecen, Hungary

Ancient DNA is highly degraded, resulting in very short sequences. Reads generated with modern high-throughput sequencing machines are generally longer than ancient DNA molecules, therefore the reads often contain some portion of the sequencing adaptors. It is crucial to remove those adaptors, as they can interfere with downstream analysis. Furthermore, overlapping portions when DNA has been read forward and backward (paired-end) can be merged to correct sequencing errors and improve read quality. Several tools have been developed for adapter trimming and read merging, however, no one has attempted to evaluate their accuracy and evaluate their potential impact on downstream analyses. Through the simulation of sequencing data, seven commonly used tools were analyzed in their ability to reconstruct ancient DNA sequences through read merging. The analyzed tools exhibit notable differences in their abilities to correct sequence errors and identify the correct read overlap, but the most substantial difference is observed in their ability to calculate quality scores for merged bases. Selecting the most appropriate tool for a given project depends on several factors, although some tools such as fastp have some shortcomings, whereas others like leeHom outperform the other tools in most aspects. While the choice of tool did not result in a measurable difference when analyzing population genetics using principal component analysis, it is important to note that downstream analyses that are sensitive to wrongly merged reads or that rely on quality scores can be significantly impacted by the choice of tool.

Cutadapt

fastp


Trimmomatic

TrimGalore


AdapterRemoval

BBduk

Physalia
Courses



What if you forget to trim the adapters: soil sample



Perhaps there was a lake in the region where the soil sample came from?

No, you simply forgot to trim your adapters :)))

It turns out the Carp genome is full of Illumina adapters.

One of the first things we teach people in our [NGS courses](#) is how to remove adapters. It's not hard – we use [CutAdapt](#), but many other tools exist. It's simple, but really important – with De Bruijn graphs you will get paths through the graphs converging on kmers from adapters; and with OLC assemblers you will get spurious overlaps. With gap-fillers, it's possible to fill the gaps with sequences ending in adapters, and this may be what happened in the Carp genome.

Why then are we finding such elementary mistakes in such important papers? Why aren't reviewers picking up on this? It's frustrating.

This is a separate, but related issue, to genomic contamination – [the Wheat genome](#) has PhiX in it; [tons of bacterial genomes do too](#); and [lots of bacterial genes were problematically included in the Tardigrade genome](#) and declared as horizontal gene transfer.

Bioinformatics Bits and Bobs

Rare and random blog posts about bioinformatics, genomics and evolution.

Monday, 29 September 2014

Why you should QC your reads AND your assembly

The genome sequence of the Common Carp *Cyprinus carpio* was published in [Nature last week](#). By coincidence, I was doing some QC on some domesticated Ferret (*Mustela putorius furo*) reads, which had thrown some kmer warnings in the [FastQC tool](#). I blasted the kmers in NCBI and was quite perplexed by the number of hits that I found in the carp genome. Nearly all of the first 150 hits were all from the carp genome. Anyway, I looked a bit further into my odd kmers and it turns out that they were the ends of some Illumina adapter sequences that had presumably been incorporated into the paired-reads on the shorter ends of the insert size. This then took me back to the Carp Genome - what had crept into that?



*Knut och Alice
Wallenbergs
Stiftelse*

