

Reactive strategies with longer memory

ICSD 2024


Nikoleta E. Glynatsi, Ethan Akin, Martin A. Nowak, Christian Hilbe



MAX-PLANCK-GESELLSCHAFT




MAX-PLANCK-GESELLSCHAFT






MAX-PLANCK-GESELLSCHAFT




$$\begin{matrix} C & D \end{matrix}$$

$$\begin{matrix} C & (r & s \\ D & t & p) \end{matrix}$$


$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left(\begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$




$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left(\begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$





$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left(\begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$




$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left(\begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$



1	2	3	$n - 1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	\cdots	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$



1	2	3	$n - 1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	\cdots	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$




C

D

C

C

C



D


C

C

D

C

1	2	3	$n - 1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	\cdots	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$




C

D

C

C

C




D

C


C

D

C



1	2	3	$n - 1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	\cdots	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$




C
 D

D
 C



C
 C


C
 D



C
 C




1	2	3	$n-1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$	$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$	$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$	\cdots	$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$



	C	D	C	C
	D	C	C	D








	1	2	3	$n-1$	n
1	$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
2	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	\dots	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$
	C	D	C	C	C
	D	C	C	D	C









1	2	3	$n-1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$	$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$	$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$	\cdots	$\begin{matrix} C \\ D \end{matrix} \left(\begin{matrix} r & s \\ t & p \end{matrix} \right)$



	C	D	C	C
	D	C	C	D







Can we say anything about Nash equilibria in repeated games for any n ?





?


Nash
equilibria

Memory- n

Can we say anything about Nash equilibria in repeated games for any n ?

1	2	3	$n - 1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	\cdots	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$

	C	D	C	C	C
	D	C	C	D	C



Definitions

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Examples.

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Examples.

A reactive-1 strategy can be defined as: $\mathbf{p} = (p_C, p_D)$

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Examples.

A reactive-1 strategy can be defined as: $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as: $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Examples.

A reactive-1 strategy can be defined as: $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as: $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as: $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Examples.

A reactive-1 strategy can be defined as: $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as: $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as: $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

Tit for tat (1,0) Random $(\frac{1}{2}, \frac{1}{2})$

Definitions

Definition 1.

A reactive- n strategy can be defined as 2^n -dimensional vector $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$ with $0 \leq p_{\mathbf{h}^{-i}} \leq 1$ where \mathbf{h}^{-i} refers to an n -history of the co-player from the space of all possible co-player histories.

Examples.

A reactive-1 strategy can be defined as: $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as: $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$


A reactive-3 strategy can be defined as: $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

Tit for tat (1,0) Random $(\frac{1}{2}, \frac{1}{2})$


Definition 2.

A strategy \mathbf{p} for a repeated game is a Nash equilibrium if it is a best response to itself.

That is $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\sigma, \mathbf{p})$ for all other strategies σ .



p



p

Nash equilibria in higher n

Nash equilibria in higher n


Theorem. A reactive strategy $\mathbf{p} \in \mathcal{R}_n$ is a Nash equilibrium if and only if $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$ for all pure self-reactive strategies $\tilde{\mathbf{p}}$.

Nash equilibria in higher n

Theorem. A reactive strategy $\mathbf{p} \in \mathcal{R}_n$ is a Nash equilibrium if and only if $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$ for all pure self-reactive strategies $\tilde{\mathbf{p}}$.

1	2	3	$n - 1$	n
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	\cdots	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$

	C	D	C	C
	D	C	C	D





Nash equilibria in higher n

Theorem. A reactive strategy $\mathbf{p} \in \mathcal{R}_n$ is a Nash equilibrium if and only if $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$ for all pure self-reactive strategies $\tilde{\mathbf{p}}$.


Nash equilibria in higher n

Theorem. A reactive strategy $\mathbf{p} \in \mathcal{R}_n$ is a Nash equilibrium if and only if $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$ for all pure self-reactive strategies $\tilde{\mathbf{p}}$.




Nash equilibria in higher n

Theorem. A reactive strategy $\mathbf{p} \in \mathcal{R}_n$ is a Nash equilibrium if and only if $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$ for all pure self-reactive strategies $\tilde{\mathbf{p}}$.




$$\mathbf{p} = (p_C, p_D)$$




4

Nash equilibria in higher n

Theorem. A reactive strategy $\mathbf{p} \in \mathcal{R}_n$ is a Nash equilibrium if and only if $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$ for all pure self-reactive strategies $\tilde{\mathbf{p}}$.



$$\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$$



Donation game for $n = 2$

$$\begin{array}{cc} & \begin{matrix} C & D \end{matrix} \\ \begin{matrix} C \\ D \end{matrix} & \left(\begin{matrix} b - c & -c \\ b & 0 \end{matrix} \right) \end{array}$$

$$b > c > 0$$

Donation game for $n = 2$

Donation game for $n = 2$


Theorem. A reactive-2 strategy $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$ is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} = 1, \quad \frac{p_{CD} + p_{DC}}{2} < 1 - \frac{1}{2} \cdot \frac{c}{b}, \quad p_{DD} \leq 1 - \frac{c}{b}.$$

Donation game for $n = 2$

Theorem. A reactive-2 strategy $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$ is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,


$$p_{CC} = 1, \quad \frac{p_{CD} + p_{DC}}{2} < 1 - \frac{1}{2} \cdot \frac{c}{b}, \quad p_{DD} \leq 1 - \frac{c}{b}.$$



Donation game for $n = 2$

Theorem. A reactive-2 strategy $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$ is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} = 1, \quad \frac{p_{CD} + p_{DC}}{2} < 1 - \frac{1}{2} \cdot \frac{c}{b}, \quad p_{DD} \leq 1 - \frac{c}{b}.$$




Theorem. A reactive-2 strategy $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$ is a defective Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} \leq \frac{c}{b}, \quad \frac{p_{CD} + p_{DC}}{2} \leq \frac{c}{2b}, \quad p_{DD} = 0.$$

Donation game for $n = 2$


Theorem. A reactive-2 strategy $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$ is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} = 1, \quad \frac{p_{CD} + p_{DC}}{2} < 1 - \frac{1}{2} \cdot \frac{c}{b}, \quad p_{DD} \leq 1 - \frac{c}{b}.$$




Theorem. A reactive-2 strategy $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$ is a defective Nash equilibrium if and only if its entries satisfy the conditions,


$$p_{CC} \leq \frac{c}{b}, \quad \frac{p_{CD} + p_{DC}}{2} \leq \frac{c}{2b}, \quad p_{DD} = 0.$$




Evolutionary simulations




Evolutionary simulations




Evolutionary simulations



Evolutionary simulations








1. Algorithm to verify whether a given reactive- n strategy is an equilibrium.

```
input: p,n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in pure\_self\_reactive\_strategies$  do
    if p is not a best response  $\tilde{p}$  to then
        isNash ← False;
return (p, isNash);
```

1. Algorithm to verify whether a given reactive- n strategy is an equilibrium.

```
input: p, n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in \text{pure\_self\_reactive\_strategies}$  do
    if  $p$  is not a best response  $\tilde{p}$  to  $\tilde{p}$  then
        isNash ← False;
return (p, isNash);
```


2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$.




1. Algorithm to verify whether a given reactive- n strategy is an equilibrium.

```
input:  $p, n$ 
pure_self_reactive_strategies  $\leftarrow \{\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash  $\leftarrow$  True;
for  $\tilde{p} \in$  pure_self_reactive_strategies do
    if  $p$  is not a best response  $\tilde{p}$  to  $\tilde{p}$  then
        isNash  $\leftarrow$  False;
return ( $p$ , isNash);
```

2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$.




3. Longer memory helps sustain cooperation.




1. Algorithm to verify whether a given reactive- n strategy is an equilibrium.

```
input: p, n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in$  pure_self_reactive_strategies do
    if p is not a best response  $\tilde{p}$  to then
        isNash ← False;
return (p, isNash);
```

2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$.



3. Longer memory helps sustain cooperation.



More information

Conditional cooperation with longer memory

<https://arxiv.org/abs/2402.02437>


@NikoletaGlyn

<https://nikoleta-v3.github.io/>


1. Algorithm to verify whether a given reactive- n strategy is an equilibrium.

```
input: p, n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in$  pure_self_reactive_strategies do
    if p is not a best response  $\tilde{p}$  to then
        isNash ← False;
return (p, isNash);
```

2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$.



3. Longer memory helps sustain cooperation.



More information

Conditional cooperation with longer memory

<https://arxiv.org/abs/2402.02437>

@NikoletaGlyn

<https://nikoleta-v3.github.io/>

