

# **Reactive strategies with longer memory**


**MMEE 2024**

**Nikoleta E. Glynatsi, Ethan Akin, Martin A. Nowak, Christian Hilbe**




---

MAX-PLANCK-GESELLSCHAFT



# Social Behavior


## Understand Cooperation




$$\begin{matrix} C & D \end{matrix}$$

$$\begin{matrix} C & (r & s \\ D & t & p) \end{matrix}$$


$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left( \begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$




$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left( \begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$



$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left( \begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$





$$\begin{matrix} & C & D \\ \begin{matrix} C \\ D \end{matrix} & \left( \begin{matrix} r & s \\ t & p \end{matrix} \right) \end{matrix}$$




1	2	3	$n - 1$	$n$
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$\cdots$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$

---



1	2	3	$n - 1$	$n$
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\cdots$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$




$C$

$D$

$C$

$C$

$C$



$D$

$C$


$C$

$D$

$C$

1	2	3	$n - 1$	$n$
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$\cdots$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$

---




$C$

$D$

$C$

$C$

$C$




$D$

$C$


$C$

$D$

$C$



1	2	3	$n - 1$	$n$
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$\cdots$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$




$C$   
 $D$

$D$   
 $C$


$C$   
 $C$

$C$   
 $D$

$C$   
 $C$





	1	2	3	$n - 1$	$n$
1	$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
2	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\dots$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$
3	$C$	$D$	$C$	$C$	$C$
4	$D$	$C$	$C$	$D$	$C$





[1] Press, W.H. and Dyson, F.J., 2012. Iterated Prisoner's Dilemma contains strategies that dominate any evolutionary opponent.

[2] Stewart, A.J. and Plotkin, J.B., 2016. Small groups and long memories promote cooperation.




[1] Press, W.H. and Dyson, F.J., 2012. Iterated Prisoner's Dilemma contains strategies that dominate any evolutionary opponent.

[2] Stewart, A.J. and Plotkin, J.B., 2016. Small groups and long memories promote cooperation.




[1] Press, W.H. and Dyson, F.J., 2012. Iterated Prisoner's Dilemma contains strategies that dominate any evolutionary opponent.

[2] Stewart, A.J. and Plotkin, J.B., 2016. Small groups and long memories promote cooperation.





[1] Press, W.H. and Dyson, F.J., 2012. Iterated Prisoner's Dilemma contains strategies that dominate any evolutionary opponent.



[2] Stewart, A.J. and Plotkin, J.B., 2016. Small groups and long memories promote cooperation.





[1] Press, W.H. and Dyson, F.J., 2012. Iterated Prisoner's Dilemma contains strategies that dominate any evolutionary opponent.

[2] Stewart, A.J. and Plotkin, J.B., 2016. Small groups and long memories promote cooperation.







Can we say anything about Nash equilibria in repeated games for any  $n$ ?




Can we say anything about Nash equilibria in repeated games for any  $n$ ?

1	2	3	$n - 1$	$n$
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$	$\cdots$	$\begin{matrix} C & (r & s) \\ D & (t & p) \end{matrix}$

---

	$C$	$D$	$C$	$C$	$C$
	$D$	$C$	$C$	$D$	$C$



# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as:  $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as:  $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as:  $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as:  $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as:  $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

## Examples 2.

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as:  $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as:  $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

## Examples 2.

Tit for tat (1,0)

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as:  $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as:  $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

## Examples 2.

Tit for tat (1,0)

Random (1/2,1/2)

# Definitions

## Definition 1.

A reactive- $n$  strategy can be defined as  $2^n$ -dimensional vector  $\mathbf{p} = (p_{\mathbf{h}^{-i}})_{\mathbf{h}^{-i} \in H^{-i}}$  with  $0 \leq p_{\mathbf{h}^{-i}} \leq 1$  where  $\mathbf{h}^{-i}$  refers to an  $n$ -history of the co-player from the space of all possible co-player histories.

## Examples.

A reactive-1 strategy can be defined as:  $\mathbf{p} = (p_C, p_D)$

A reactive-2 strategy can be defined as:  $\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$

A reactive-3 strategy can be defined as:  $\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD})$

## Examples 2.

Tit for tat (1,0)

Random (1/2,1/2)

Two for Two Tats (1,1,1,0)

# **Definitions Continue**

# Definitions Continue

## Definition 3.

A strategy is considered pure if all conditional cooperation probabilities are either zero or one. If all cooperation probabilities are strictly between zero and one, the strategy is described as stochastic.

# Definitions Continue

## Definition 3.

A strategy is considered pure if all conditional cooperation probabilities are either zero or one. If all cooperation probabilities are strictly between zero and one, the strategy is described as stochastic.

## Examples 3.

Tit for tat  $(1,0)$   $\leftarrow$  pure

Random  $(1/2,1/2)$   $\leftarrow$  stochastic

Two for Two Tats  $(1,1,1,0)$   $\leftarrow$  pure

# Definitions Continue

## Definition 3.

A strategy is considered pure if all conditional cooperation probabilities are either zero or one. If all cooperation probabilities are strictly between zero and one, the strategy is described as stochastic.

## Examples 3.

Tit for tat  $(1,0) \leftarrow$  pure


Random  $(1/2,1/2) \leftarrow$  stochastic

Two for Two Tats  $(1,1,1,0) \leftarrow$  pure


## Definition 4.

A strategy  $\mathbf{p}$  for a repeated game is a Nash equilibrium if it is a best response to itself.

That is  $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\sigma, \mathbf{p})$  for all other strategies  $\sigma$ .



p



p

# Nash equilibria in higher $n$

# Nash equilibria in higher $n$

We established the following technical results:

# Nash equilibria in higher $n$

We established the following technical results:

# Nash equilibria in higher $n$



We established the following technical results:

1. Against reactive strategies, any feasible payoff can be generated with self-reactive strategies.

# Nash equilibria in higher $n$

We established the following technical results:



1. Against reactive strategies, any feasible payoff can be generated with self-reactive strategies.



# Nash equilibria in higher $n$

We established the following technical results:


1. Against reactive strategies, any feasible payoff can be generated with self-reactive strategies.




# Nash equilibria in higher $n$

We established the following technical results:

1. Against reactive strategies, any feasible payoff can be generated with self-reactive strategies.
2. To any reactive strategy, there is a best response among the pure self-reactive strategies.

1	2	3	$n - 1$	$n$
$C \ D$	$C \ D$	$C \ D$	$C \ D$	$C \ D$
$C \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$	$\dots$	$D \begin{pmatrix} r & s \\ t & p \end{pmatrix}$
<hr/>				
	$C$	$D$	$C$	$C$
	$D$	$C$	$C$	$D$





# Nash equilibria in higher $n$

**Theorem.** A reactive strategy  $\mathbf{p} \in \mathcal{R}_n$  is a Nash equilibrium if and only if  $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$  for all pure self-reactive strategies  $\tilde{\mathbf{p}}$ .


# Nash equilibria in higher $n$

**Theorem.** A reactive strategy  $\mathbf{p} \in \mathcal{R}_n$  is a Nash equilibrium if and only if  $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$  for all pure self-reactive strategies  $\tilde{\mathbf{p}}$ .




# Nash equilibria in higher $n$

**Theorem.** A reactive strategy  $\mathbf{p} \in \mathcal{R}_n$  is a Nash equilibrium if and only if  $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$  for all pure self-reactive strategies  $\tilde{\mathbf{p}}$ .




$$\mathbf{p} = (p_C, p_D)$$




4

# Nash equilibria in higher $n$

**Theorem.** A reactive strategy  $\mathbf{p} \in \mathcal{R}_n$  is a Nash equilibrium if and only if  $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$  for all pure self-reactive strategies  $\tilde{\mathbf{p}}$ .



$$\mathbf{p} = (p_{CC}, p_{CD}, p_{DC}, p_{DD})$$



# Nash equilibria in higher $n$

**Theorem.** A reactive strategy  $\mathbf{p} \in \mathcal{R}_n$  is a Nash equilibrium if and only if  $\pi(\mathbf{p}, \mathbf{p}) \geq \pi(\tilde{\mathbf{p}}, \mathbf{p})$  for all pure self-reactive strategies  $\tilde{\mathbf{p}}$ .



$$\mathbf{p} = (p_{CCC}, p_{CCD}, p_{CDC}, p_{CDD}, p_{DCC}, p_{DCD}, p_{DDC}, p_{DDD}) \quad 256$$

## Donation game for $n = 2$ & $n = 3$

$$\begin{array}{cc} C & D \\ \begin{matrix} C \\ D \end{matrix} & \left( \begin{matrix} b - c & -c \\ b & 0 \end{matrix} \right) \end{array}$$

$$b > c > 0$$

# Cooperative Nash

# Cooperative Nash

**Theorem.** A reactive-2 strategy  $\mathbf{p}$  is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} = 1, \quad \frac{p_{CD} + p_{DC}}{2} < 1 - \frac{1}{2} \cdot \frac{c}{b}, \quad p_{DD} \leq 1 - \frac{c}{b}.$$

# Cooperative Nash

**Theorem.** A reactive-2 strategy  $\mathbf{p}$  is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} = 1, \quad \frac{p_{CD} + p_{DC}}{2} < 1 - \frac{1}{2} \cdot \frac{c}{b}, \quad p_{DD} \leq 1 - \frac{c}{b}.$$

**Theorem.** A reactive-3 strategy  $\mathbf{p}$  is a cooperative Nash equilibrium if and only if its entries satisfy the conditions,

$$\begin{aligned} p_{CCC} &= 1 & \frac{p_{CDC} + p_{DCD}}{2} &\leq 1 - \frac{1}{2} \cdot \frac{c}{b} \\ \frac{p_{CCD} + p_{CDC} + p_{DCC}}{3} &\leq 1 - \frac{1}{3} \cdot \frac{c}{b} & \frac{p_{CDD} + p_{DCD} + p_{DDC}}{3} &\leq 1 - \frac{2}{3} \cdot \frac{c}{b} \\ \frac{p_{CCD} + p_{CDD} + p_{DCC} + p_{DDC}}{4} &\leq 1 - \frac{1}{2} \cdot \frac{c}{b} & p_{DDD} &\leq 1 - \frac{c}{b} \end{aligned}$$

# Defective Nash

# Defective Nash

**Theorem.** A reactive-2 strategy  $\mathbf{p}$  is a defective Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} \leq \frac{c}{b}, \quad \frac{p_{CD} + p_{DC}}{2} \leq \frac{c}{2b}, \quad p_{DD} = 0.$$

# Defective Nash



**Theorem.** A reactive-2 strategy  $\mathbf{p}$  is a defective Nash equilibrium if and only if its entries satisfy the conditions,

$$p_{CC} \leq \frac{c}{b}, \quad \frac{p_{CD} + p_{DC}}{2} \leq \frac{c}{2b}, \quad p_{DD} = 0.$$


**Theorem.** A reactive-3 strategy  $\mathbf{p}$  is a defecting Nash strategy if and only if its entries satisfy the conditions,

$$\begin{aligned} p_{CCC} &\leq \frac{c}{b}, & \frac{p_{CDC} + p_{DCD}}{2} &\leq \frac{1}{2} \cdot \frac{c}{b} \\ \frac{p_{CCD} + p_{CDC} + p_{DCC}}{3} &\leq \frac{2}{3} \cdot \frac{c}{b}, & \frac{p_{CDD} + p_{DCD} + p_{DDC}}{3} &\leq \frac{1}{3} \cdot \frac{c}{b} \\ \frac{p_{CCD} + p_{CDD} + p_{DCC} + p_{DDC}}{4} &\leq \frac{1}{2} \cdot \frac{c}{b}, & p_{DDD} &= 0. \end{aligned}$$


# Cooperative & Defective Nash




# Evolutionary simulations




# Evolutionary simulations




# Evolutionary simulations




# Evolutionary simulations



# Evolutionary simulations



# Simulations with error






# 1. Algorithm to verify whether a given reactive- $n$ strategy is an equilibrium.

```
input: p,n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in pure\_self\_reactive\_strategies$  do
    if p is not a best response  $\tilde{p}$  to then
        isNash ← False;
return (p, isNash);
```

## 1. Algorithm to verify whether a given reactive- $n$ strategy is an equilibrium.

```
input: p, n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in \text{pure\_self\_reactive\_strategies}$  do
    if  $p$  is not a best response  $\tilde{p}$  to  $\tilde{p}$  then
        isNash ← False;
return (p, isNash);
```


## 2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$ .




# 1. Algorithm to verify whether a given reactive- $n$ strategy is an equilibrium.

```
input:  $p, n$ 
pure_self_reactive_strategies  $\leftarrow \{\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash  $\leftarrow$  True;
for  $\tilde{p} \in$  pure_self_reactive_strategies do
    if  $p$  is not a best response  $\tilde{p}$  to  $\tilde{p}$  then
        isNash  $\leftarrow$  False;
return ( $p$ , isNash);
```

# 2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$ .




# 3. Longer memory helps sustain cooperation.




# 1. Algorithm to verify whether a given reactive- $n$ strategy is an equilibrium.

```
input:  $p, n$ 
pure_self_reactive_strategies  $\leftarrow \{\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash  $\leftarrow$  True;
for  $\tilde{p} \in$  pure_self_reactive_strategies do
| if  $p$  is not a best response  $\tilde{p}$  to  $\tilde{p}$  then
| | isNash  $\leftarrow$  False;
return ( $p$ , isNash);
```

# 2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$ .



# 3. Longer memory helps sustain cooperation.



# More information

Conditional cooperation with longer memory

<https://arxiv.org/abs/2402.02437>


@NikoletaGlyn

<https://nikoleta-v3.github.io/>


# 1. Algorithm to verify whether a given reactive- $n$ strategy is an equilibrium.

```
input: p, n
pure_self_reactive_strategies ← { $\tilde{p} \mid \tilde{p} \in \{0,1\}^{2^n}\}$ ;
isNash ← True;
for  $\tilde{p} \in$  pure_self_reactive_strategies do
    if p is not a best response  $\tilde{p}$  to then
        isNash ← False;
return (p, isNash);
```

# 2. Fully characterize cooperative & defective equilibria for $n = 2$ and $n = 3$ .



# 3. Longer memory helps sustain cooperation.



## More information

Conditional cooperation with longer memory

<https://arxiv.org/abs/2402.02437>

@NikoletaGlyn

<https://nikoleta-v3.github.io/>

