



The emptying of the Kakhovka reservoir: Sentinel-2 water, the day the bed fell dry and the bed afterwards (not observed is not dry)


(a) S2 water 06-05
2106 km² (99% observed)
the day before the breach




(b) S2 water 06-08
709 km² (37% observed)
IoU with the model 0.90



(c) S2 water 06-13
288 km² (17% observed)
IoU with the model 0.85




(d) S2 water 06-20
648 km² (100% observed)
the model ends 06-13




Sentinel-2 water
(a: p25 water3,
b-d: p15 crosscheck)
■ water
□ observed, not water
■ not observed (≠ dry)

(e) day the bed fell dry
317 km² by 06-13 (model)
+1166 km² by 06-20 (S2)



(f) S2 bed classes
07-05 (93% observed)



(g) S2 bed classes
09-08 (99% observed)




Sentinel-2 bed classes
(f, g; p25 k10e)

■ open water
■ shallow or mixed water
■ wet sediment
■ dry bare sediment
■ sparse herbaceous
■ dense herbaceous
■ reed or flooded vegetation
■ built hard surface
■ ambiguous

day the bed fell dry (e)

■ 06-06-07 (model)
■ 06-08-09 (model)
■ 06-10-11 (model)
■ 06-12-13 (model)
■ by 06-20 (Sentinel-2)
■ water on 06-20 (Sentinel-2)
■ not observed 06-20

(h) pool water area over time



— model (p95f sloped surface), 26 May - 13 June
○ model, upper estimate (Nikopol bound)
■ Sentinel-2, pool observed ($\geq 90\%$)
■ Sentinel-2, water share of the observed part
▼ Yi et al. 2025, Sentinel-1 (authors' archive)