



LUND
UNIVERSITY

Instrumental Variables and regression discontinuity designs


Nils Droste

2022 ClimBEco course



Instrumental Variables

Instrumental Variable (IV)



Using exogenous variation in instrument to close back-door. Image source: [Huntington-Klein 2018](#)



Instrumental Variables

An exemplary study

The Impact of the Women's March on the U.S. House Election*

Magdalena Larreboure

Felipe González

April 10, 2021

Three million people participated in the Women's March against discrimination in 2017, the largest single-day protest in U.S. history. We show that protesters in the March increased political preferences for women and people from ethnic minorities in the following federal election, the 2018 House of Representatives Election. Using **daily weather** shocks as exogenous drivers of attendance at the March, we show that protesters increased turnout at the Election and the vote shares obtained by minorities, particularly women, irrespective of their party affiliation. We conclude that protests can help to empower historically underrepresented groups through changes in local political preferences.

Larreboure2021

2022 ClimBEco course



Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression
Discontinuity

intuition

notation

examples

Instrumental Variables

But there may be plenty of causal pathways in reality

Introduction

Instrumental Variables

intuition

formalities



technicalities

Regression
Discontinuity

intuition

notation

examples



Weather IV and (in)dep vars, in general (left) and with temporal variations (right). Source: **Mellon2020**

Instrumental Variables

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation

examples



Two-Stage Least Square (2SLS) estimator

1. stage: regress Z on X :

$$X_i = \alpha_1 + \beta_1 Z_i + \gamma_1 W_i + \varepsilon_{1,i} \quad (1)$$

and predict the variation in X
explained by Z : $\hat{X} = \beta_1 Z_i$.

2. stage: plug in \hat{X} to estimate the
variation in Y not explained by
confounder W :

$$Y_i = \alpha_2 + \beta_2 \hat{X}_i + \gamma_2 W_i + \varepsilon_{2,i} \quad (2)$$

conditions

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation

examples

There are important conditions to consider

- *relevance* of instrument for predicting $Y \rightarrow E((\hat{X}_i|Z = 1) - (\hat{X}_i|Z = 0)) \neq 0$, aka Z is correlated with X , and thus with Y .



conditions

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation

examples



There are important conditions to consider

- *relevance* of instrument for predicting $Y \rightarrow E((\hat{X}_i|Z = 1) - (\hat{X}_i|Z = 0)) \neq 0$, aka Z is correlated with X , and thus with Y .
- *exclusion* restriction of Z being independent of Y : $E(\epsilon_i, Z_i|W_i) = 0$, aka no backdoor $Z \rightarrow Y$ or endogeneity, i.e. no relation with omitted variables.

conditions

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



There are important conditions to consider

- *relevance* of instrument for predicting $Y \rightarrow E((\hat{X}_i|Z=1) - (\hat{X}_i|Z=0)) \neq 0$, aka Z is correlated with X , and thus with Y .
- *exclusion* restriction of Z being independent of Y : $E(\epsilon_i, Z_i|W_i) = 0$, aka no backdoor $Z \rightarrow Y$ or endogeneity, i.e. no relation with omitted variables.

Now, let us see how to formulate this in the potential outcome notation.

conditions

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



There are important conditions to consider

- *relevance* of instrument for predicting $Y \rightarrow E((\hat{X}_i|Z = 1) - (\hat{X}_i|Z = 0)) \neq 0$, aka Z is correlated with X , and thus with Y .
- *exclusion* restriction of Z being independent of Y : $E(\epsilon_i, Z_i|W_i) = 0$, aka no backdoor $Z \rightarrow Y$ or endogeneity, i.e. no relation with omitted variables.

Now, let us see how to formulate this in the potential outcome notation.

For this let treatment or participation again be denoted by D , now as a function of the instrument $\rightarrow D_i(Z_i)$, the *intention to treat*.

notation

Step by step

- Imbens and Angrist (**Imbens1994**) formulate local average treatment effect (LATE)
 - for the *subpopulation* responding to instrument Z , that is those who participate $P(1)$ in treatment D

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



notation

Step by step

- Imbens and Angrist (**Imbens1994**) formulate local average treatment effect (LATE)
 - for the *subpopulation* responding to instrument Z , that is those who participate $P(1)$ in treatment D

$$E(Y_i|Z_i = 1) - E(Y_i|Z_i = 0) \quad (3)$$



notation

Step by step

- Imbens and Angrist (**Imbens1994**) formulate local average treatment effect (LATE)

- for the *subpopulation* responding to instrument Z , that is those who participate $P(1)$ in treatment D

$$E(Y_i|Z_i = 1) - E(Y_i|Z_i = 0) \quad (3)$$

- here the LATE is given by $P(1) \cdot E[Y_i(1) - Y_i(0)|D_i(1) = 1]$



notation

Step by step

- Imbens and Angrist (**Imbens1994**) formulate local average treatment effect (LATE)

- for the *subpopulation* responding to instrument Z , that is those who participate $P(1)$ in treatment D

$$E(Y_i|Z_i = 1) - E(Y_i|Z_i = 0) \quad (3)$$

- here the LATE is given by $P(1) \cdot E[Y_i(1) - Y_i(0)|D_i(1) = 1]$
- as long as participation $P(1) > P(0)$ and $D_i(1) \geq D_i(0) \forall i$, aka monotonic (or \leq , respectively)

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



notation

Step by step

- Imbens and Angrist (**Imbens1994**) formulate local average treatment effect (LATE)

- for the *subpopulation* responding to instrument Z , that is those who participate $P(1)$ in treatment D

$$E(Y_i|Z_i = 1) - E(Y_i|Z_i = 0) \quad (3)$$

- here the LATE is given by $P(1) \cdot E[Y_i(1) - Y_i(0)|D_i(1) = 1]$
- as long as participation $P(1) > P(0)$ and $D_i(1) \geq D_i(0) \forall i$, aka monotonic (or \leq , respectively)

Why?



Who's gonna be "treated"

Consider Angrist, Imbens and Rubin (**Angrist1996a**)

		$Z_i = 0$	
		$D_i(0) = 0$	$D_i(0) = 1$
$Z_i = 1$	$D_i(1) = 0$	Nevertaker	Defier
	$D_i(1) = 1$	Complier	Always-taker

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation

examples



Who's gonna be "treated"

Consider Angrist, Imbens and Rubin (**Angrist1996a**)

		$Z_i = 0$	
		$D_i(0) = 0$	$D_i(0) = 1$
$Z_i = 1$	$D_i(1) = 0$	Nevertaker	Defier
	$D_i(1) = 1$	Complier	Always-taker

- *"if people are more likely, on average, to participate given $Z = w$ than given $Z = z$, then anyone who would participate given $Z = z$ must also participate given $Z = w$ " (Imbens1994)*



Who's gonna be "treated"

Consider Angrist, Imbens and Rubin (**Angrist1996a**)

		$Z_i = 0$	
		$D_i(0) = 0$	$D_i(0) = 1$
$Z_i = 1$	$D_i(1) = 0$	Nevertaker	Defier
	$D_i(1) = 1$	Complier	Always-taker

- *"if people are more likely, on average, to participate given $Z = w$ than given $Z = z$, then anyone who would participate given $Z = z$ must also participate given $Z = w$ " (Imbens1994)*

→ assumes existence of only one of compliers or defiers, e.g. *no one* defies



Who's gonna be "treated"

Consider Angrist, Imbens and Rubin (**Angrist1996a**)

		$Z_i = 0$	
		$D_i(0) = 0$	$D_i(0) = 1$
$Z_i = 1$	$D_i(1) = 0$	Nevertaker	Defier
	$D_i(1) = 1$	Complier	Always-taker

- *"if people are more likely, on average, to participate given $Z = w$ than given $Z = z$, then anyone who would participate given $Z = z$ must also participate given $Z = w$ " (Imbens1994)*

→ assumes existence of only one of compliers or defiers, e.g. *no one* defies

- allows valid estimate of LATE, but may not always be realistic



Relaxing monotonicity assumption

de Chaisemartin (**DeChaisemartin2017**) shows IVs can be valid without strong monotonicity

- *"If there are defiers in the population, we only know that 2SLS estimates a weighted difference between the effect of the treatment among compliers and defiers"*

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Relaxing monotonicity assumption

de Chaisemartin (**DeChaisemartin2017**) shows IVs can be valid without strong monotonicity

- *"If there are defiers in the population, we only know that 2SLS estimates a weighted difference between the effect of the treatment among compliers and defiers"*
- a weak solution: $P(C_F) = P(F)$ and $E(Y(1) - Y(0)|C_F) = E(Y(1) - Y(0)|F)$

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Relaxing monotonicity assumption

de Chaisemartin (**DeChaisemartin2017**) shows IVs can be valid without strong monotonicity

- *"If there are defiers in the population, we only know that 2SLS estimates a weighted difference between the effect of the treatment among compliers and defiers"*
- a weak solution: $P(C_F) = P(F)$ and $E(Y(1) - Y(0)|C_F) = E(Y(1) - Y(0)|F)$
- *"is satisfied if a subgroup of compliers accounts for the same percentage of the population as defiers and has the same LATE"*

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Relaxing monotonicity assumption

de Chaisemartin (**DeChaisemartin2017**) shows IVs can be valid without strong monotonicity

- *"If there are defiers in the population, we only know that 2SLS estimates a weighted difference between the effect of the treatment among compliers and defiers"*
- a weak solution: $P(C_F) = P(F)$ and $E(Y(1) - Y(0)|C_F) = E(Y(1) - Y(0)|F)$
- *"is satisfied if a subgroup of compliers accounts for the same percentage of the population as defiers and has the same LATE"*
- I believe this can be approached with matching, too. See Murray et al. (**Murray2021**) who suggest to estimate the intention to treat $D(Z)$ with logistic regression, providing leeway for a propensity score or other matching approach (cf. **Hirano2003**; **Rosenbaum1984**).

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Weak instruments

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation

examples



When instruments are only weakly correlated with treatment, reconsider

$$Y_i = \alpha_2 + \beta_2 D_i + \gamma_2 W_i + \varepsilon_i \quad (4)$$

$$D_i = \alpha_1 + \beta_1 Z_i + \gamma_1 W_i + v_i \quad (5)$$

A condition was relevance, i.e. $E((D_i|Z = 1) - (D_i|Z = 0)) \neq 0$, or $\text{Cov}(Z_i, D_i|W_i) \neq 0$

- to estimate IV, $\hat{\beta}_2 = \frac{\text{Cov}(Y_i, Z_i)}{\text{Cov}(Z_i, D_i|W_i)}$
- problematic when $\text{Cov}(Z_i, D_i|W_i) \rightarrow 0$ as $\Delta\beta_2$ grows large even for small variations

Weak instruments

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



A range of techniques for robust parameter estimation in weak IV 2SLS

- F-test for strong enough instruments (**Stock2003**)

Weak instruments

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



A range of techniques for robust parameter estimation in weak IV 2SLS

- F-test for strong enough instruments (**Stock2003**)
- heteroskedasticity and autocorrelation robust for just identified models (**Chernozhukov2008**)

Weak instruments

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



A range of techniques for robust parameter estimation in weak IV 2SLS

- F-test for strong enough instruments (**Stock2003**)
- heteroskedasticity and autocorrelation robust for just identified models (**Chernozhukov2008**)
- heteroskedasticity, autocorrelation and cluster robust in a more general setting (**Montiel2013**)

Weak instruments

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



A range of techniques for robust parameter estimation in weak IV 2SLS

- F-test for strong enough instruments (**Stock2003**)
- heteroskedasticity and autocorrelation robust for just identified models (**Chernozhukov2008**)
- heteroskedasticity, autocorrelation and cluster robust in a more general setting (**Montiel2013**)
- a more powerful test with t-ratio critical value adjustments for significance testing (**Lee2020**)

Weak instruments

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



A range of techniques for robust parameter estimation in weak IV 2SLS

- F-test for strong enough instruments (**Stock2003**)
- heteroskedasticity and autocorrelation robust for just identified models (**Chernozhukov2008**)
- heteroskedasticity, autocorrelation and cluster robust in a more general setting (**Montiel2013**)
- a more powerful test with t-ratio critical value adjustments for significance testing (**Lee2020**)

→ We will look into some (basic) testing in the seminar.

intermediate summary

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Instrumental variables allow us to

- isolate a treatment effect by looking at the outcomes of exogeneously caused treatment variation
- it is considered a very robust causal inference, but assumptions are *somewhat* crucial
- mainly it is theory and reason that make a "valid instrument"
- there is loads of tests, I do not think they alone suffice

A first thought experiment

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Available online at www.sciencedirect.com



Ecological Economics 55 (2005) 527–538

**ECOLOGICAL
ECONOMICS**

www.elsevier.com/locate/ecolecon

ANALYSIS

Environmental pressure group strength and air pollution: An empirical analysis

Seth Binder, Eric Neumayer*

*Department of Geography and Environment and Center for Environmental Policy and Governance (CEPG),
London School of Economics and Political Science, Houghton Street, London WC2A 2AE, UK*

Received 7 December 2003; received in revised form 22 October 2004; accepted 14 December 2004

Available online 24 February 2005

A first thought experiment

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity


intuition

notation

examples



Do you think this is a valid instrument?



reformulating **Binder2005**: a (partial DAG)

Regression discontinuity designs (RDD) – intuition

Suppose we believe there is an effect for which assignment is non-random, but the cut-off at which treatment is assigned is quasi-random (cf.

Thistlewaite2016).

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Regression discontinuity designs (RDD) – intuition

Suppose we believe there is an effect for which assignment is non-random, but the cut-off at which treatment is assigned is quasi-random (cf. **Thistlewaite2016**).


Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



The border between Haiti and the Dominican Republic. Image source: **Wuepper2020**



concept


Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



The RDD concept and the effect of bin size choice. Image source: **Cattaneo2016**

Introduction

Instrumental Variables


intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples





Regression-Discontinuity-Design (RDD)



Focussing on effects just around the cutoff value. Image source: [Huntington-Klein 2018](#)

DAGs

Regression-Discontinuity-Design (RDD)



Focussing on effects just around the cutoff value. Image source: [Huntington-Klein 2018](#)

→ Do you see the IV in RDD?



notation

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples

Let us formulate in the potential outcomes notation. Suppose there is a outcome $(Y(1), Y(0))$ that depends on treatment D and covariate X .



notation

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples

Let us formulate in the potential outcomes notation. Suppose there is a outcome $(Y(1), Y(0))$ that depends on treatment D and covariate X .

While $Y(X)$ is assumed to be continuous, treatment D kicks in at a quasi-random threshold of \bar{x} , such that



notation

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples

Let us formulate in the potential outcomes notation. Suppose there is a outcome $(Y(1), Y(0))$ that depends on treatment D and covariate X .

While $Y(X)$ is assumed to be continuous, treatment D kicks in at a quasi-random threshold of \bar{x} , such that

$$D_i = \begin{cases} 1 & \text{if } x_i \geq \bar{x} \\ 0 & \text{if } x_i < \bar{x} \end{cases} \quad (6)$$



notation

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Let us formulate in the potential outcomes notation. Suppose there is a outcome $(Y(1), Y(0))$ that depends on treatment D and covariate X .

While $Y(X)$ is assumed to be continuous, treatment D kicks in at a quasi-random threshold of \bar{x} , such that

$$D_i = \begin{cases} 1 & \text{if } x_i \geq \bar{x} \\ 0 & \text{if } x_i < \bar{x} \end{cases} \quad (6)$$

The identifying assumption is again that treatment assignment is independent of outcomes $E(Y(1) - Y(0) \perp D | X = \bar{x})$.

Sharp RDD

The cutcoff at \bar{x} can be **sharp**

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Sharp RDD

The cutcoff at \bar{x} can be **sharp**

- in which case there is no overlap on both sides of \bar{x}

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation


examples



Sharp RDD

The cutoff at \bar{x} can be **sharp**


- in which case there is no overlap on both sides of \bar{x}
- we assume the outcomes would have been smooth in the absence of treatment (aka extrapolate a "bin" beyond the threshold)



Sharp RDD

The cutoff at \bar{x} can be **sharp**

- in which case there is no overlap on both sides of \bar{x}
- we assume the outcomes would have been smooth in the absence of treatment (aka extrapolate a "bin" beyond the threshold)



Imbens2008


- and measure $\tau_{srd} = \lim_{x \rightarrow \bar{x}} E[Y(1)|X = x] - \lim_{x \leftarrow \bar{x}} E[Y(0)|X = x]$



Sharp RDD

The cutoff at \bar{x} can be **sharp**

- in which case there is no overlap on both sides of \bar{x}
- we assume the outcomes would have been smooth in the absence of treatment (aka extrapolate a "bin" beyond the threshold)



Imbens2008

- and measure $\tau_{srd} = \lim_{x \rightarrow \bar{x}} E[Y(1)|X = x] - \lim_{x \leftarrow \bar{x}} E[Y(0)|X = x]$
- D is not just correlated but a deterministic function of x (once we know x and \bar{x} , we know D)

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



estimator

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



estimator

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



RDD estimation

$$Y_i = \alpha_i + \beta X_{it} + \gamma t_i + \varepsilon_{it} \quad (7)$$


where t indicates treatment cutoff values \bar{x} :

$$t_i = \begin{cases} 1 & \text{if } x_i \geq \bar{x} \\ 0 & \text{if } x_i < \bar{x} \end{cases} \quad (8)$$

This would often include polynomial terms to allow for non-linear functional forms (but should not, cf. **Gelman2019**). Another typical approach is a local linear regression (which is displayed in the animation) or smoothing functions.


fuzzy RDD

Suppose the data did **not** look like this




fuzzy RDD

but rather looked like this



fuzzy RDD

So we need to evaluate




Heiss 2020



fuzzy RDD

So we need to evaluate



Heiss 2020

This is literally an IV setting where a different probability on two sides of the cutoff predicts participation.



Fuzzy RDD

The cutoff at \bar{x} is be **fuzzy**

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Fuzzy RDD

The cutoff at \bar{x} is be **fuzzy**

- because of deniers or nevertakers etc, there is overlap on both sides of \bar{x}

Introduction

Instrumental Variables

intuition

formalities

technicalities

Regression Discontinuity

intuition

notation


examples



Fuzzy RDD

The cutoff at \bar{x} is be **fuzzy**

- because of deniers or nevertakers etc, there is overlap on both sides of \bar{x}
- probabilities differ: $\lim_{x \rightarrow \bar{x}} Pr(Y(1)|X = x) \neq \lim_{x \leftarrow \bar{x}} Pr(Y(0)|X = x)$




Imbens2008



Fuzzy RDD

The cutoff at \bar{x} is be **fuzzy**

- because of deniers or nevertakers etc, there is overlap on both sides of \bar{x}
- probabilities differ: $\lim_{x \rightarrow \bar{x}} Pr(Y(1)|X = x) \neq \lim_{x \leftarrow \bar{x}} Pr(Y(0)|X = x)$



Imbens2008


- if unconfounded, $\tau_{frdd} = E[Y(1)|D = 1, X = \bar{x}] - E[Y(0)|D = 1, X = \bar{x}]$



Fuzzy RDD

The cutoff at \bar{x} is be **fuzzy**

- because of deniers or nevertakers etc, there is overlap on both sides of \bar{x}
- probabilities differ: $\lim_{x \rightarrow \bar{x}} Pr(Y(1)|X = x) \neq \lim_{x \leftarrow \bar{x}} Pr(Y(0)|X = x)$



Imbens2008

- if unconfounded, $\tau_{frdd} = E[Y(1)|D = 1, X = \bar{x}] - E[Y(0)|D = 1, X = \bar{x}]$
- which we can estimate with 2SLS, predicting D in first stage, plugging estimates into second stage

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



examples

There are discontinuities in space



Wuepper2020a



examples

There are discontinuities in space


Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Regression discontinuities in covariates but not in vegetation potential, **Wuepper2020a**



examples

Time


Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



Hausman2018



examples

Rules

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples

The Causal Effect of Radical Right Success on Mainstream Parties' Policy Positions: A Regression Discontinuity Approach

TARIK ABOU-CHADI AND WERNER KRAUSE*

This article investigates how the success of radical right parties affects the policy positions of mainstream parties. We do this using a regression discontinuity approach that allows us to causally attribute mainstream parties' positional changes to radical right strength independent of public opinion as a potential confounder. Making use of exogenous variation created through differences in electoral thresholds, we empirically demonstrate that radical right success, indeed, causally affects mainstream parties' positions. This is true for mainstream left as well as mainstream right parties. These findings make an important contribution to the broader literature on party competition as they indicate that other parties' behavior and not only public opinion plays a crucial role in explaining parties' policy shift.

Keywords: radical right; party competition; immigration.

AbouChadi2018



examples

Rules


Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples




Hotelling-Downs Model of 2 Party Competition. Image Source: [Daniel Corradi Stevens](#)



examples

Rules



Mainstream party position on cultural position. Image source: **AbouChadi2018**



examples

There can be kinks, aka slope shifts


Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples




Ganong2018




examples

Multiple breaks



examples

Multiple breaks



Angrist1999



software

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples



available packages

- rdd
- rdrobust
- rdlocrand
- rddensity
- rdmulti
- rdpower

intermediate summary

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples

Regression discontinuity designs

- identify a causal effect at a (quasi-)randomly occurring break point that introduces treatment
- are the youngest "classical" causal inference methods and seen as favorable
- use breaks that can occur in space, time, institutions, etc. pp.



References I

Introduction

Instrumental Variables

intuition
formalities
technicalities

Regression Discontinuity

intuition
notation
examples

