

Designing and Evaluating a Likelihood-Ratio Test for IRT models

Methodology and Statistics for the Behavioural,

Biomedical and Social Sciences

Nina van Gerwen (1860852)

Supervisor: Dave Hessen

Candidate Journal: *Applied Psychological Measurement*

13th of October, 2022

1 Introduction

In organisations, a lot of effort tends to be dedicated to researching important organising variables, such as job performance and personality traits. These types of variables are latent constructs. This entails that they are not directly visible. Thus, to measure latent constructs we have to infer them based on responses to a test or questionnaire. To achieve this, Item Response Theory (IRT) is often used. A large issue in IRT, however, is that you must use a model that correctly describes the answering process of the test in order to gain correct conclusions. This means that you should investigate whether the model you use, fits the data you acquired. If a wrong model is used, it can lead to dire consequences such as faults in the validity of your measurement (Crişan et al., 2017; Jiao & Lau, 2003; Zhao & Hambleton, 2017) and by extension your conclusion. In an organisation, this could for example translate into issues such as discharging the wrong people. Hence, it is vital that you test the appropriateness of your model (i.e., model-fit).

Currently in IRT research, there are only two procedures that are applicable to every model that allow you to measure model-fit. Namely, the χ^2 -difference test and the Pearson's χ^2 -test. The χ^2 -difference test, however, requires your null model to be nested under an alternative model. Furthermore, the test is sensitive to large sample sizes, where it will always reject a model (Curran et al., 1996). This requirement of nesting also forms a serious issue for an often used model in IRT, namely the three-parameter logistic (3PL) model. This is because the 3PL model can only be nested under the four-parameter logistic (4PL) model, which suffers from consistency and estimation issues (Barton & Lord, 1981; Loken & Rulison, 2010). This in turn entails that the χ^2 -difference test often cannot be properly used under the 3PL model. Another option to assess model-fit is to calculate fit indices, which are standardised indicators of model appropriateness. For an overview of current fit indices in IRT and their limitations, see Nye et al. (2020). Besides these indices, fit indices from Structural Equation Modeling can also be used in IRT. However, hardly any research has been done towards the use of the Tucker-Lewis Index (TLI; Tucker and Lewis, 1973) and the Comparative Fit Index (CFI; Bentler, 1990) in IRT. We found only one paper examining the CFI (Yang, 2020) and two papers investigating the TLI (Cai et al., 2021; Yang, 2020). However, we believe that the calculations for the CFI and TLI by Yang might not be correct due to the baseline model used, which could have affected

the results. To summarise, there are (a) too few model-fit measures available in IRT to test the 3PL model and (b) insufficient research on the CFI and TLI, which we will attempt to address.

Our proposed research would (a) develop and test the performance of a model-fit Likelihood Ratio (LR) test that is applicable to all IRT models and (b) test the performance of the CFI and TLI with a complete-independence baseline model. More specifically, we will answer the following three research questions:

1. What is the robustness (i.e., empirical α and power) of the χ^2 -test associated with our LR?
2. How does the performance of our developed test compare to the performance of the χ^2 -difference and Pearson's χ^2 -test?
3. What is the performance of the TLI and CFI with a complete-independence baseline model in IRT?

2 Analytic strategy

In order to answer the research questions, we will conduct a simulation study. When simulating data, we will vary the following four variables:

- Test items
- Sample size
- Model types
- Number of group randomisation

For a complete overview of the conditions that will be used for the different variables, see *Table 1* below.

Table 1

Overview of Simulation Conditions for all Variables

Variable	Conditions	Description
Test items	5 - 10 - 20	The total number of items that the test will consist of
Sample size	20 - 50 - 100 - 200 - 500	The total number of observations that will be available for each item
Model type	1PL - 2PL - 3PL	The models that we will use as the basis for both data generation and model-fitting
Group randomisation	2 - 3 - 4	The number of groups that the total dataset gets divided into for the LR calculations

Note. 1PL = one-parameter logistic model; 2PL = two-parameter logistic model; 3PL = three-parameter logistic model.

Each particular condition will be replicated 500 times and in each replication the following LR will be calculated:

$$\frac{\prod_{i=1}^n f(x_i)}{\prod_{j=1}^g \prod_{i=1}^{n_g} f(x_{ij})} = \frac{L_0}{\prod_{j=1}^g L_j} \quad (1)$$

where $f(x)$ is the probability distribution, n is the total sample size, g is the group number and n_g is the sample size in group g . The groups will be gained by randomly assigning the dataset to the g groups. According to Wilk's theorem (Wilks, 1938), this LR will then asymptotically follow a χ^2 -distribution with the following formula:

$$-2\ln \frac{L_0}{\prod_{j=1}^g L_j} \rightarrow \chi^2(\Delta) \quad (2)$$

where Δ is the difference in the number of estimated parameters between the two likelihoods. This entails that the LR can be used for Null Hypothesis Significance testing. Then, due to the fact that we know the data-generating model, we can calculate the proportion of times that the test rejects the correct model (i.e., β : type II error). Furthermore, we can also calculate the proportion of times that the test accepts a wrong model (i.e., empirical α). Knowing these values, we can compare the multiple tests with one another,

where a test with higher values for either proportions will be noted as performing worse. This will inform us of the scenarios where our test works properly and scenarios where the test lacks power. To measure the performance of the TLI and CFI, we can calculate the proportion of times that the fit index improves (i.e., approaches the value 1) when the correct model is used compared to the other models. Finally, we will provide an empirical example by testing the LR test, CFI and TLI on an existing dataset and interpreting their results.

The proposed research would be conducted in R (R. C. Team, 2021) through the use of RStudio (R. Team, 2020). As for packages, we will use the MASS (Venables & Ripley, 2002), lavaan (Rosseel, 2012) and ltm (Rizopoulos, 2006) packages. Approval for the ethical consent for the simulated dataset has been requested. The approval for the empirical example, however, has to wait until a proper dataset has been found.

References

- Barton, M. A., & Lord, F. M. (1981). An upper asymptote for the three-parameter logistic item-response model. *ETS Research Report Series, 1981*(1), i–8.
- Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological bulletin, 107*(2), 238.
- Cai, L., Chung, S. W., & Lee, T. (2021). Incremental model fit assessment in the case of categorical data: Tucker–lewis index for item response theory modeling. *Prevention Science, 1*–12.
- Crısan, D. R., Tendeiro, J. N., & Meijer, R. R. (2017). Investigating the practical consequences of model misfit in unidimensional irt models [PMID: 28804181]. *Applied Psychological Measurement, 41*(6), 439–455. <https://doi.org/10.1177/0146621617695522>
- Curran, P., West, S., & Finch, J. (1996). The robustness of test statistics to nonnormality and specification error in confirmatory factor analysis. *Psychological Methods, 1*(1), 16–29. <https://doi.org/10.1037/1082-989X.1.1.16>
- Jiao, H., & Lau, A. C. (2003). The effects of model misfit in computerized classification test. *Annual meeting of the National Council of Educational Measurement, Chicago, IL.*
- Kang, T., & Cohen, A. S. (2007). Irt model selection methods for dichotomous items. *Applied Psychological Measurement, 31*(4), 331–358. <https://doi.org/10.1177/0146621606292213>
- Krammer, G. (2018). The andersen likelihood ratio test with a random split criterion lacks power. *Journal of modern applied statistical methods: JMASM, 17*, eP2685. <https://doi.org/10.22237/jmasm/1555594442>
- Loken, E., & Rulison, K. L. (2010). Estimation of a four-parameter item response theory model. *British Journal of Mathematical and Statistical Psychology, 63*(3), 509–525.
- Maydeu-Olivares, A. (2013). Goodness-of-fit assessment of item response theory models. *Measurement Interdisciplinary Research and Perspectives, 11*, 71–101. <https://doi.org/10.1080/15366367.2013.831680>
- Maydeu-Olivares, A., & Joe, H. (2014). Assessing approximate fit in categorical data analysis. *Multivariate Behavioral Research, 49*. <https://doi.org/10.1080/00273171.2014.911075>

- Nye, C. D., Joo, S.-H., Zhang, B., & Stark, S. (2020). Advancing and evaluating irt model data fit indices in organizational research. *Organizational Research Methods*, 23(3), 457–486. <https://doi.org/10.1177/1094428119833158>
- Orlando, M., & Thissen, D. (2000). Likelihood-based item-fit indices for dichotomous item response theory models. *Applied Psychological Measurement*, 24(1), 50–64. <https://doi.org/10.1177/01466216000241003>
- Rizopoulos, D. (2006). Ltm: An r package for latent variable modelling and item response theory analyses. *Journal of Statistical Software*, 17(5), 1–25. <https://doi.org/10.18637/jss.v017.i05>
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36. <https://doi.org/10.18637/jss.v048.i02>
- Team, R. C. (2021). *R: A language and environmental for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria. <https://www.R-project.org/>
- Team, R. (2020). *Rstudio: Integrated development environment for r*. RStudio, PBC. Boston, MA. <http://www.rstudio.com/>
- Tucker, L. R., & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, 38(1), 1–10.
- Venables, W. N., & Ripley, B. D. (2002). *Modern applied statistics with s* (Fourth) [ISBN 0-387-95457-0]. Springer. <https://www.stats.ox.ac.uk/pub/MASS4/>
- Wilks, S. S. (1938). The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses. *The Annals of Mathematical Statistics*, 9(1), 60–62. <https://doi.org/10.1214/aoms/1177732360>
- Yang, X. (2020). *Comparing global model-data fit indices in item response theory applications* [PhD dissertation]. Florida State University, College of Education.
- Zhao, Y., & Hambleton, R. (2017). Practical consequences of item response theory model misfit in the context of test equating with mixed-format test data. *Frontiers in Psychology*, 8. <https://doi.org/10.3389/fpsyg.2017.00484>