

imci-selected-v0 machine-readable rule set - Domain Expert Crosscheck

Source: WHO Integrated Management of Childhood Illness, Chart Booklet, March 2014

Population: Children aged 2 to 59 months

Review each EdgeIMCI-encoded rule derived from the WHO IMCI chart: does its condition, classification, and action set preserve the selected source logic? These are not WHO-authored machine-readable rules and do not represent complete IMCI. Tick the box in the last column if correct, or write a note if something is wrong.

Rule ID	Clinical Area	If (condition)	Then	Actions / Note	WHO page	Correct?
IMCI-GDS-UNABLE-TO-DRINK	General danger signs	Child has: unable to drink or breastfeed	Very severe disease	Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral	p.5 (chart 1 of 76)	[]
IMCI-GDS-VOMITS-EVERYTHING	General danger signs	Child has: vomits everything	Very severe disease	Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral	p.5 (chart 1 of 76)	[]
IMCI-GDS-CONVULSIONS-HISTORY	General danger signs	Child has: had convulsions	Very severe disease	Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral	p.5 (chart 1 of 76)	[]
IMCI-GDS-LETHARGIC-OR-UNCONSCIOUS	General danger signs	Child has: lethargic or unconscious	Very severe disease	Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral	p.5 (chart 1 of 76)	[]
IMCI-GDS-CONVULSING-NOW	General danger signs	Child has: convulsing now	Very severe disease	Give diazepam (if convulsing now); Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral	p.5 (chart 1 of 76)	[]
IMCI-RESP-FAST-BREATHING-2-12M	Cough or difficult breathing	Child aged 2 to 12 months with respiratory rate ≥ 50 breaths/min	Sets: fast breathing = yes (used by rule IMCI-RESP-PNEUMONIA-FAST-BREATHING)	Not a classification rule - threshold check that feeds the respiratory classification rules below	p.6 (chart 2 of 76)	[]
IMCI-RESP-FAST-BREATHING-12-60M	Cough or difficult breathing	Child aged 12 to 60 months with respiratory rate ≥ 40 breaths/min	Sets: fast breathing = yes (used by rule IMCI-RESP-PNEUMONIA-FAST-BREATHING)	Not a classification rule - threshold check that feeds the respiratory classification rules below	p.6 (chart 2 of 76)	[]
IMCI-RESP-SEVERE-DANGER-SIGN	Cough or difficult breathing	Any general danger sign is present	Severe pneumonia or very severe disease	Give first dose of appropriate antibiotic; Urgent referral	p.6 (chart 2 of 76)	[]

Rule ID	Clinical Area	If (condition)	Then	Actions / Note	WHO page	Correct?
IMCI-RESP-SEVERE-STRIDOR	Cough or difficult breathing	Child has: stridor when calm	Severe pneumonia or very severe disease	Give first dose of appropriate antibiotic; Urgent referral	p.6 (chart 2 of 76)	[]
IMCI-RESP-PNEUMONIA-CHEST-INDRAWING	Cough or difficult breathing	Child has: chest indrawing	Pneumonia	Give oral amoxicillin for 5 days; Soothe throat and relieve cough; Advise when to return immediately; Follow up in 3 days	p.6 (chart 2 of 76)	[]
IMCI-RESP-PNEUMONIA-FAST-BREATHING	Cough or difficult breathing	Fast breathing detected (threshold met)	Pneumonia	Give oral amoxicillin for 5 days; Soothe throat and relieve cough; Advise when to return immediately; Follow up in 3 days	p.6 (chart 2 of 76)	[]
IMCI-RESP-COUGH-OR-COLD	Cough or difficult breathing	None of the above conditions met (fallback)	Cough or cold	Soothe throat and relieve cough; Advise when to return immediately; Follow up in 5 days if not improving	p.6 (chart 2 of 76)	[]
IMCI-DIARRHOEA-SEVERE-DEHYDRATION	Diarrhoea dehydration	At least 2 of: lethargic or unconscious = yes; sunken eyes = yes; drinking status in (Unable to drink, Drinks poorly); skin pinch = Goes back very slowly	Severe dehydration	If no other severe classification: Give fluid for severe dehydration (Plan C) If other severe classification present: Urgent referral; Frequent ORS sips during referral; Continue breastfeeding	p.7 (chart 3 of 76)	[]
IMCI-DIARRHOEA-SOME-DEHYDRATION	Diarrhoea dehydration	At least 2 of: restless or irritable = yes; sunken eyes = yes; drinking status = Drinks eagerly / thirsty; skin pinch = Goes back slowly	Some dehydration	If no other severe classification: Give fluid, zinc and food (Plan B); Advise when to return immediately; Follow up in 5 days if not improving If other severe classification present: Urgent referral; Frequent ORS sips during referral; Continue breastfeeding	p.7 (chart 3 of 76)	[]
IMCI-DIARRHOEA-NO-DEHYDRATION	Diarrhoea dehydration	None of the above conditions met (fallback)	No dehydration	Give fluid, zinc and food (Plan A); Advise when to return immediately; Follow up in 5 days if not improving	p.7 (chart 3 of 76)	[]