

# IMCI Rules - Domain Expert Crosscheck

Source: WHO Integrated Management of Childhood Illness, Chart Booklet, March 2014

Population: Children aged 2 to 59 months

Review each rule: does the condition, classification, and actions match the WHO IMCI chart? Tick the box in the last column if correct, or write a note if something is wrong.

| Rule ID                                  | Clinical Area                | If (condition)                                                         | Then                                                                         | Actions / Note                                                                                                                                               | WHO page            | Correct? |
|------------------------------------------|------------------------------|------------------------------------------------------------------------|------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|----------|
| <b>IMCI-GDS-UNABLE-TO-DRINK</b>          | General danger signs         | Child has: unable to drink or breastfeed                               | Very severe disease                                                          | Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral                                    | p.5 (chart 1 of 76) | [ ]      |
| <b>IMCI-GDS-VOMITS-EVERYTHING</b>        | General danger signs         | Child has: vomits everything                                           | Very severe disease                                                          | Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral                                    | p.5 (chart 1 of 76) | [ ]      |
| <b>IMCI-GDS-CONVULSIONS-HISTORY</b>      | General danger signs         | Child has: had convulsions                                             | Very severe disease                                                          | Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral                                    | p.5 (chart 1 of 76) | [ ]      |
| <b>IMCI-GDS-LETHARGIC-OR-UNCONSCIOUS</b> | General danger signs         | Child has: lethargic or unconscious                                    | Very severe disease                                                          | Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral                                    | p.5 (chart 1 of 76) | [ ]      |
| <b>IMCI-GDS-CONVULSING-NOW</b>           | General danger signs         | Child has: convulsing now                                              | Very severe disease                                                          | Give diazepam (if convulsing now); Complete assessment quickly; Give pre-referral treatment immediately; Prevent low blood sugar; Keep warm; Urgent referral | p.5 (chart 1 of 76) | [ ]      |
| <b>IMCI-RESP-FAST-BREATHING-2-12M</b>    | Cough or difficult breathing | Child aged 2 to 12 months with respiratory rate $\geq 50$ breaths/min  | Sets: fast breathing = yes (used by rule IMCI-RESP-PNEUMONIA-FAST-BREATHING) | Not a classification rule - threshold check that feeds the respiratory classification rules below                                                            | p.6 (chart 2 of 76) | [ ]      |
| <b>IMCI-RESP-FAST-BREATHING-12-60M</b>   | Cough or difficult breathing | Child aged 12 to 60 months with respiratory rate $\geq 40$ breaths/min | Sets: fast breathing = yes (used by rule IMCI-RESP-PNEUMONIA-FAST-BREATHING) | Not a classification rule - threshold check that feeds the respiratory classification rules below                                                            | p.6 (chart 2 of 76) | [ ]      |
| <b>IMCI-RESP-SEVERE-DANGER-SIGN</b>      | Cough or difficult breathing | Any general danger sign is present                                     | Severe pneumonia or very severe disease                                      | Give first dose of appropriate antibiotic; Urgent referral                                                                                                   | p.6 (chart 2 of 76) | [ ]      |

| Rule ID                                    | Clinical Area                | If (condition)                                                                                                                                            | Then                                    | Actions / Note                                                                                                                                                                                                                                                           | WHO page            | Correct? |
|--------------------------------------------|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|----------|
| <b>IMCI-RESP-SEVERE-STRIDOR</b>            | Cough or difficult breathing | Child has: stridor when calm                                                                                                                              | Severe pneumonia or very severe disease | Give first dose of appropriate antibiotic; Urgent referral                                                                                                                                                                                                               | p.6 (chart 2 of 76) | [ ]      |
| <b>IMCI-RESP-PNEUMONIA-CHEST-INDRAWING</b> | Cough or difficult breathing | Child has: chest indrawing                                                                                                                                | Pneumonia                               | Give oral amoxicillin for 5 days; Soothe throat and relieve cough; Advise when to return immediately; Follow up in 3 days                                                                                                                                                | p.6 (chart 2 of 76) | [ ]      |
| <b>IMCI-RESP-PNEUMONIA-FAST-BREATHING</b>  | Cough or difficult breathing | Fast breathing detected (threshold met)                                                                                                                   | Pneumonia                               | Give oral amoxicillin for 5 days; Soothe throat and relieve cough; Advise when to return immediately; Follow up in 3 days                                                                                                                                                | p.6 (chart 2 of 76) | [ ]      |
| <b>IMCI-RESP-COUGH-OR-COLD</b>             | Cough or difficult breathing | None of the above conditions met (fallback)                                                                                                               | Cough or cold                           | Soothe throat and relieve cough; Advise when to return immediately; Follow up in 5 days if not improving                                                                                                                                                                 | p.6 (chart 2 of 76) | [ ]      |
| <b>IMCI-DIARRHOEA-SEVERE-DEHYDRATION</b>   | Diarrhoea dehydration        | At least 2 of: lethargic or unconscious = yes; sunken eyes = yes; drinking status in (Unable to drink, Drinks poorly); skin pinch = Goes back very slowly | Severe dehydration                      | If no other severe classification: Give fluid for severe dehydration (Plan C)<br><br>If other severe classification present: Urgent referral; Frequent ORS sips during referral; Continue breastfeeding                                                                  | p.7 (chart 3 of 76) | [ ]      |
| <b>IMCI-DIARRHOEA-SOME-DEHYDRATION</b>     | Diarrhoea dehydration        | At least 2 of: restless or irritable = yes; sunken eyes = yes; drinking status = Drinks eagerly / thirsty; skin pinch = Goes back slowly                  | Some dehydration                        | If no other severe classification: Give fluid, zinc and food (Plan B); Advise when to return immediately; Follow up in 5 days if not improving<br><br>If other severe classification present: Urgent referral; Frequent ORS sips during referral; Continue breastfeeding | p.7 (chart 3 of 76) | [ ]      |
| <b>IMCI-DIARRHOEA-NO-DEHYDRATION</b>       | Diarrhoea dehydration        | None of the above conditions met (fallback)                                                                                                               | No dehydration                          | Give fluid, zinc and food (Plan A); Advise when to return immediately; Follow up in 5 days if not improving                                                                                                                                                              | p.7 (chart 3 of 76) | [ ]      |