

Tensor Calculus
J.L. Synge and A.Schild (Dover Publication)
Solutions to exercices

Bernard Carrette

December 1, 2021

Remarks and warnings

Some notation conventions

$$\partial_r a_{mn} \equiv \frac{\partial a_{mn}}{\partial x^r}$$

$$\Gamma_{mn}^r \equiv \left\{ \begin{matrix} r \\ mn \end{matrix} \right\} \quad \text{Christoffel symbol of the second kind}$$

Contents

1	Spaces and Tensors	3
1.1	p5-exercise	4
1.2	p6-exercise	6
1.3	p8-exercise	7
1.4	p8-exercise	8
1.5	p8-clarification on expression 1.210	9
1.6	p9-exercise	10
1.7	p11-exercise	11
1.8	p12-exercise	12
1.9	p14-exercise	13
1.10	p16-exercise	15
1.11	p16-exercise	16
1.12	p18-exercise	17
1.13	p19-exercise	18
1.14	p19-exercise	19
1.15	p21-exercise	20
1.16	p23-exercise 1.	22
1.17	p23-exercise 2.	23
1.18	p23-exercise 3.	24
1.19	p23-exercise 4.	25
1.20	p23-exercise 5.	26
1.21	p23-exercise 6.	27
1.22	p24-exercise 7.	28
1.23	p24-exercise 8.	29
1.24	p24-exercise 9.	30
1.25	p23-exercise 10.	32
1.26	p23-exercise 11.	33
1.27	p23-exercise 12.	34
1.28	p23-exercise 13.	36
1.29	p23-exercise 14.	37
1.30	p23-exercise 15.	38
1.31	p23-exercise 16.	40

Spaces and Tensors

1.1 p5-exercise

The parametric equations of a hypersurface in V_n are

$$\begin{aligned} x^1 &= a \cos(u^1) \\ x^2 &= a \sin(u^1) \cos(u^2) \\ x^3 &= a \sin(u^1) \sin(u^2) \cos(u^3) \\ &\vdots \\ x^{N-1} &= a \sin(u^1) \sin(u^2) \sin(u^3) \dots \sin(u^{N-2}) \cos(u^{N-1}) \\ x^N &= a \sin(u^1) \sin(u^2) \sin(u^3) \dots \sin(u^{N-2}) \sin(u^{N-1}) \end{aligned}$$

where a is a constant. Find the single equation of the hyperspace in the form 1.103.

We have:

$$\begin{aligned} (x^N)^2 + (x^{N-1})^2 &= a^2 \prod_{i=1}^{N-2} \sin^2(u^i) (\cos^2(u^{N-1}) + \sin^2(u^{N-1})) \\ &= a^2 \prod_{i=1}^{N-2} \sin^2(u^i) \\ &= a^2 \prod_{i=1}^{N-3} \sin^2(u^i) \sin^2(u^{N-2}) \\ &= a^2 \prod_{i=1}^{N-3} \sin^2(u^i) (1 - \cos^2(u^{N-2})) \\ &= a^2 \prod_{i=1}^{N-3} \sin^2(u^i) - a^2 \prod_{i=1}^{N-3} \sin^2(u^i) \cos^2(u^{N-2}) \\ &= a^2 \prod_{i=1}^{N-3} \sin^2(u^i) - (x^{N-2})^2 \end{aligned}$$

giving

$$(x^N)^2 + (x^{N-1})^2 + (x^{N-2})^2 = a^2 \prod_{i=1}^{N-3} \sin^2(u^i)$$

In general, by recursion

$$\sum_{i=0}^k (x^{N-i})^2 = a^2 \prod_{i=1}^{N-k-1} \sin^2(u^i) \quad (k \leq N-2)$$

be $k = N - 2$ ($N - k - 1 = 1$) and in the left term put $j = N - i$ (j goes from 2 to N), we get

$$\begin{aligned}\sum_{j=2}^N (x^j)^2 &= a^2 \prod_{i=1}^1 \sin^2(u^i) \\ &= a^2 (1 - \cos^2(u^1)) \\ &= a^2 - (x^1)^2\end{aligned}$$

and thus the equation of the hyperspace is given by

$$\sum_{j=1}^N (x^j)^2 - a^2 = 0$$

Determine whether the points $(\frac{1}{2}a, 0, 0, \dots, 0)$, $(0, 0, \dots, 0, 2a)$ lie on the same or opposite sides of the hyperspace.

For $(\frac{1}{2}a, 0, 0, \dots, 0)$ we have $\sum_{j=1}^N (x^j)^2 - a^2 = -\frac{3a^2}{4} < 0$ and for $(0, 0, \dots, 0, 2a)$ we have $\sum_{j=1}^N (x^j)^2 - a^2 = \frac{3a^2}{4} > 0$.

So the points lie on opposite sides of the hyperplane.



1.2 p6-exercise

Let U_2 and W_2 be subspaces of V_N . Show that if $N = 3$ they will in general intersect in a curve; if $N = 4$ they will in general intersect in a finite number of points; and if $N > 4$ they will not in general intersect at all.

We have (see 1.102 page 5): $x^r = f^r(u^1, u^2, \dots, u^M) \quad (r = 1, 2, \dots, N)$

Case $N=3$:

For U_2 we have:

$$x^r = \phi^r(u^1, u^2) \quad (r = 1, 2, 3)$$

For W_2 we have:

$$x^r = \psi^r(v^1, v^2) \quad (r = 1, 2, 3)$$

The intersect of the two hyperplanes is given by the N equations:

$$\phi^r(u^1, u^2) = \psi^r(v^1, v^2) \quad (r = 1, 2, 3)$$

So we have 3 equations in 4 unknown u^1, u^2, v^1, v^2 and can choose (fix) one e.g. u^1 and solve the set of equations for u^2, v^1, v^2 giving

$$x^r = \theta^r(u^1) \quad (r = 1, 2, 3)$$

This is an equation of a curve in space (1 parameter equation)

Case $N=4$:

Using the same reasoning as with $N=3$, we get 4 equations for 4 unknown u^1, u^2, v^1, v^2 .

Provided that the set of equation does not degenerate, these 4 equations will determine u^1, u^2, v^1, v^2 without any degree of freedom. So we get points as solutions. This solution does not to be unique, e.g. if the $\phi^r(u^1, u^2)$ are quadratic form, then the solutions

$$(u^1, u^2, v^1, v^2)$$

$$(-u^1, u^2, v^1, v^2)$$

$$(u^1, -u^2, v^1, v^2)$$

$$(-u^1, -u^2, v^1, v^2)$$

are possible.

Case $N=5$: There are more equations than variables. If the equations are not linear dependent, no solutions will be found.



1.3 p8-exercise

Show that $(a_{rst} + a_{str} + a_{srt})x^r x^s x^t = 3a_{rst}x^r x^s x^t$

$(a_{rst} + a_{str} + a_{srt})x^r x^s x^t = a_{rst}x^r x^s x^t + a_{rts}x^r x^s x^t + a_{srt}x^r x^s x^t$ so by just renaming the dummy indices e.g. for the second term $r \mapsto s$, $s \mapsto t$ and $t \mapsto r$ we get the desired result.



1.4 p8-exercise

If $\phi = a_{rs}x^r x^s$, show that

$$\frac{\partial \phi}{\partial x^r} = (a_{rs} + a_{sr})x^s \quad , \quad \frac{\partial^2 \phi}{\partial x^r \partial x^s} = a_{rs} + a_{sr}$$

Simplify these expressions in the case where $a^{rs} = a^{sr}$

We have

$$\frac{\partial \phi}{\partial x^t} = \frac{\partial a_{rs}}{\partial x^t} x^r x^s + a_{rs} \frac{\partial x^r}{\partial x^t} x^s + a_{rs} x^r \frac{\partial x^s}{\partial x^t} \quad (1)$$

$$= \frac{\partial a_{rs}}{\partial x^t} x^r x^s + a_{rs} \delta_t^r x^s + a_{rs} x^r \delta_t^s \quad (2)$$

$$= \frac{\partial a_{rs}}{\partial x^t} x^r x^s + a_{ts} x^s + a_{rt} x^r \quad (3)$$

$$= \frac{\partial a_{rs}}{\partial x^t} x^r x^s + a_{ts} x^s + a_{st} x^s \quad (\text{rename dummy variable in third term}) \quad (4)$$

$$= \frac{\partial a_{rs}}{\partial x^t} x^r x^s + (a_{ts} + a_{st})x^s \quad (5)$$

Replace x^t by x^r , we get

$$\frac{\partial \phi}{\partial x^r} = \frac{\partial a_{rs}}{\partial x^r} x^r x^s + (a_{rs} + a_{sr})x^s \quad (6)$$

So the asked expression is only true if a_{rs} is not a function of the x^s . Assuming that a_{rs} is not a function of the x^s , take the partial derivative of (6) with respect to x^t , we get

$$\frac{\partial^2 \phi}{\partial x^r \partial x^t} = (a_{rs} + a_{sr}) \frac{\partial x^s}{\partial x^t} \quad (7)$$

$$= (a_{rs} + a_{sr}) \delta_t^s \quad (8)$$

$$= (a_{rt} + a_{tr}) \quad (9)$$

Replace x^t by x^s , and we get the proposed expression.



1.5 p8-clarification on expression 1.210

$$\frac{\partial^2 x^{,q}}{\partial x^p \partial x^s} + \frac{\partial^2 x^r}{\partial x^{,m} \partial x^{,n}} \frac{\partial x^{,m}}{\partial x^p} \frac{\partial x^{,n}}{\partial x^s} \frac{\partial x^{,q}}{\partial x^r} = 0$$

From 1.209:

$$\frac{\partial^2 x^r}{\partial x^{,m} \partial x^{,n}} \frac{\partial x^{,m}}{\partial x^p} \frac{\partial x^{,n}}{\partial x^s} + \frac{\partial x^r}{\partial x^{,n}} \frac{\partial^2 x^{,n}}{\partial x^p \partial x^s} = 0 \quad (1)$$

multiply (1) with $\frac{\partial x^{,q}}{\partial x^r}$

$$\frac{\partial^2 x^r}{\partial x^{,m} \partial x^{,n}} \frac{\partial x^{,m}}{\partial x^p} \frac{\partial x^{,n}}{\partial x^s} \frac{\partial x^{,q}}{\partial x^r} + \frac{\partial x^r}{\partial x^{,n}} \frac{\partial^2 x^{,n}}{\partial x^p \partial x^s} \frac{\partial x^{,q}}{\partial x^r} = 0 \quad (2)$$

$$\Leftrightarrow \frac{\partial x^r}{\partial x^{,n}} \frac{\partial^2 x^{,n}}{\partial x^p \partial x^s} \frac{\partial x^{,q}}{\partial x^r} + \frac{\partial^2 x^r}{\partial x^{,m} \partial x^{,n}} \frac{\partial x^{,m}}{\partial x^p} \frac{\partial x^{,n}}{\partial x^s} \frac{\partial x^{,q}}{\partial x^r} = 0 \quad (3)$$

$$\text{in the first term we get} \quad \frac{\partial x^{,q}}{\partial x^r} \frac{\partial x^r}{\partial x^{,n}} = \frac{\partial x^{,q}}{\partial x^{,n}} = \delta_n^q \quad (4)$$

(3) becomes

$$\frac{\partial^2 x^{,n}}{\partial x^p \partial x^s} \delta_n^q + \frac{\partial^2 x^r}{\partial x^{,m} \partial x^{,n}} \frac{\partial x^{,m}}{\partial x^p} \frac{\partial x^{,n}}{\partial x^s} \frac{\partial x^{,q}}{\partial x^r} = 0 \quad (5)$$

$$\Leftrightarrow \frac{\partial^2 x^{,q}}{\partial x^p \partial x^s} + \frac{\partial^2 x^r}{\partial x^{,m} \partial x^{,n}} \frac{\partial x^{,m}}{\partial x^p} \frac{\partial x^{,n}}{\partial x^s} \frac{\partial x^{,q}}{\partial x^r} = 0 \quad (6)$$



1.6 p9-exercise

If A_s^r are the elements of a determinant A, and B_s^r the elements of a determinant B, show that the element of the product determinant is $A_n^r B_s^n$. Hence show that the product of the two jacobians

$$J = \left| \frac{\partial x^r}{\partial x^s} \right|, \quad J' = \left| \frac{\partial x'^r}{\partial x^s} \right|$$

is unity.

Remark: Some nitpick about the formulation: A_s^r are not the elements of a determinant A, but elements of the matrix A which gives $\det\{A\}$ provided that A is square (which is not explicitly mentioned.). The same remark for B and $A_n^r B_s^n$.

Be A_k^i the elements of matrix A and B_j^k the elements of matrix B and $C = A.B$ the resulting matrix of the multiplication of A and B, then

$$C_j^i = A_k^i B_j^k$$

are the elements of matrix C. Now, put $A_k^i = \frac{\partial x^i}{\partial x'^k}$ and $B_j^k = \frac{\partial x'^k}{\partial x^j}$ then,

$$C_j^i = A_k^i B_j^k \tag{1}$$

$$= \frac{\partial x^i}{\partial x'^k} \frac{\partial x'^k}{\partial x^j} \tag{2}$$

$$= \delta_k^i \tag{3}$$

So $C = JJ'$ becomes the unity matrix.



1.7 p11-exercise

Show that a finite contravariant vector determines the ratios of the components of an infinitesimal displacement. (Consider the transformation of the equation $dx^r = \theta T^r$, where θ is an arbitrary factor which does not change under the transformation. Alternatively, show that the equations $T^r dx^s - T^s dx^r = 0$ remain true when we transform the coordinates.)

Be T^q a contravariant vector.

$$T^{,q} = T^r \frac{\partial x^{,q}}{\partial x^r} \quad (\text{by definition}) \quad (1)$$

Be θ a small infinitesimal factor invariant for a coordinate transformation, define

$$dx^r = \theta T^r \quad (2)$$

$$(3)$$

then

$$\frac{dx^r}{dx^s} = \frac{\theta T^r}{\theta T^s} \quad (4)$$

$$\Leftrightarrow T^s dx^r - T^r dx^s = 0 \quad (5)$$

Alternatively, multiply (5) with $\partial_{x^r} x^{,q}$, then

$$\frac{\partial x^{,q}}{\partial x^r} dx^r T^s - \frac{\partial x^{,q}}{\partial x^r} dx^s T^r = 0 \quad (6)$$

$$\Leftrightarrow \frac{\partial x^{,q}}{\partial x^r} dx^r T^s - dx^s T^{,q} = 0 \quad (\text{use (1) in the second term}) \quad (7)$$

$$\Leftrightarrow dx^{,q} T^s - dx^s T^{,q} = 0 \quad (8)$$

$$(9)$$

Multiply (8) with $\partial_{x^s} x^{,p}$, then

$$dx^{,q} T^s \partial_{x^s} x^{,p} - dx^s T^{,q} \partial_{x^s} x^{,p} = 0 \quad (10)$$

$$\Leftrightarrow T^{,p} dx^{,q} - T^{,q} dx^{,p} = 0 \quad (\text{use (1) in the first term}) \quad (11)$$

and thus

$$\frac{dx^{,q}}{dx^{,p}} = \frac{T^{,q}}{T^{,p}}$$



1.8 p12-exercise

Write down the equation of transformation, analogous to 1.305, of a contravariant tensor of the third order. Solve the equation so as to express the unprimed components in terms of the primed components.

Be

$$T^{,uvw} = T^{rst} \frac{\partial x^{,u}}{\partial x^r} \frac{\partial x^{,v}}{\partial x^s} \frac{\partial x^{,w}}{\partial x^t} \quad (\text{by definition}) \quad (1)$$

a contravariant vector.

Multiply (1) by $\frac{\partial x^n}{\partial x^{,u}}$

$$T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} = T^{rst} \frac{\partial x^{,u}}{\partial x^r} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^{,v}}{\partial x^s} \frac{\partial x^{,w}}{\partial x^t} \quad (2)$$

$$\Leftrightarrow T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} = T^{rst} \delta_r^n \frac{\partial x^{,v}}{\partial x^s} \frac{\partial x^{,w}}{\partial x^t} \quad (3)$$

$$\Leftrightarrow T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} = T^{nst} \frac{\partial x^{,v}}{\partial x^s} \frac{\partial x^{,w}}{\partial x^t} \quad (4)$$

Multiply (4) by $\frac{\partial x^m}{\partial x^{,v}}$

$$T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} = T^{nst} \frac{\partial x^{,v}}{\partial x^s} \frac{\partial x^m}{\partial x^{,v}} \frac{\partial x^{,w}}{\partial x^t} \quad (5)$$

$$\Leftrightarrow T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} = T^{nst} \delta_s^m \frac{\partial x^{,w}}{\partial x^t} \quad (6)$$

$$\Leftrightarrow T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} = T^{nmt} \frac{\partial x^{,w}}{\partial x^t} \quad (7)$$

Multiply (7) by $\frac{\partial x^p}{\partial x^{,w}}$

$$T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} \frac{\partial x^p}{\partial x^{,w}} = T^{nmt} \frac{\partial x^{,w}}{\partial x^t} \frac{\partial x^p}{\partial x^{,w}} \quad (8)$$

$$\Leftrightarrow T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} \frac{\partial x^p}{\partial x^{,w}} = T^{nmt} \delta_t^p \quad (9)$$

$$\Leftrightarrow T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} \frac{\partial x^p}{\partial x^{,w}} = T^{nmp} \quad (10)$$

Giving

$$T^{nmp} = T^{,uvw} \frac{\partial x^n}{\partial x^{,u}} \frac{\partial x^m}{\partial x^{,v}} \frac{\partial x^p}{\partial x^{,w}}$$



1.9 p14-exercise

For a transformation from on set of rectangular Cartesian coordinates to another in Euclidean 3-space, show that the law of transformation of a contravariant vector is precisely the same as that of a covariant vector. Can this statements be extended to cover tensor of higher orders?

We have to prove that, given that,

$$T^{,i} = T^j \frac{\partial x^{,i}}{\partial x^j} \quad T_i = T_j \frac{\partial x^j}{\partial x^{,i}}$$

that also

$$T^{,i} = T^j \frac{\partial x^j}{\partial x^{,i}} \quad T_i = T_j \frac{\partial x^{,i}}{\partial x^j} \quad (1)$$

$$\Leftrightarrow \frac{\partial x^j}{\partial x^{,i}} = \frac{\partial x^{,i}}{\partial x^j} \quad (2)$$

Be

$$\hat{e}^{,i} = g_k^i \hat{e}^k \quad \text{and} \quad \hat{e}^i = h_k^i \hat{e}^{,k} \quad (3)$$

the transformation rules from one set of (rectangular Cartesian) basis vectors to another set of (rectangular Cartesian) basis vectors. Then,

$$\langle \hat{e}^{,i}, \hat{e}^{,j} \rangle = \langle g_k^i \hat{e}^k, g_k^j \hat{e}^k \rangle \quad \text{and} \quad \langle \hat{e}^i, \hat{e}^j \rangle = \langle h_k^i \hat{e}^{,k}, h_k^j \hat{e}^{,k} \rangle \quad (4)$$

$$\Leftrightarrow \delta_j^p = g_k^p g_k^j \quad \text{and} \quad \delta_j^p = h_k^p h_k^j \quad (5)$$

$$(6)$$

Be \vec{v} a random vector in the Euclidean space,

$$\vec{v} = x^j \hat{e}^j = x^{,j} \hat{e}^{,j} \quad (7)$$

then

$$(3) \Rightarrow x^j \hat{e}^j = x^j h_k^j \hat{e}^{,k} \quad \text{and} \quad x^{,j} \hat{e}^{,j} = x^{,j} g_k^j \hat{e}^k \quad (8)$$

$$\Rightarrow x^{,j} = x^m h_j^m \quad \text{and} \quad x^m = x^{,j} g_m^j \quad (9)$$

$$\Rightarrow x^{,j} = x^{,i} g_m^i h_j^m \quad \text{and} \quad x^m = x^k h_j^k g_m^j \quad (10)$$

$$\Rightarrow \delta_j^p = g_k^p h_j^k \quad \text{and} \quad \delta_j^p = g_j^k h_k^p \quad (11)$$

$$(5) \Rightarrow g_k^p g_k^j = g_k^p h_j^k \quad \text{and} \quad h_k^p h_k^j = g_j^k h_k^p \quad (12)$$

$$\Rightarrow g_k^j = h_j^k \quad \text{and} \quad h_k^j = g_j^k \quad (13)$$

From (9)

$$x^j = x^{,m} g_j^m \text{ and } x^{,k} = x^n h_k^n \quad (14)$$

$$\Rightarrow \frac{\partial x^{,k}}{\partial x^j} = \frac{\partial x^n}{\partial x^j} h_k^n \text{ and } \frac{\partial x^j}{\partial x^{,k}} = \frac{\partial x^{,m}}{\partial x^{,k}} g_j^m \quad (15)$$

$$\Leftrightarrow \frac{\partial x^{,k}}{\partial x^j} = \delta_j^n h_k^n \text{ and } \frac{\partial x^j}{\partial x^{,k}} = \delta_k^m g_j^m \quad (16)$$

$$\Leftrightarrow \frac{\partial x^{,k}}{\partial x^j} = h_k^j \text{ and } \frac{\partial x^j}{\partial x^{,k}} = g_j^k \quad (17)$$

$$(13) \Rightarrow \frac{\partial x^{,k}}{\partial x^j} = \frac{\partial x^j}{\partial x^{,k}} \quad (18)$$

So (13) matches (2), proving the assertion.

Can this statements be extended to cover tensor of higher orders? Consider

$$T^{i,j,\dots,n} = T^{r,s,\dots,w} \frac{\partial x^{,i}}{\partial x^r} \frac{\partial x^{,j}}{\partial x^s} \dots \frac{\partial x^{,n}}{\partial x^w} \text{ and } T^{r,s,\dots,w} = T^{i,j,\dots,n} \frac{\partial x^r}{\partial x^{,i}} \frac{\partial x^s}{\partial x^{,j}} \dots \frac{\partial x^w}{\partial x^{,n}}$$

Using the same reasoning as in (1) to (2) we need

$$\frac{\partial x^{,i}}{\partial x^r} \frac{\partial x^{,j}}{\partial x^s} \dots \frac{\partial x^{,n}}{\partial x^w} = \frac{\partial x^r}{\partial x^{,i}} \frac{\partial x^s}{\partial x^{,j}} \dots \frac{\partial x^w}{\partial x^{,n}}$$

As the conclusion (18) is independent of the order of the tensor, it is obvious that the above equality yields. Hence, the answer is YES.



1.10 p16-exercise

In a space of 4 dimensions, the tensor A_{rst} is skew-symmetric in the last pair of suffixes. Show that only 24 of the 64 components may be chosen arbitrarily. If the further condition $A_{rst} + A_{str} + A_{trs} = 0$ is imposed, show that that only 20 components may be chosen arbitrarily.

We have, as A is skew-symmetric in the last pair of suffixes

$$A_{rst} = -A_{rts} \Rightarrow s = t: A_{rst} = 0$$

So, for each r (4 possible choices as $N = 4$) we have $4 \times 4 / 2 - 4 = 6$ degrees of freedom. [we have the term $4 \times 4 / 2$ as the tensor is (skew-)symmetric, e.g. once we choose element a_{12} , then a_{21} is also known. The term -4 takes into account the diagonal element which are 0 and thus cannot be chosen.] So, we have $4 \times 6 = 24$ degrees of freedom.

What about the supplementary constraint $A_{rst} + A_{str} + A_{trs} = 0$:

Consider the two possible excluding cases:

$$\text{i) } r = s \neq t \quad (\Leftrightarrow r = t \neq s)$$

This case gives - without the additional constraint (1) - $4 \times (4 \times 3 / 2 - 4) = 8$ degrees of freedom. Does the constraint (1) reduces this degree of freedom?

We have,

$$A_{rst} + A_{str} + A_{trs} = 0 \quad (1)$$

$$\Rightarrow \underbrace{A_{rrt} + A_{rtt}}_{= 0 \text{ (non-diagonal terms)}} + \underbrace{A_{trr}}_{= 0 \text{ (diagonal terms)}} = 0 \quad (2)$$

So, no additional constraints are added by (1) to the restriction i) and the DOF remains 8.

$$\text{ii) } t \neq r \neq s \neq t$$

This case means that we have to choose a set of 3 elements out of 4 elements without repetition. This a *variation* of 3 elements out of 4.

$$V_k^n = \frac{n!}{(n-k)!} \text{ giving } V_3^4 = \frac{4!}{(4-3)!} = 24$$

The constraint (1) gives us 24 equations but as $A_{rst} = -A_{rts}$ only 12 equations have to be considered. So, with the additional constraints the DOF becomes $24 - 12 = 12$.

As i) and ii) are independent and excluding events we can add the DOF of both events and we get $8 + 12 = 20$ DOF.



1.11 p16-exercise

If A^{rs} is skew-symmetric and B_{rs} is symmetric, prove that $A^{rs}B_{rs} = 0$. Hence show that the quadratic form $a_{ij}x^i x^j$ is unchanged if a_{ij} is replaced by its symmetric part.

We can split the summation $A^{rs}B_{rs}$ in three subsummations:

$$A^{rs}B_{rs} = A^{rs}B_{rs}|_{r=s} \tag{1}$$

$$+ A^{rs}B_{rs}|_{r>s} \tag{2}$$

$$+ A^{rs}B_{rs}|_{r<s} \tag{3}$$

We have:

$$(1) = 0 \text{ as } A^{kk} = 0 \text{ (skew-symmetric)}$$

$$(2)+(3) = A^{rs}B_{rs}|_{r>s} + A^{rs}B_{rs}|_{r<s}$$

As $A^{rs} = -A^{sr}$ and $B^{rs} = B^{sr}$ we can write (2)+(3) as :

$$A^{rs}B_{rs}|_{r>s} + (-A^{sr})B_{sr}|_{r>s} = 0$$

So, $A^{rs}B_{rs} = 0$

Consider the quadratic form $\phi = a_{ij}x^i x^j$

Be $A_{ij} = (a_{ij})$ and $B_{ij} = (x^i x^j)$, then it is obvious that B_{ij} is symmetric and that $C_{ij} = -A_{ij}$ is the form where $-a_{ij}$ is replaced by its symmetric part (skew-symmetric). Hence $\phi = a_{ij}x^i x^j = a_{ij}b^{ij} = 0$ and so is $\phi = c_{ij}b^{ij} = 0$



1.12 p18-exercise

What are the values (in a space of N dimensions) of the following contractions formed from the Kronecker delta?

$$\delta_m^m, \delta_n^m \delta_m^n, \delta_n^m \delta_r^n \delta_m^r$$

We can split the summation $A^{rs} B_{rs}$ in three subsummations:

$$\delta_m^m = N \tag{1}$$

$$\delta_n^m \delta_m^n = \delta_m^m = N \tag{2}$$

$$\delta_n^m \delta_r^n \delta_m^r = \delta_n^m \delta_m^n = \delta_m^m = N \tag{3}$$



1.13 p19-exercise

If X^r, Y^r are arbitrary contravariant vectors and $a_{rs}X^rY^s$ is an invariant, then a_{rs} are the components of a covariant tensor of the second order.

We have to prove that

$$a'_{rs} = a_{ij} \frac{\partial x^i}{\partial x'^r} \frac{\partial x^j}{\partial x'^s} \text{ or } a_{ij} = a'_{rs} \frac{\partial x'^r}{\partial x^i} \frac{\partial x'^s}{\partial x^j} \quad (1)$$

$a_{rs}X^rY^s$ is an invariant, means

$$a'_{rs}X'^rY'^s = a_{rs}X^rY^s \quad (2)$$

As X^r, Y^r are arbitrary contravariant vectors, we have

$$X'^r = X^i \frac{\partial x'^r}{\partial x^i} \text{ and } Y'^s = Y^j \frac{\partial x'^s}{\partial x^j} \quad (3)$$

(3) in (2) gives

$$a'_{rs}X^i \frac{\partial x'^r}{\partial x^i} Y^j \frac{\partial x'^s}{\partial x^j} = a_{rs}X^rY^s \quad (4)$$

$$\Leftrightarrow a'_{rs} \frac{\partial x'^r}{\partial x^i} \frac{\partial x'^s}{\partial x^j} X^i Y^j = a_{ij} X^i Y^j \quad (5)$$

$$\Leftrightarrow (a'_{rs} \frac{\partial x'^r}{\partial x^i} \frac{\partial x'^s}{\partial x^j} - a_{ij}) X^i Y^j = 0 \quad (6)$$

As X^r, Y^r are arbitrary contravariant vectors, we conclude that

$$a'_{rs} \frac{\partial x'^r}{\partial x^i} \frac{\partial x'^s}{\partial x^j} - a_{ij} = 0 \quad (7)$$

$$\Leftrightarrow a_{ij} = a'_{rs} \frac{\partial x'^r}{\partial x^i} \frac{\partial x'^s}{\partial x^j} \quad (8)$$

(8) = (1): OK



1.14 p19-exercise

If X_{rs} is an arbitrary covariant tensor of the second order, and $A_r^{mn} X_{mn}$ is a covariant vector, then A_r^{mn} has the mixed tensor character indicated by the positions of its suffixes

We have to prove that

$$A_r^{vw} = A_k^{mn} \frac{\partial x^k}{\partial x^{,r}} \frac{\partial x^{,v}}{\partial x^m} \frac{\partial x^{,w}}{\partial x^n} \quad (1)$$

We have

$$P_r = A_r^{mn} X_{mn} \quad (2)$$

is a covariant vector

$$\Rightarrow P_r^{,} = A_k^{mn} X_{mn} \frac{\partial x^k}{\partial x^{,r}} \quad (3)$$

but X_{mn} is a covariant tensor

$$\Rightarrow X_{mn} = X_{ps} \frac{\partial x^{,p}}{\partial x^m} \frac{\partial x^{,s}}{\partial x^n} \quad (4)$$

So (4) in (3) gives

$$P_r^{,} = A_k^{mn} X_{ps} \frac{\partial x^{,p}}{\partial x^m} \frac{\partial x^{,s}}{\partial x^n} \frac{\partial x^k}{\partial x^{,r}} \quad (5)$$

$$\Leftrightarrow P_r^{,} = A_k^{mn} \underbrace{\frac{\partial x^{,p}}{\partial x^m} \frac{\partial x^{,s}}{\partial x^n} \frac{\partial x^k}{\partial x^{,r}}}_{(*)} X_{ps} \quad (6)$$

Putting (*) as $A_r^{ps} = A_k^{mn} \frac{\partial x^{,p}}{\partial x^m} \frac{\partial x^{,s}}{\partial x^n} \frac{\partial x^k}{\partial x^{,r}}$ we see that (6) has the form (2) and that A_r^{ps} obeys the rule of a mixed tensor (1).



1.15 p21-exercise

If A_{rs} is a skew-symmetric covariant tensor, prove that B_{rst} defined as

$$B_{rst} = \partial_r A_{st} + \partial_s A_{tr} + \partial_t A_{rs}$$

is a covariant tensor, and that it is skew-symmetric in all pairs of suffixes.

We have A_{rs} is a covariant tensor

$$A_{ij} = A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^i} \frac{\partial x^\beta}{\partial x^j} \quad (1)$$

$$\Rightarrow B_{rst} = \partial_r (A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t}) + \partial_s (A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^t} \frac{\partial x^\beta}{\partial x^r}) + \partial_t (A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s}) \quad (2)$$

Note that

$$\partial_k (A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t}) = \partial_k (A_{\alpha\beta}) \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t} + A_{\alpha\beta} \partial_k (\frac{\partial x^\alpha}{\partial x^s}) \frac{\partial x^\beta}{\partial x^t} + A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^s} \partial_k (\frac{\partial x^\beta}{\partial x^t}) \quad (3)$$

$$(4)$$

so,

$$\begin{aligned} B_{rst} &= \partial_r A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t} + \underbrace{A_{\alpha\beta} \partial_r \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t}}_{*} + \underbrace{A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^s} \partial_r \frac{\partial x^\beta}{\partial x^t}}_{**} \\ &\quad + \partial_s A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^t} \frac{\partial x^\beta}{\partial x^r} + \underbrace{A_{\alpha\beta} \partial_s \frac{\partial x^\alpha}{\partial x^t} \frac{\partial x^\beta}{\partial x^r}}_{***} + \underbrace{A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^t} \partial_s \frac{\partial x^\beta}{\partial x^r}}_{*} \\ &\quad + \partial_t A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s} + \underbrace{A_{\alpha\beta} \partial_t \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s}}_{**} + \underbrace{A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^r} \partial_t \frac{\partial x^\beta}{\partial x^s}}_{***} \end{aligned} \quad (5)$$

In (5) consider the two terms with (*)

$$T = A_{\alpha\beta} \partial_r \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t} + A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^t} \partial_s \frac{\partial x^\beta}{\partial x^r} \quad (6)$$

$$= A_{\alpha\beta} \frac{\partial^2 x^\alpha}{\partial x^s \partial x^r} \frac{\partial x^\beta}{\partial x^t} + A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^t} \frac{\partial^2 x^\beta}{\partial x^r \partial x^s} \quad (7)$$

$$= A_{\alpha\beta} \frac{\partial^2 x^\alpha}{\partial x^s \partial x^r} \frac{\partial x^\beta}{\partial x^t} + A_{\beta\alpha} \frac{\partial x^\beta}{\partial x^t} \frac{\partial^2 x^\alpha}{\partial x^r \partial x^s} \text{ (by renaming dummy variables)} \quad (8)$$

As $A_{ij} = -A_{ji}$ (skew-symmetric tensor), we get $T = 0$. The same yields for the (**) and (***) terms. So, B_{rst} reduces to

$$B_{rst} = \partial_r A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t} + \partial_s A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^t} \frac{\partial x^\beta}{\partial x^r} + \partial_t A_{\alpha\beta} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s} \quad (9)$$

$$\Leftrightarrow B_{rst} = \frac{\partial A_{\alpha\beta}}{\partial x^\gamma} \frac{\partial x^\gamma}{\partial x^r} \frac{\partial x^\alpha}{\partial x^s} \frac{\partial x^\beta}{\partial x^t} + \frac{\partial A_{\alpha\beta}}{\partial x^\gamma} \frac{\partial x^\gamma}{\partial x^s} \frac{\partial x^\alpha}{\partial x^t} \frac{\partial x^\beta}{\partial x^r} + \frac{\partial A_{\alpha\beta}}{\partial x^\gamma} \frac{\partial x^\gamma}{\partial x^t} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s} \quad (10)$$

By adequate renaming of the dummy variable in the 3 terms:

$$\begin{bmatrix} 1^{st} term \\ 2^{nd} term \\ 3^{rd} term \end{bmatrix} \longrightarrow \begin{bmatrix} \gamma \rightarrow \alpha & \alpha \rightarrow \beta & \beta \rightarrow \gamma \\ \beta \rightarrow \alpha & \gamma \rightarrow \beta & \alpha \rightarrow \gamma \\ \alpha \rightarrow \alpha & \beta \rightarrow \beta & \gamma \rightarrow \gamma \end{bmatrix}$$

we get

$$B_{rst} = \left(\frac{\partial A_{\beta\gamma}}{\partial x^\alpha} + \frac{\partial A_{\gamma\alpha}}{\partial x^\beta} + \frac{\partial A_{\alpha\beta}}{\partial x^\gamma} \right) \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s} \frac{\partial x^\gamma}{\partial x^t} \quad (11)$$

$$\Leftrightarrow B_{rst} = \underbrace{(\partial_\alpha A_{\beta\gamma} + \partial_\beta A_{\gamma\alpha} + \partial_\gamma A_{\alpha\beta})}_{(****)} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^\beta}{\partial x^s} \frac{\partial x^\gamma}{\partial x^t} \quad (12)$$

The expression (****) has exactly the required form $B_{rst} = \partial_r A_{st} + \partial_s A_{tr} + \partial_t A_{rs}$ and is transformed (12) according the rules of a covariant tensor.

Let's prove now that it is skew-symmetric in all pairs of suffixes. We have to consider the following permutations

$$\begin{bmatrix} rst \\ rts \\ srt \\ str \\ trs \\ tsr \end{bmatrix}$$

E.g. srt

$$B_{rts} = \partial_r A_{ts} + \partial_t A_{sr} + \partial_s A_{rt} \quad (13)$$

$$= -\partial_r A_{st} - \partial_t A_{rs} - \partial_s A_{tr} \quad (14)$$

$$= -B_{rst} \quad (15)$$

The same calculations can be done for the other permutations.



1.16 p23-exercise 1.

In a V_4 there are two 2-spaces with equations

$$x^r = f^r(u^1, u^2), \quad x^r = g^r(u^3, u^4)$$

Prove that if these 2-spaces have a curve of intersection, then the determinantal equation

$$\left| \frac{\partial x^r}{\partial u^s} \right| = 0$$

is satisfied along the curve.

Having a curve means that one of the parameters u^i can be freely chosen while the other 3 are determined by the chosen parameter.

We have,

$$\left| \frac{\partial x^r}{\partial u^s} \right| = \begin{vmatrix} \frac{\partial x^1}{\partial u^1} & \frac{\partial x^1}{\partial u^2} & \frac{\partial x^1}{\partial u^3} & \frac{\partial x^1}{\partial u^4} \\ \vdots & \vdots & \vdots & \vdots \\ \frac{\partial x^4}{\partial u^1} & \frac{\partial x^4}{\partial u^2} & \frac{\partial x^4}{\partial u^3} & \frac{\partial x^4}{\partial u^4} \end{vmatrix} \quad (1)$$

Suppose we choose u^4 as parameter. This means $u^i = \phi^i(u^4)$ for $i=1,2,3$ and thus we can write

$$\frac{\partial x^i}{\partial u^4} = \frac{\partial x^i}{\partial u^j} \frac{d\phi^j}{du^4} + \frac{\partial x^i}{\partial u^4} \quad \text{with } j=1,2,3 \quad i = 1,2,3,4 \quad (2)$$


$$\Rightarrow \frac{\partial x^i}{\partial u^j} \frac{d\phi^j}{du^4} = 0 \quad (3)$$

This means that in (1) the three first columns are not linearly independent and thus have $\left| \frac{\partial x^r}{\partial u^s} \right| = 0$



1.17 p23-exercise 2.

In Euclidean space of three dimensions, write down the equations of transformation between rectangular Cartesian coordinates x, y, z and spherical polar coordinates r, θ, ϕ . Find the Jacobian of the transformation. Where is it zero or infinite?



We use the latitude ψ instead of the co-latitude ϕ .

$$\begin{cases} x = r \cos(\psi) \cos(\theta) \\ y = r \cos(\psi) \sin(\theta) \\ z = r \sin(\psi) \end{cases}$$

Partial differentiating of (x, y, z) with respect to (r, ψ, θ) gives the Jacobian

$$J = \begin{vmatrix} \cos(\psi) \cos(\theta) & -r \sin(\psi) \cos(\theta) & -r \cos(\psi) \sin(\theta) \\ \cos(\psi) \sin(\theta) & -r \sin(\psi) \sin(\theta) & r \cos(\psi) \cos(\theta) \\ \sin(\psi) & r \cos(\psi) & 0 \end{vmatrix} \quad (1)$$

$$J = \cos(\psi) \cos(\theta) (-r^2) \cos^2(\psi) \cos(\theta) \quad (2)$$

$$+ r \sin(\psi) \cos(\theta) (-r \cos(\psi) \cos(\theta) \sin(\psi)) \quad (3)$$

$$- r \cos(\psi) \sin(\theta) (r \cos^2(\psi) \sin(\theta) + r \sin^2(\psi) \sin(\theta)) \quad (4)$$

$$= -r^2 \cos^3(\psi) \cos^2(\theta) - r^2 \sin^2(\psi) \cos^2(\theta) \cos(\psi) - r^2 \cos(\psi) \sin^2(\theta) \quad (5)$$

Noting that the 2nd term in (5) can be written as $-r^2 \cos^2(\theta) \cos(\psi) + r^2 \cos^2(\theta) \cos^3(\psi)$, we get

$$J = -r^2 (\cos^3(\psi) \cos^2(\theta) + \cos^2(\theta) \cos(\psi) - \cos^3(\psi) \cos^2(\theta) + \cos(\psi) \sin^2(\theta)) \quad (6)$$

$$= -r^2 \cos(\psi) \quad (7)$$

$J=0$: for $r = 0$ or $\psi = \frac{\pi}{2} |_{r \in (-\infty, +\infty)}$ and $J \rightarrow \pm\infty$ or $\mp\infty$ for $r \rightarrow \pm\infty |_{\psi \neq 0}$. But what about the case $r \rightarrow \pm\infty |_{\psi \rightarrow 0}$? This case is not determined as long as no path is chosen in the (r, ψ) configuration space.



1.18 p23-exercise 3.

If X, Y, Z are the components of a contravariant vector for rectangular Cartesian coordinates in Euclidean 3-space, find its components for spherical polar coordinates.

Be x^α the components of a contravariant vector in spherical polar coordinates and x^i its components in rectangular Cartesian coordinates. As we have

$$\begin{aligned} x^\rho &= \sqrt{x^j x^j} \\ x^\theta &= \text{atan} \frac{x^2}{x^1} \\ x^\phi &= \text{asin} \frac{x^3}{\sqrt{x^j x^j}} \end{aligned} \quad \text{and} \quad A^\alpha = A^i \frac{\partial x^\alpha}{\partial x^i} \quad (1)$$

$$\Rightarrow [A^\alpha] = \left[A^i \frac{\partial x^\alpha}{\partial x^i} \right] = \begin{bmatrix} \frac{x^1}{\sqrt{x^j x^j}} & \frac{x^2}{\sqrt{x^j x^j}} & \frac{x^3}{\sqrt{x^j x^j}} \\ -\frac{x^2}{(x^1)^2 + (x^2)^2} & \frac{x^1}{(x^1)^2 + (x^2)^2} & 0 \\ -\frac{x^3 x^1}{(x^j x^j) \sqrt{(x^1)^2 + (x^2)^2}} & -\frac{x^3 x^2}{(x^j x^j) \sqrt{(x^1)^2 + (x^2)^2}} & \frac{\sqrt{(x^1)^2 + (x^2)^2}}{(x^j x^j)} \end{bmatrix} \begin{bmatrix} A^1 \\ A^2 \\ A^3 \end{bmatrix} \quad (2)$$

◆

1.19 p23-exercise 4.

In a space of three dimensions, how many different expressions are represented by the product $A_{np}^m B_{rs}^{pq} C_{tu}^s$? How many terms occur in each such expression, when written out explicitly?

As we have V_3 and considering that in $A_{np}^m B_{rs}^{pq} C_{tu}^s$ the six indices m, n, q, r, t, u are not dummy indices, we get 3^6 different expressions (first choose m : you have three choices, then n : also three choices giving 3×3 possibilities, etc for q, r, t, u).

For the second question, as in $A_{np}^m B_{rs}^{pq}$ there is only summation on over index (p) we get three terms for this part. As the summation with $A_{np}^m B_{rs}^{pq}$ and C_{tu}^s occurs only on one index also (s) we get 3×3 terms in the expression.



1.20 p23-exercise 5.

If A is an invariant in V_n , are the second derivatives $\frac{\partial^2 A}{\partial x^r \partial x^s}$ the components of a tensor?

As A is invariant (note: different alphabets in the indices indicates different coordinate systems):

$$A(x^\rho) = A(x^i) \quad (1)$$

$$\Rightarrow \frac{\partial A(x^\rho)}{\partial x^i} = \frac{\partial A(x^j)}{\partial x^i} \quad (2)$$

To simplify the notation, we put $A(x^\rho) = A'$ and $A(x^j) = A'$ then (2) can be written as

$$\frac{\partial A'}{\partial x^\rho} \frac{\partial x^\rho}{\partial x^i} = \frac{\partial A}{\partial x^i} \quad (3)$$

Conclusion: $\frac{\partial A}{\partial x^i}$ is a covariant tensor.

Consider now $\frac{\partial A}{\partial x^i} = \frac{\partial A'}{\partial x^\rho} \frac{\partial x^\rho}{\partial x^i}$. Then,

$$\frac{\partial^2 A}{\partial x^i \partial x^j} = \frac{\partial^2 A'}{\partial x^\rho \partial x^j} \frac{\partial x^\rho}{\partial x^i} + \frac{\partial A'}{\partial x^\rho} \frac{\partial^2 x^\rho}{\partial x^i \partial x^j} \quad (4)$$

$$\Leftrightarrow \frac{\partial^2 A}{\partial x^i \partial x^j} = \frac{\partial^2 A'}{\partial x^\rho \partial x^\gamma} \frac{\partial x^\gamma}{\partial x^j} \frac{\partial x^\rho}{\partial x^i} + \frac{\partial A'}{\partial x^\rho} \frac{\partial^2 x^\rho}{\partial x^i \partial x^j} \quad (5)$$

The first term on the right side, behaves as covariant tensor but the presence of the second term makes that generally, $\frac{\partial^2 A}{\partial x^i \partial x^j}$ has not a tensor character. This is only when $\frac{\partial A'}{\partial x^\rho} \frac{\partial^2 x^\rho}{\partial x^i \partial x^j} = 0$, which means that x^ρ, x^i are a linear map of each other.



1.21 p23-exercise 6.

Suppose that in V_2 the components of a contravariant tensor field T^{mn} in a coordinate system x^r are

$$T^{11} = 1 \quad T^{12} = 0$$

$$T^{21} = 1 \quad T^{22} = 0$$

Find the components T^{mn} in a coordinate system x'^r , where

$$x'^1 = (x^1)^2 \quad x'^2 = (x^2)^2$$

Write down the values of these components in particular at the point $x^1 = 1, x^2 = 0$.

As we have a contravariant tensor field :

$$T'^{mn} = T^{ij} \frac{\partial x'^m}{\partial x^i} \frac{\partial x'^n}{\partial x^j} \quad (1)$$

$$\begin{aligned} x'^1 = (x^1)^2 &\Rightarrow \frac{\partial x'^1}{\partial x^1} = 2x^1 & \frac{\partial x'^1}{\partial x^2} &= 0 \\ x'^2 = (x^2)^2 &\Rightarrow \frac{\partial x'^2}{\partial x^1} &= 0 & \frac{\partial x'^2}{\partial x^2} = 2x^2 \end{aligned} \quad (2)$$

$$(3)$$

$$\Rightarrow T'^{11} = 4(x^1)^2 + 4(x^2)^2 \quad (4)$$

$$\Rightarrow T'^{12} = T'^{21} = 0 \quad (5)$$

$$\Rightarrow T'^{22} = 4(x^1)^2 + 4(x^2)^2 \quad (6)$$

The components in at the point $x^1 = 1, x^2 = 0$ are

$$T'(1, 0) = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}$$



1.22 p24-exercise 7.

Given that if $T_{mnr s}$ is a covariant tensor, and

$$T_{mnr s} + T_{mnsr} = 0$$

in a coordinate system x^p , establish directly that

$$T_{mnr s} + T_{mnsr} = 0$$

in any other coordinate system x, q .

Note: in the following, different alphabets in the indices indicates different coordinate systems.
As we $T_{mnr s}$ is a covariant tensor :

$$T_{\alpha\beta\gamma\delta} = T_{mnr s} \frac{\partial x^m}{\partial x^\alpha} \frac{\partial x^n}{\partial x^\beta} \frac{\partial x^r}{\partial x^\gamma} \frac{\partial x^s}{\partial x^\delta} \quad (1)$$

$$\Rightarrow T_{\alpha\beta\gamma\delta} + T_{\alpha\beta\delta\gamma} = T_{mnr s} \frac{\partial x^m}{\partial x^\alpha} \frac{\partial x^n}{\partial x^\beta} \frac{\partial x^r}{\partial x^\gamma} \frac{\partial x^s}{\partial x^\delta} + T_{mnr s} \frac{\partial x^m}{\partial x^\alpha} \frac{\partial x^n}{\partial x^\beta} \frac{\partial x^r}{\partial x^\delta} \frac{\partial x^s}{\partial x^\gamma} \quad (2)$$

Now, swap the dummy indices r and s in the second term on the right and as $T_{mnr s} = -T_{mnsr}$:

$$T_{\alpha\beta\gamma\delta} + T_{\alpha\beta\delta\gamma} = T_{mnr s} \frac{\partial x^m}{\partial x^\alpha} \frac{\partial x^n}{\partial x^\beta} \frac{\partial x^r}{\partial x^\gamma} \frac{\partial x^s}{\partial x^\delta} + T_{mnsr} \frac{\partial x^m}{\partial x^\alpha} \frac{\partial x^n}{\partial x^\beta} \frac{\partial x^s}{\partial x^\delta} \frac{\partial x^r}{\partial x^\gamma} \quad (3)$$

$$= (T_{mnr s} + T_{mnsr}) \frac{\partial x^m}{\partial x^\alpha} \frac{\partial x^n}{\partial x^\beta} \frac{\partial x^s}{\partial x^\delta} \frac{\partial x^r}{\partial x^\gamma} \quad (4)$$

$$= 0 \quad (5)$$



1.23 p24-exercise 8.

Prove that if A_r is a covariant vector, then $\frac{\partial A_r}{\partial x^s} - \frac{\partial A_s}{\partial x^r}$ is a skew-symmetric covariant tensor of the second order (use the notation of 1.7).

Be $B_{rs} = \frac{\partial A_r}{\partial x^s} - \frac{\partial A_s}{\partial x^r}$.

i) B_{rs} is skew-symmetric: It is obvious that:

$$-B_{rs} = -\frac{\partial A_r}{\partial x^s} + \frac{\partial A_s}{\partial x^r} = \frac{\partial A_s}{\partial x^r} - \frac{\partial A_r}{\partial x^s} \equiv B_{sr}$$

ii) B_{rs} is covariant:

Note: in the following, different alphabets in the indices indicates different coordinate systems.

Let

$$C_{\alpha\beta} = (\partial_s A_r - \partial_r A_s) X_\alpha^r X_\beta^s. \quad (1)$$

We know that $A_i = A_\gamma X_i^\gamma$ as A_i is covariant. Hence,

$$\partial_j A_i = \partial_j A_\gamma X_i^\gamma + A_\gamma \partial_j X_i^\gamma \quad (2)$$

$$= \partial_\alpha A_\gamma X_j^\alpha X_i^\gamma + A_\gamma \partial_j X_i^\gamma \quad (3)$$

Using (3), we compute the first term in (1)

$$\partial_s A_r X_\alpha^r X_\beta^s = \partial_\rho A_\gamma X_s^\rho X_r^\gamma X_\alpha^r X_\beta^s + A_\gamma \partial_s X_r^\gamma X_\alpha^r X_\beta^s \quad (4)$$

$$= \partial_\rho A_\gamma X_\beta^\rho X_\alpha^\gamma + A_\gamma \partial_s X_r^\gamma X_\alpha^r X_\beta^s \quad (5)$$

$$= \partial_\rho A_\gamma \delta_\beta^\rho \delta_\alpha^\gamma + A_\gamma \partial_s X_r^\gamma X_\alpha^r X_\beta^s \quad (6)$$

$$= \partial_\beta A_\alpha + A_\gamma \partial_s X_r^\gamma X_\alpha^r X_\beta^s \quad (7)$$

In the same way, we get for the second term in (1)

$$\partial_r A_s X_\alpha^s X_\beta^r = \partial_\alpha A_\beta + A_\gamma \partial_r X_s^\gamma X_\alpha^r X_\beta^s \quad (8)$$

And thus,

$$C_{\alpha\beta} = (\partial_s A_r - \partial_r A_s) X_\alpha^r X_\beta^s = \partial_\beta A_\alpha + A_\gamma \partial_s X_r^\gamma X_\alpha^r X_\beta^s - \partial_\alpha A_\beta - A_\gamma \partial_r X_s^\gamma X_\alpha^r X_\beta^s \quad (9)$$

$$\Rightarrow \partial_\beta A_\alpha - \partial_\alpha A_\beta = (\partial_s A_r - \partial_r A_s) X_\alpha^r X_\beta^s \quad (10)$$

So, i) and (10) proves that $\frac{\partial A_r}{\partial x^s} - \frac{\partial A_s}{\partial x^r}$ is skew-symmetric tensor of the second order.



1.24 p24-exercise 9.

Let $x^r, \bar{x}^r, y^r, \bar{y}^r$ be four systems of coordinates. Examine the tensor character of $\frac{\partial x^r}{\partial y^s}$ with respect to the following transformations:

- i) A transformation $x^r = f^r(\bar{x}^1, \dots, \bar{x}^N)$, with y^r unchanged;
- ii) A transformation $y^r = g^r(\bar{y}^1, \dots, \bar{y}^N)$, with x^r unchanged;

Note: in the following, different alphabets in the indices indicates different coordinate systems.

i) Let's compute the expression $A(\alpha, \beta) = \frac{\partial x^r}{\partial y^s} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^s}{\partial x^\beta}$. Obviously, the right side is an expression of a (possible) mixed tensor of the second order ($\frac{\partial x^r}{\partial y^s}$) under transformation from the (r) coordinate system to the (α) coordinate system. Then,

$$A(\alpha, \beta) = \frac{\partial x^r}{\partial y^s} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^s}{\partial x^\beta} \quad (1)$$

$$= \frac{\partial x^\alpha}{\partial y^s} \frac{\partial x^s}{\partial x^\beta} \quad (2)$$

$$= \frac{\partial x^\alpha}{\partial y^\rho} \frac{\partial y^\rho}{\partial y^s} \frac{\partial x^s}{\partial x^\beta} \quad (3)$$

If we consider the \bar{y}^r coordinate system as the y^ρ coordinate system and as $\bar{y}^r = y^r$ then $\frac{\partial y^\rho}{\partial y^s} = \delta_s^\rho$ and we get from (3)

$$A(\alpha, \beta) = \frac{\partial x^\alpha}{\partial y^\rho} \frac{\partial y^\rho}{\partial y^s} \frac{\partial x^s}{\partial x^\beta} \quad (4)$$

$$= \frac{\partial x^\alpha}{\partial y^\rho} \delta_s^\rho \frac{\partial x^s}{\partial x^\beta} \quad (5)$$

$$= \frac{\partial x^\alpha}{\partial y^\rho} \frac{\partial x^\rho}{\partial x^\beta} \quad (6)$$

$$= \frac{\partial x^\alpha}{\partial y^\rho} \delta_\beta^\rho \quad (7)$$

$$= \frac{\partial x^\alpha}{\partial y^\beta} \quad (8)$$

$$(1) \text{ and } (8) \Rightarrow \frac{\partial x^\alpha}{\partial y^\beta} = \frac{\partial x^r}{\partial y^s} \frac{\partial x^\alpha}{\partial x^r} \frac{\partial x^s}{\partial x^\beta} \quad (9)$$

So $A(r, s) = \frac{\partial x^r}{\partial y^s}$ is a mixed tensor of type A_s^r

ii) Let's compute the expression $A(\alpha, \beta) = \frac{\partial x^r}{\partial y^s} \frac{\partial y^\alpha}{\partial y^r} \frac{\partial y^s}{\partial y^\beta}$. Obviously, the right side is an expression of a (possible) mixed tensor of the second order ($\frac{\partial x^r}{\partial y^s}$) under transformation from the (r) coordinate

system to the (α) coordinate system. Then,

$$A(\alpha, \beta) = \frac{\partial x^r}{\partial y^s} \frac{\partial y^\alpha}{\partial y^r} \frac{\partial y^s}{\partial y^\beta} \quad (10)$$

$$= \frac{\partial x^r}{\partial y^\rho} \frac{\partial y^\rho}{\partial y^s} \frac{\partial y^\alpha}{\partial y^r} \frac{\partial y^s}{\partial y^\beta} \quad (11)$$

$$= \frac{\partial x^r}{\partial y^\rho} \frac{\partial y^\rho}{\partial y^\beta} \frac{\partial y^\alpha}{\partial y^r} \quad (12)$$

$$= \frac{\partial x^r}{\partial y^\rho} \delta_\beta^\rho \frac{\partial y^\alpha}{\partial y^r} \quad (13)$$

$$= \frac{\partial x^r}{\partial y^\beta} \frac{\partial y^\alpha}{\partial y^r} \quad (14)$$

$$= \frac{\partial x^r}{\partial x^\sigma} \frac{\partial x^\sigma}{\partial y^\beta} \frac{\partial y^\alpha}{\partial y^r} \quad (15)$$

If we consider the \bar{x}^r coordinate system as the x^σ coordinate system and as $\bar{x}^r = x^r$ then $\frac{\partial x^\sigma}{\partial x^r} = \delta_r^\sigma$ and we get from (15)

$$A(\alpha, \beta) = \frac{\partial x^r}{\partial x^\sigma} \frac{\partial x^\sigma}{\partial y^\beta} \frac{\partial y^\alpha}{\partial y^r} \quad (16)$$

$$= \delta_\sigma^r \frac{\partial x^\sigma}{\partial y^\beta} \frac{\partial y^\alpha}{\partial y^r} \quad (17)$$

$$= \frac{\partial x^\sigma}{\partial y^\beta} \frac{\partial y^\alpha}{\partial y^\sigma} \quad (18)$$

$$= \frac{\partial x^\sigma}{\partial y^\beta} \delta_\sigma^\alpha \quad (19)$$

$$= \frac{\partial x^\alpha}{\partial y^\beta} \quad (20)$$

$$(10) \text{ and } (19) \Rightarrow \frac{\partial x^\alpha}{\partial y^\beta} = \frac{\partial x^r}{\partial y^s} \frac{\partial y^\alpha}{\partial y^r} \frac{\partial y^s}{\partial y^\beta} \quad (21)$$

So $A(r, s) = \frac{\partial x^r}{\partial y^s}$ is a mixed tensor of type A_s^r



1.25 p23-exercise 10.

If x^r, y^r, z^r are three systems of coordinates, prove the following rule for the multiplication of Jacobians.

$$\left| \frac{\partial x^m}{\partial y^n} \right| \left| \frac{\partial y^r}{\partial z^s} \right| = \left| \frac{\partial x^t}{\partial z^u} \right|$$

As we have

$$\frac{\partial x^t}{\partial z^u} = \frac{\partial x^t}{\partial y^k} \frac{\partial y^k}{\partial z^u} \quad (1)$$

$$\begin{bmatrix} \frac{\partial x^1}{\partial z^1} & \cdots & \frac{\partial x^1}{\partial z^N} \\ \vdots & \vdots & \vdots \\ \frac{\partial x^N}{\partial z^1} & \cdots & \frac{\partial x^N}{\partial z^N} \end{bmatrix} = \begin{bmatrix} \frac{\partial x^1}{\partial y^k} \frac{\partial y^k}{\partial z^1} & \cdots & \frac{\partial x^1}{\partial y^k} \frac{\partial y^k}{\partial z^N} \\ \vdots & \vdots & \vdots \\ \frac{\partial x^N}{\partial y^k} \frac{\partial y^k}{\partial z^1} & \cdots & \frac{\partial x^N}{\partial y^k} \frac{\partial y^k}{\partial z^N} \end{bmatrix} \quad (2)$$

$$= \begin{bmatrix} \frac{\partial x^1}{\partial y^1} & \cdots & \frac{\partial x^1}{\partial y^N} \\ \vdots & \vdots & \vdots \\ \frac{\partial x^N}{\partial y^1} & \cdots & \frac{\partial x^N}{\partial y^N} \end{bmatrix} \begin{bmatrix} \frac{\partial y^1}{\partial z^1} & \cdots & \frac{\partial y^1}{\partial z^N} \\ \vdots & \vdots & \vdots \\ \frac{\partial y^N}{\partial z^1} & \cdots & \frac{\partial y^N}{\partial z^N} \end{bmatrix} \quad (3)$$

$$\Rightarrow \left| \frac{\partial x^m}{\partial y^n} \right| \left| \frac{\partial y^r}{\partial z^s} \right| = \left| \frac{\partial x^t}{\partial z^u} \right| \quad (4)$$



1.26 p23-exercise 11.

Prove that with respect to transformations

$$x^{,r} = C_{rs}x^s$$

where the coefficients are constants satisfying

$$C_{mr}C_{ms} = \delta_s^r$$

contravariant and covariant vectors have the same formula of transformation

$$A^{,r} = C_{rs}A^s, A_{,r} = C_{rs}A_s$$

i) $A^{,r} = C_{rs}A^s$

Be $A^{,r} = A^s \frac{\partial x^{,r}}{\partial x^s}$ and as $x^{,r} = C_{rs}x^s$ we have $\frac{\partial x^{,r}}{\partial x^s} = C_{rs}$. Hence,

$$A^{,r} = C_{rs}A^s$$

.

i) $A_{,r} = C_{rs}A_s$

Be $A_{,r} = A_s \frac{\partial x^s}{\partial x^{,r}}$ and as $x^{,r} = C_{rs}x^s$ we have

$$\frac{\partial x^{,r}}{\partial x^{,t}} = C_{rs} \frac{\partial x^s}{\partial x^{,t}} \quad (1)$$

$$\Rightarrow \delta_t^r = C_{rs} \frac{\partial x^s}{\partial x^{,t}} \quad (2)$$

Now, multiply (2) by C_{rq} . We get,

$$\delta_t^r C_{rq} = C_{rq} C_{rs} \frac{\partial x^s}{\partial x^{,t}} \quad (3)$$

$$C_{tq} = C_{rq} C_{rs} \frac{\partial x^s}{\partial x^{,t}} \quad (4)$$

$$\text{as } C_{mr}C_{ms} = \delta_s^r \Rightarrow C_{tq} = \delta_s^q \frac{\partial x^s}{\partial x^{,t}} \quad (5)$$

$$\Rightarrow C_{tq} = \frac{\partial x^q}{\partial x^{,t}} \text{ or } C_{rs} = \frac{\partial x^s}{\partial x^{,r}} \quad (6)$$

$$\text{as } A_{,r} = A_s \frac{\partial x^s}{\partial x^{,r}} \Rightarrow A_{,r} = C_{rs} \frac{\partial x^s}{\partial x^{,r}} \quad (7)$$



1.27 p23-exercise 12.

Prove that

$$\frac{\partial \ln \left| \frac{\partial x^m}{\partial y^n} \right|}{\partial x^r} = \frac{\partial^2 y^m}{\partial x^r \partial x^n} \frac{\partial x^n}{\partial y^m}$$

Lemma:

Be A a square matrix $N \times N$; Be f a C^1 function $f : \mathbb{R}^{N \times N} \rightarrow \mathbb{R}$.

Define A' as $(A')_{ij} = \frac{df}{dA_{ij}}$. Then,

$$(\ln |A|)' = (A^{-1})^T \text{ with } f = |A|$$

Proof:

By definition of the determinant, we have

$$|A| = A_{iK} C_K^i \quad (\text{no summation on } K!) \quad (1)$$

with $(C_K^i) = (-1)^{i+K} M_K^i$ being the cofactor of element A_{iK} and M_K^i the minor $(N-1) \times (N-1)$ matrix associated with the cofactor A_{iK} . Be $C = (C_{ij}^i)$ the $N \times N$ matrix formed with all possible cofactor elements C_j^i ($i, j = 1 \dots, N$).

We have

$$A^{-1} = \frac{C^T}{|A|} \quad (2)$$

$$\Rightarrow (A^{-1})^T = \frac{C}{|A|} \quad (3)$$

$$\text{differentiating (1)} \Rightarrow \frac{\partial |A|}{\partial A_{mn}} = \frac{\partial A_{iK}}{\partial A_{mn}} C_K^i + A_{iK} \frac{\partial C_K^i}{\partial A_{mn}} \quad (4)$$

$$\text{we have for } i = m \quad \begin{aligned} \frac{\partial A_{iK}}{\partial A_{mn}} &= 1 & K &= n \\ \frac{\partial A_{iK}}{\partial A_{mn}} &= 0 & K &\neq n \end{aligned} \quad (5)$$

Also, $\forall K : \frac{\partial C_K^i}{\partial A_{in}} = 0$ as by definition of the cofactor matrix, A_{ij} is not contained in C_{ij} .

Hence, (4) becomes

$$\frac{\partial |A|}{\partial A_{ij}} = C_j^i \quad (6)$$

$$\text{But, } \frac{\partial \ln |A|}{\partial A_{ij}} = \frac{\frac{\partial |A|}{\partial A_{ij}}}{|A|} \quad (7)$$

$$(6) \text{ and } (7) \text{ gives } \frac{\partial \ln |A|}{\partial A_{ij}} = \frac{C_j^i}{|A|} \quad (8)$$

$$(3) \text{ and } (8) \text{ gives } \frac{\partial \ln |A|}{\partial A_{ij}} = \frac{(A_{ij}^{-1})^T |A|}{|A|} = (A_{ij}^{-1})^T \quad (9)$$

$$\Rightarrow (\ln |A|)' = (A^{-1})^T \quad (10)$$

Now the main proof:

Let,

$$A \equiv [a_{mn}] = \left[\frac{\partial y^m}{\partial x^n} \right] \quad (11)$$

$$\Rightarrow \frac{\partial \ln |A|}{\partial x^r} = \frac{\partial \ln |A|}{\partial a_{mn}} \frac{\partial a_{mn}}{\partial x^r} \quad (12)$$

$$\text{from (10) we get } \frac{\partial \ln |A|}{\partial a_{mn}} = (A^{-1})_{mn}^T \quad (13)$$

$$\text{But } A \text{ is a Jacobian, so } (A^{-1})_{mn} = \frac{\partial x^m}{\partial y^n} \quad (14)$$

$$\text{and thus } (A^{-1})_{mn}^T = \frac{\partial x^n}{\partial y^m} \quad (15)$$

$$(13) \text{ can be written as } \frac{\partial \ln |A|}{\partial x^r} = \frac{\partial x^n}{\partial y^m} \frac{\partial a_{mn}}{\partial x^r} \quad (16)$$

$$\Rightarrow \frac{\partial \ln |A|}{\partial x^r} = \frac{\partial x^n}{\partial y^m} \frac{\partial^2 y^m}{\partial x^r \partial x^n} \quad (17)$$

◆

1.28 p23-exercise 13.

Consider the quantities $\frac{dx^r}{dt}$ for a particle moving in the plane. If x^r are the rectangular Cartesian coordinates, are these quantities the components of a contravariant or covariant vector with respect to rotation of the axes? Are they components of a vector with respect to transformation to any curvilinear coordinates (e.g. polar coordinates)?

Note: we suppose that by a rotation of the axes, the problem means a fixed rotation and not a rotation varying in time.

i) Be $v^r = \frac{dx^r}{dt}$ and consider v^α the same object but in another the coordinate system. A rotation of the axes implies the linear form

$$x^\alpha = R_k^\alpha x^k \quad \text{with } R_k^\alpha \neq R_k^\alpha(x^k) \quad (1)$$

$$\Rightarrow \frac{\partial x^\alpha}{\partial x^r} = R_k^\alpha \delta_r^k \quad (2)$$

$$\Rightarrow R_r^\alpha = \frac{\partial x^\alpha}{\partial x^r} \quad (3)$$

Consider $v^\alpha = \frac{dx^\alpha}{dt}$

$$v^\alpha = \frac{dx^\alpha}{dt} \quad (4)$$

$$(1) \Rightarrow v^\alpha = R_k^\alpha \frac{dx^k}{dt} \quad (5)$$

$$\Rightarrow v^\alpha = R_k^\alpha v^k \quad (6)$$

$$(3) \Rightarrow v^\alpha = v^k \frac{\partial x^\alpha}{\partial x^r} \quad (7)$$

Conclusion: v^k is a contravariant vector.

ii) Are they components of a vector with respect to transformation to any curvilinear coordinates (e.g. polar coordinates)?

We know that

$$dx^\alpha = \frac{\partial x^\alpha}{\partial x^r} dx^r \quad (8)$$

$$\Rightarrow \frac{dx^\alpha}{dt} = \frac{\partial x^\alpha}{\partial x^r} \frac{dx^r}{dt} \quad (9)$$

$$\Rightarrow v^\alpha = v^r \frac{\partial x^\alpha}{\partial x^r} \quad (10)$$

So v^r is a contravariant vector in general. Note that this proof is more straightforward than the prove in i).



1.29 p23-exercise 14.

Consider the question raised in No. 13 for the acceleration $\frac{d^2 x^r}{dt^2}$.

From exercise 13. we know that

$$\frac{dx^\alpha}{dt} = \frac{\partial x^\alpha}{\partial x^r} \frac{dx^r}{dt} \quad (1)$$

$$\Rightarrow \frac{d^2 x^\alpha}{dt^2} = \frac{d^2 x^r}{dt^2} \frac{\partial x^\alpha}{\partial x^r} + \frac{d \frac{\partial x^\alpha}{\partial x^r}}{dt} \frac{dx^r}{dt} \quad (2)$$

$$= \frac{d^2 x^r}{dt^2} \frac{\partial x^\alpha}{\partial x^r} + \frac{\partial^2 x^\alpha}{\partial x^r \partial x^m} \frac{dx^m}{dt} \frac{dx^r}{dt} \quad (3)$$

The second term on the right does not vanish in general, hence $\frac{d^2 x^r}{dt^2}$ has not a tensor character.



1.30 p23-exercise 15.

It is well known that the equation of an ellipse may be written

$$ax^2 + 2hxy + by^2 = 1$$

What is the tensor character of a, h, b with respect to transformation to any Cartesian coordinates (rectangular or oblique) in the plane?

Consider the transformation from a (w, z) coordinate system to a (x, y) coordinate system. For the considered type of transformation we have

$$\begin{aligned} x &= \alpha w + \beta z \\ y &= \gamma w + \delta z \end{aligned} \quad (1)$$

$$\text{consider} \quad \begin{aligned} ax^2 + 2hxy + by^2 &= 1 \\ pw^2 + 2q wz + rz^2 &= 1 \end{aligned} \quad (2)$$

the two representations of the same ellipse in the respective coordinate systems. Plugging (1) in (2):

$$a\alpha^2 w^2 + 2a\alpha\beta wz + \alpha\beta^2 z^2 + 2h\alpha\gamma w^2 + \beta\delta z^2 + 2h(\alpha\delta + \gamma\beta)wz + b\gamma^2 w^2 + 2b\gamma\delta wz + \delta^2 z^2 = 1 \quad (3)$$

$$(4)$$

Rearranging and equating the terms in w^2, wz, z^2 in (2) gives

$$p = a\alpha^2 + 2h\alpha\gamma + b\gamma^2 \quad (5)$$

$$q = a\alpha\beta + h(\alpha\delta + \gamma\beta) + \gamma\delta \quad (6)$$

$$r = a\beta^2 + 2h\beta\delta + b\delta^2 \quad (7)$$

Consider the following objects

$$(A_{ij}) = \begin{bmatrix} a & h \\ h & b \end{bmatrix} \quad (8)$$

$$(A_{ij})' = \begin{bmatrix} a'_{11} & a'_{12} \\ a'_{21} & a'_{22} \end{bmatrix} \quad (9)$$

$$\text{we calculate} \quad A'_{ij} = A_{km} \frac{\partial x^k}{\partial x'^i} \frac{\partial x^m}{\partial x'^j} \quad (10)$$

with $(x^1, x^2) = (w, z)$ and $(x'^1, x'^2) = (x, y)$. We have,

$$\frac{\partial x^1}{\partial x'^1} = \alpha, \frac{\partial x^1}{\partial x'^2} = \beta, \frac{\partial x^2}{\partial x'^1} = \gamma, \frac{\partial x^2}{\partial x'^2} = \delta \quad (11)$$

$$\begin{aligned} (10) \text{ and } (11) \quad \Rightarrow \quad \begin{aligned} a'_{11} &= a\alpha^2 + 2h\alpha\gamma + b\gamma^2 \\ a'_{22} &= a\beta^2 + 2h\beta\delta + b\delta^2 \\ a'_{12} &= a'_{21} = a\alpha\beta + h(\alpha\delta + \gamma\beta) + b\gamma\delta \end{aligned} \end{aligned} \quad (12)$$

Combining (5), (6), (7) and (12) we get

$$p = a'_{11}, r = a'_{22}, q = a'_{12} = a'_{21}$$

and so (9) becomes

$$(A_{ij})' = \begin{bmatrix} p & q \\ q & r \end{bmatrix}$$

Considering (10) we see that $\begin{bmatrix} a & h \\ h & b \end{bmatrix}$ transforms to $\begin{bmatrix} p & q \\ q & r \end{bmatrix}$ according to the rules of a covariant tensor of order two.



1.31 p23-exercise 16.

Matter is distributed in a plane and A, B, H are the moments and product of inertia with respect to rectangular axes Oxy in a plane. Examine the tensor character of the set of quantities A, B, H under rotation of the axes. What notation would you suggest for moments and product of inertia in order to exhibit the tensor character? What simple invariant can be formed from A, B, H ?

Consider the transformation from a (x^1, x^2) coordinate system to a (y^1, y^2) coordinate system. For the considered type of transformation we have

$$\begin{aligned} y^1 &= \alpha x^1 + \beta x^2 \\ y^2 &= \gamma x^1 + \delta x^2 \end{aligned} \quad (1)$$

$$\begin{aligned} \text{Be} \quad A &= \sum_{\rho} m_{\rho} (x^{2,\rho})^2 & A' &= \sum_{\rho} m_{\rho} (y^{2,\rho})^2 \\ B &= \sum_{\rho} m_{\rho} (x^{1,\rho})^2 & B' &= \sum_{\rho} m_{\rho} (y^{1,\rho})^2 \\ H &= \sum_{\rho} m_{\rho} x^{1,\rho} x^{2,\rho} & H' &= \sum_{\rho} m_{\rho} y^{1,\rho} y^{2,\rho} \end{aligned} \quad (2)$$

the moments and product of inertia, ρ being the index of summation over all the points with mass m_{ρ} .

For the sake of notational simplicity we consider only one point of mass as the linearity of A, B, H related to ρ ensures the validity of the next steps for all points in the plane.

From (1) and (2) we have:

$$\frac{A'}{m_{\rho}} = \gamma^2 (x^1)^2 + 2\gamma\delta x^1 x^2 + \delta^2 (x^2)^2 \quad (3)$$

$$\frac{B'}{m_{\rho}} = \alpha^2 (x^1)^2 + 2\alpha\beta x^1 x^2 + \beta^2 (x^2)^2 \quad (4)$$

$$\frac{H'}{m_{\rho}} = \alpha\gamma (x^1)^2 + (\gamma\beta + \alpha\delta) x^1 x^2 + \beta\delta (x^2)^2 \quad (5)$$

$$\begin{aligned} \text{Note that} \quad \frac{\partial y^1}{\partial x^1} &= \alpha & \frac{\partial y^1}{\partial x^2} &= \beta \\ \frac{\partial y^2}{\partial x^1} &= \gamma & \frac{\partial y^2}{\partial x^2} &= \delta \end{aligned} \quad (6)$$

$$(6) \text{ in } (4): \quad \frac{B'}{m_{\rho}} = (x^1)^2 \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^1} + 2(x^1)(x^2) \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + (x^2)^2 \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^2} \quad (7)$$

$$= (x^1)^2 \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^1} + (x^1)(x^2) \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + (x^2)(x^1) \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + (x^2)^2 \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^2} \quad (8)$$

Repeating the same calculations for $\frac{A'}{m_{\rho}}$ and $\frac{H'}{m_{\rho}}$ gives:

$$\begin{aligned} \frac{A'}{m_{\rho}} &= (x^1)^2 \frac{\partial y^2}{\partial x^1} \frac{\partial y^2}{\partial x^1} + (x^1)(x^2) \frac{\partial y^2}{\partial x^1} \frac{\partial y^2}{\partial x^2} + (x^2)(x^1) \frac{\partial y^2}{\partial x^2} \frac{\partial y^2}{\partial x^1} + (x^2)^2 \frac{\partial y^2}{\partial x^2} \frac{\partial y^2}{\partial x^2} \\ \frac{B'}{m_{\rho}} &= (x^1)^2 \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^1} + (x^1)(x^2) \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + (x^2)(x^1) \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + (x^2)^2 \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^2} \\ \frac{H'}{m_{\rho}} &= (x^1)^2 \frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^1} + (x^1)(x^2) \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^1} + (x^2)(x^1) \frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^2} + (x^2)^2 \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^2} \end{aligned} \quad (9)$$

Now, define

$$(K_{ij}) = \begin{bmatrix} (x^1)^2 & (x^1)(x^2) \\ (x^2)(x^1) & (x^2)^2 \end{bmatrix} \quad (K_{ij})' = \begin{bmatrix} (y^1)^2 & (y^1)(y^2) \\ (y^2)(y^1) & (y^2)^2 \end{bmatrix} \quad (10)$$

Then (9) can be written as

$$\begin{aligned} \frac{A'}{m_\rho} &= (y^1)^2 = K^{11} \frac{\partial y^2}{\partial x^1} \frac{\partial y^2}{\partial x^1} + K^{12} \frac{\partial y^2}{\partial x^1} \frac{\partial y^2}{\partial x^2} + K^{21} \frac{\partial y^2}{\partial x^2} \frac{\partial y^2}{\partial x^1} + K^{22} \frac{\partial y^2}{\partial x^2} \frac{\partial y^2}{\partial x^2} \\ \frac{B'}{m_\rho} &= (y^2)^2 = K^{11} \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^1} + K^{12} \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + K^{21} \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^1} + K^{22} \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^2} \end{aligned} \quad (11)$$

$$\frac{H'}{m_\rho} = (y^1)(y^2) = K^{11} \frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^1} + K^{12} \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^1} + K^{21} \frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^2} + K^{22} \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^2} \quad (12)$$

$$\begin{aligned} K^{,11} &= K^{11} \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^1} + K^{12} \frac{\partial y^1}{\partial x^1} \frac{\partial y^1}{\partial x^2} + K^{21} \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^1} + K^{22} \frac{\partial y^1}{\partial x^2} \frac{\partial y^1}{\partial x^2} \\ \text{or} \quad K^{,22} &= K^{11} \frac{\partial y^2}{\partial x^1} \frac{\partial y^2}{\partial x^1} + K^{12} \frac{\partial y^2}{\partial x^1} \frac{\partial y^2}{\partial x^2} + K^{21} \frac{\partial y^2}{\partial x^2} \frac{\partial y^2}{\partial x^1} + K^{22} \frac{\partial y^2}{\partial x^2} \frac{\partial y^2}{\partial x^2} \end{aligned} \quad (13)$$

$$K^{,12} = K^{,21} = K^{11} \frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^1} + K^{12} \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^1} + K^{21} \frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^2} + K^{22} \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^2}$$

So the object (K_{ij}) transforms according (13) like a contravariant second order tensor.

Now, consider $|K_{ij}|$, obviously $|K_{ij}| = 0$, but so is also $|K_{,ij}|$.

$\Rightarrow |K_{ij}|$ is an invariant under the considered transformation

